

Stochastic Expression of the Interferon- β Gene

Mingwei Zhao¹, Jiangwen Zhang^{2,3}, Hemali Phatnani^{3,9}, Stefanie Scheu⁴, Tom Maniatis^{1,3*}

1 Department of Molecular and Cellular Biology, Harvard University, Cambridge, Massachusetts, United States of America, **2** FAS Research Computing, Harvard University, Cambridge, Massachusetts, United States of America, **3** Columbia University College of Physicians and Surgeons, Department of Biochemistry and Molecular Biophysics, New York, New York, United States of America, **4** Institute of Medical Microbiology and Hospital Hygiene, Universität Düsseldorf, Düsseldorf, Germany

Abstract

Virus infection of mammalian cells induces the production of high levels of type I interferons (IFN α and β), cytokines that orchestrate antiviral innate and adaptive immunity. Previous studies have shown that only a fraction of the infected cells produce IFN. However, the mechanisms responsible for this stochastic expression are poorly understood. Here we report an in depth analysis of IFN-expressing and non-expressing mouse cells infected with Sendai virus. Mouse embryonic fibroblasts in which an internal ribosome entry site/yellow fluorescent protein gene was inserted downstream from the endogenous IFN β gene were used to distinguish between the two cell types, and they were isolated from each other using fluorescence-activated cell sorting methods. Analysis of the separated cells revealed that stochastic IFN β expression is a consequence of cell-to-cell variability in the levels and/or activities of limiting components at every level of the virus induction process, ranging from viral replication and expression, to the sensing of viral RNA by host factors, to activation of the signaling pathway, to the levels of activated transcription factors. We propose that this highly complex stochastic IFN β gene expression evolved to optimize both the level and distribution of type I IFNs in response to virus infection.

Citation: Zhao M, Zhang J, Phatnani H, Scheu S, Maniatis T (2012) Stochastic Expression of the Interferon- β Gene. *PLoS Biol* 10(1): e1001249. doi:10.1371/journal.pbio.1001249

Academic Editor: Hidde L. Ploegh, Whitehead Institute, United States of America

Received: July 23, 2011; **Accepted:** December 7, 2011; **Published:** January 24, 2012

Copyright: © 2012 Zhao et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work is supported by US National Institutes of Health grant 5R01AI020642-26 (to T.M.) (<http://grants.nih.gov/grants/oer.htm>) and Deutsche Forschungsgemeinschaft grant SCHE692/3-1 (to S.S.) (http://www.dfg.de/en/research_funding/programmes/index.html). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abbreviations: DI, defective interfering; dsRNA, double-stranded RNA; FACS, fluorescence-activated cell sorting; h.p.i., hours post-infection; HAU, hemagglutination units; IFN, interferon; ISG, interferon-stimulated gene; ISH, in situ hybridization; MEF, mouse embryonic fibroblast; NP, nucleoprotein; poly I:C, polyribonucleosinic polyribocytidylic acid; qPCR, quantitative PCR; qRT-PCR, quantitative reverse transcription PCR; SeV, Sendai virus

* E-mail: tm2472@mail.cumc.columbia.edu

⁹ These authors contributed equally to this work.

Introduction

Eukaryotic cells respond to extracellular signals and environmental stresses by coordinately activating specific sets of genes. Signals from the cell surface or cytoplasm trigger signaling pathways that culminate in the binding of distinct combinations of coordinately activated transcription factors to promoter and enhancer elements that regulate gene expression. A well-characterized example of this is the activation of type I interferon (IFN) gene expression in response to virus infection or double-stranded RNA (dsRNA) treatment [1,2]. After infection, viral RNA is detected in the cytoplasm by one of two RNA helicases, retinoic acid-inducible gene I (RIG-I) or melanoma differentiation-associated gene 5 (MDA5), which respond to different types of viruses [3]. RIG-I recognizes short dsRNA or panhandle RNA bearing a 5' triphosphate group [3], and its activity is positively regulated by the ubiquitin E3 ligase tripartite motif 25 (Trim25) [4]. When RIG-I or MDA5 bind to RNA, they form heterodimers, undergo a conformational change, and expose a critical N-terminal caspase-recruiting domain (CARD) [5,6]. This domain interacts with the CARD domain of the downstream adaptor protein mitochondrial antiviral signaling (MAVS) (also known as IPS-1/Cardif/VISA) on the mitochondrial membrane [7]. The association of RIG-I with MAVS initiates the recruitment of adaptor proteins and leads to the activation of the transcription

factors IFN regulatory factors 3 and 7 (IRF3 and IRF7) and NF- κ B by the TANK-binding kinase 1 (TBK1) [8–10] and IKK α and IKK β , respectively [7,11]. Activated IRF3/IRF7 and NF- κ B translocate into the nucleus and, along with the transcription factors ATF2/cJun, bind the IFN- β gene enhancer and recruit additional transcription components to form an enhanceosome [12]. This complex signaling and promoter recognition mechanism functions to coordinately activate a specific set of transcription factors that recognize the unique enhancer sequence of the IFN β gene and thus specifically activate IFN gene expression.

Early in situ hybridization (ISH) studies revealed that induction of IFN β expression by virus infection or dsRNA treatment in both human and mouse cells is stochastic [13,14]. That is, only a fraction of the infected cells express IFN β . This “noisy” expression is not due to genetic variation within the cell population, as multiple subclones of individual cells display the same low percentage of cells expressing IFN β [14]. In addition, different mouse and human cell lines display different percentages of expressed cells, and the levels of IFN β gene expression can be increased in low expressing cell lines by fusing them with high expressing lines, or by treating low expressing lines with IFN β [13,14]. These studies suggest that stochastic expression of the IFN β gene is a consequence of cell-to-cell differences in limiting cellular components required for IFN induction, and that one or more of the limiting factors are inducible by IFN β [13].

Author Summary

Eukaryotic cells can respond to extracellular signals by triggering the activation of specific genes. Viral infection of mammalian cells, for example, induces a high level of expression of type I interferons (IFN α and β), proteins required for antiviral immunity that protects cells from the infection. Previous studies have shown that the expression of the IFN β gene is stochastic, and under optimal conditions only a fraction of the infected cells express the IFN β gene. At present neither the mechanisms nor functions of this interesting phenomenon are well understood. We have addressed this question by analyzing IFN-expressing and non-expressing mouse cells that were infected with the highly transmissible Sendai virus. We show that stochastic IFN β gene expression is a consequence of cell-to-cell differences in limiting levels and/or activities of virus components at every level of the virus induction process, from viral replication to expression. These differences include the sensing of viral RNA by host factors, the activation of the signaling pathway, and the levels of activated transcription factors. Our findings reveal the complexity of the regulatory mechanisms controlling stochastic IFN β gene expression. We propose that the stochastic expression of IFN allows for an even distribution of IFN, thus avoiding over-expression of IFN in infected cells.

Stochastic expression has been observed with a number of other cytokine genes, including *IL-2* [15], *IL-4* [16,17], *IL-10* [18], *IL-5*, and *IL-13* [19]. In many of these cases, expression is both stochastic and monoallelic. Recent studies of IFN β gene expression revealed that stochastic expression in human cells is initially monoallelic, and becomes biallelic later in the induction [20,21]. In one study the stochastic expression of the IFN β gene was proposed to be a consequence of intrinsic noise due to stochastic enhanceosome assembly [21]. Subsequently, an analysis of human HeLa cells identified a specific set of Alu-repetitive DNA sequences bearing NF- κ B binding sites that associate with the IFN β gene through interchromosomal interactions, and in so doing are thought to increase the local concentration of NF- κ B. Initially, only one of the two chromosomes associates with the specialized NF- κ B binding sequence, resulting in early monoallelic expression. Secretion of IFN leads to an increased expression of limiting factors (most likely IRF7, which is inducible by IFN), obviating the need for interchromosomal interactions, and leading to the activation of the second IFN β allele [20]. More recently, heterogeneity in the infecting viruses, rather than cell cycle differences, has been proposed to be the primary source of IFN stochastic expression [22]. Many functions have been proposed for biological noise, ranging from cell fate decisions during development to survival in fluctuating environments [23]. In the case of the IFN genes, neither the mechanisms nor functions of biological noise are well understood.

Here we report a detailed analysis of stochastic IFN β gene expression in mouse cells. We make use of an IFN-IRES-YFP reporter mouse [24] to perform a detailed analysis of differences between virus-infected cells that either express or do not express IFN β . Our results reveal a complex picture of stochastic expression of the IFN β gene, in which the levels of components required for virtually every step in the virus induction pathway are limiting. This includes components required for viral replication and expression, for sensing the presence of viral RNA by the host, and for the virus induction signaling pathway, and the transcription factors required of IFN β gene expression. Remarkably, in

spite of this complexity the percentage of expressing cells remains constant through recloning and cell division, indicating that the stochasm of clonal cells is genetically programmed.

Results

Stochastic Expression of Mouse and Human IFN β Genes

Sendai virus (SeV) infection of either mouse or human cells leads to the expression of IFN β mRNA in only a fraction of the infected cells (Figures 1A, 1B, and S1A), and the percentage of expressing cells differs between different cell lines. The time course of mouse IFN β expression determined by ISH (Figure 1B) is consistent with that from the quantitative PCR (qPCR) analysis (Figure S1B and S1C). Remarkably, the percentage of cells expressing IFN did not exceed 20%, even at the latest time point (Figure 1B). The absence of IFN β signal in the majority of cells is not an artifact of hybridization, as β -actin mRNA was detected in all cells (Figure S1D). IFN β mRNA is specifically detected with an antisense IFN β RNA probe, while no signal is detected with a sense RNA probe (Figure S1E). In addition, similar percentages of IFN β -expressing cells were detected by immunofluorescent staining using an IFN β antibody (Figure S1F), strongly supporting the reproducibility and specificity of the IFN β ISH.

As mentioned above, enhanceosome assembly and limiting amounts of NF- κ B have been proposed to be the primary limiting steps in stochastic expression of the human IFN β gene [20,21]. To determine whether this stochastic expression is unique to the IFN β gene because of the complexity of the IFN β enhanceosome, or is more general, we examined the expression of the IFN α genes, which are coinduced with IFN β , but have simple enhancer/promoters, and do not require NF- κ B [25,26]. Using either a mouse IFN α 4 or human IFN α 8 probe, we found that IFN α genes are also stochastically expressed in both mouse and human cells, respectively (Figures 1C and S1G). Although NF- κ B has been shown to be a limiting factor in the activation of the human IFN β gene [20], it is not required for IFN β expression in mouse cells [27]. Thus, in spite of this difference both the mouse and human IFN β genes are stochastically expressed. We also examined other virus-inducible genes, and found that they too are stochastically expressed (see below). Each of these virus-inducible genes requires different levels and combinations of transcription factors, yet they are all stochastic. In all of these cases (mouse and human IFN β and IFN α and the other virus-inducible genes), the common requirement is the RIG-I virus-inducible signaling pathway. We therefore carried out experiments to determine whether limiting components in this pathway contribute to the observed stochastic expression.

Separation and Characterization of IFN-Expressing and Non-Expressing Cells

To investigate the mechanism of stochastic IFN β gene expression, we made use of an IFN β reporter-knock-in mouse, in which YFP expression allows tracking of IFN β expression at a single-cell level [24]. Using IFN β /YFP homozygous mouse embryonic fibroblasts (MEFs) and fluorescence-activated cell sorting (FACS), we obtained pure populations of IFN β -producing and IFN β -negative cells upon SeV infection. As expected, IFN β mRNA is high in the YFP-positive cells, and very low in the YFP-negative cells (Figure S2A). As expected, the IFN α 2 and IFN α 4 genes are also highly expressed in the YFP-positive cells, and not in the YFP-negative cells (Figure S2A). These observations indicate that replication of the infecting virus and/or components in the RIG-I pathway are the limiting steps in the uninduced cells, rather than intrinsic differences in the IFN β and α promoters.




Figure 1. Stochastic IFN and virus-inducible gene expression. (A) Stochastic IFN β gene expression detected by ISH using a digoxigenin-labeled IFN β RNA probe. (B) Percentage of IFN β -producing cells at different times after SeV infection. (C) Mouse IFN α gene expression in primary MEFs detected by ISH using a digoxigenin-labeled IFN α 4 probe. (D) qPCR analysis illustrating the expression levels of different virus-inducible genes in sorted IFN β /YFP MEFs. doi:10.1371/journal.pbio.1001249.g001

We also detected the relative mRNA abundance of other virus-inducible genes in IFN β -expressing and non-expressing cells. As shown in Figure 1D, transcription levels of all tested inflammatory cytokine or chemokine genes are much higher in IFN β -producing cells compared to nonproducers. Considering the fact that IFN β -producing cells account for only 10% of the total cell population, we conclude that expression of all these virus-inducible genes is also stochastic and that these genes are coordinately activated with the type I IFN genes. Activation of these virus-inducible genes is known to require the RIG-I signaling pathway [28–31]. Thus, our results indicate that stochastic gene expression is due primarily to limiting components in the signaling pathway and not to gene-to-gene variation in the mechanism of gene activation.

In the case of human cells, stochastic expression of the IFN β gene is randomly monoallelic early and biallelic late in infection, and the activation of the second IFN β allele is inducible by IFN [20,21]. However, the nature of allelic expression of the IFN β gene has not been addressed in mouse cells. By using IFN β /YFP heterozygous MEFs, we showed that early after infection (<8 h

post-infection [h.p.i.]), IFN β gene expression was primarily monoallelic, while late in infection (8–16 h.p.i.), the majority of IFN β -expressing cells were both IFN β and YFP double-positive cells indicating that, as with human cells, a switch to biallelic expression also occurs in mouse cells (Figure S2B).

Previous studies have shown that the levels of IFN β gene expression can be increased by priming the cells with IFN β [13]. Using both mouse and human primary fibroblasts, we showed that IFN β pretreatment also increases the percentages of IFN β -expressing cells (Figure S3), indicating that the limiting factor(s) contributing to stochastic IFN β gene expression are, indeed, inducible by IFN β . One of these IFN-inducible factors is IRF7 ([20] and see below).

Viral Replication Is More Efficient in IFN β -Producing Cells

To examine the role of the infecting virus in stochastic IFN β gene expression, we infected primary MEFs with SeV followed by immunofluorescent staining using a SeV antibody. As shown in Figure S4A, most, if not all, of the cells are uniformly infected by

SeV, far more than could explain the small percentage of cells expressing IFN β gene. When we used increasing multiplicities of SeV (as defined by hemagglutination units [HAU]) to infect primary MEFs, we found that the percentage of IFN β -producing cells increased as the HAU was increased, reaching a maximum of approximately 18% at the peak (Figure S4B). However, as more virus was added (>200 HAU), the percentage of IFN β -producing cells decreased. Thus, the viral titer is not a limiting factor in the observed stochastic IFN β gene expression. Next, we determined viral transcript levels in both IFN β -producing and nonproducing cells. We found that the nucleoprotein (NP), matrix protein, and L polymerase protein mRNA transcripts were present at significantly

higher levels in IFN β -producing cells compared to the nonproducers (Figures 2A and S4C). In addition, higher levels of SeV NP protein were detected in IFN β -producing cells (Figure 2B).

The RNA helicase RIG-I detects viral genomic RNA and defective interfering (DI) genomes [32,33]. We therefore examined the levels of viral and DI genomes in both IFN β -producing and nonproducing cells. As shown in Figure 2C (upper panel), more SeV DI genomes were detected in IFN β -producing cells compared to IFN β -nonproducing cells at 8 and 12 h.p.i. Using a primer pair that specifically detects viral genomic RNA, we also detected more viral genomes in IFN β -producing MEFs 8 and 12 h.p.i. (Figure 2C, lower panel). These results are consistent with the




Figure 2. Viral transcription and/or replication are more efficient in IFN β -producing cells. (A) qPCR analysis illustrating the relative abundance of viral NP, matrix (M), and L polymerase protein (L) mRNA in sorted IFN β /YFP MEFs. (B) Western blots showing cytoplasmic distribution of SeV NP protein present in IFN β -producing and nonproducing cells. (C) qPCR analysis illustrating the relative abundance of SeV DI genome (upper panel), and semi-qRT-PCR analysis illustrating the relative abundance of SeV genomic RNA (lower panel) in sorted IFN β /YFP MEFs. Reverse transcriptase PCR was carried out to detect viral genomic RNA and host cell β -actin mRNA (control) using gene-specific primers. After 35 cycles (SeV genomic RNA) or 26 cycles (β -actin) of amplification, PCR products were run on a 2% agarose gel. (D) Intracellular staining using SeV antibody and FACS analysis were carried out to determine the correlation between SeV infection and IFN β expression in IFN β /YFP homozygous MEFs. IB, immunoblot.

doi:10.1371/journal.pbio.1001249.g002

observed viral NP mRNA levels (Figure S4C), and indicate that viral replication is more efficient in the IFN-producing cells. We also investigated the induction activities of total RNA extracted from both IFN β -producing and nonproducing cells. As shown in Figure S4D, total RNA from IFN β -producing cells infected for 8 or 12 h induced more IFN β expression compared to total RNA from IFN β -nonproducers at the same time points. We conclude that viral mRNA, DI genomes, and viral genomes are present at higher levels in IFN β -producing cells than in nonproducers. Thus, differences in the efficiency of viral replication/transcription contribute to the stochastic expression of the IFN β gene.

Previous studies led to the conclusion that the stochastic expression of the IFN β gene is a feature of the infecting virus, and not of the host cell [22]. To address this possibility, we determined the number of cells that have high levels of viral RNA and produce IFN β at 8 h.p.i. As shown in Figure 2D, after 8 h of virus infection, approximately 38% SeV-high cells (upper left and upper right) were detected, and about 9% YFP-positive cells (upper right and lower right). Although a higher percentage of IFN β -expressing cells was observed within the SeV-high cell population (6.56% versus 2.42%), only 17% (6.56% out of 38%) of SeV-high cells produce IFN. Thus, although cell-to-cell differences in viral replication contribute to the stochastic expression of IFN, these differences are not sufficient to explain the extent of stochastic IFN gene expression.

The RIG-I Signaling Pathway Is Activated and More Potent in IFN β -Producing Cells

To further investigate the mechanism of stochastic IFN β gene expression, we determined the localization of various components of the signaling pathway required for IFN production using nuclear and cytoplasmic fractions separated from both expressing and non-expressing cells. Consistent with the limiting component hypothesis, we detected phosphorylation and translocation of IRF3 in the YFP-positive cells, but not in the YFP-negative cells (Figure 3A). Previous studies have shown that IRF3, like IRF7, is phosphorylated by the TBK1 kinase, and translocates from the cytoplasm to the nucleus. As both IRF3 and IRF7 are activated via the RIG-I pathway, our results suggest that one or more components of the RIG-I signaling pathway are limiting in the cells that fail to express IFN. A similar result was obtained with sorted cells at 12 h.p.i. (Figure 3B).

In human cells both NF- κ B and IRF3/IRF7 are required for virus induction of the IFN β gene [12,34]. The human and mouse IFN β enhancers differ in only two nucleotides out of 45 bases. However, in mouse cells NF- κ B is required only for early antiviral activity, when the level of active IRF3 is low, but is not required for maximum levels of IFN β expression late in induction [27,35]. Consistent with this finding, we show that only a small fraction of the p65 subunit of NF- κ B translocates to the nucleus 8 h.p.i., and little difference is observed in NF- κ B localization between the YFP-positive and YFP-negative cells (Figure 3A).

The observation that IRF3 activation and translocation occurs in only a fraction of virus-infected cells suggests that upstream components in the RIG-I signaling pathway differ in IFN β -producing and nonproducing cells. Western blotting results (Figure 3C) showed that IFN β -producing cells have higher levels of both RIG-I and MDA5 than the nonproducing population. Trim25, an E3 ligase required for RIG-I activation [4], is also present at a higher level in the IFN β -producing cells (Figure 3C). The increase in protein levels appears to be a consequence of differential transcription of the tested genes, as mRNA levels of all three genes are higher in IFN β -producing cells (Figure 3D). We conclude that the IFN β -producing cells have higher levels of

essential RIG-I signaling pathway components than the IFN β -nonproducing cells. Thus, at least part of the observed stochastic expression is due to limiting RIG-I pathway components in the cells that do not express IFN.

By contrast to the RNA detectors, the protein levels for both MAVS and TBK1, two essential components of the RIG-I signaling pathway [7,9], were lower in the IFN β -producing cells (Figure 3C). However, this is likely due to the degradation and/or cleavage of the MAVS protein in infected cells [36–38]. The data of Figure 3C suggest that TBK1 is also targeted for degradation during virus infection, consistent with the observation that TBK1 is subject to proteasome-dependent degradation [39]. Thus the turnover of both MAVS and TBK1 may be required for the post-induction turn-off of IFN β gene expression [38].

Over-Expression of Individual Components of the RIG-I Signaling Pathway Increases the Percentage of Cells Expressing IFN β

We have shown that the RIG-I signaling pathway is selectively activated in IFN β -expressing cells, and this is due only in part to the cell-to-cell differences in virus infection/replication. Our results also suggest that IFN β -producing cells have a more potent signaling pathway than IFN β -non-expressing cells. To further explore this possibility, we established a series of L929 stable cell lines that express RIG-I, MDA5, or Trim25 under the control of a tetracycline-inducible promoter (Figure S5A). As shown in Figure S5B and S5C, high levels of exogenous RIG-I only slightly increased the percentage of IFN β -producing cells. A larger increase was observed with MDA5 and Trim25, but the final percentage in both cases was still under 30%. Thus, these upstream components appear to be among several limiting factors in the cell population.

Additional components in the RIG-I signaling pathway were tested using the same approach, and high percentages of IFN β -producing cells were observed (Figure 4A and 4B). While a large difference between tetracycline-negative and tetracycline-positive cells was observed with the TBK1 line, only a small difference was observed between the corresponding MAVS lines. However, a large difference was observed between the non-transformed and transformed MAVS lines, suggesting that a low level of leaky transcription in the MAVS line is sufficient to dramatically increase the number of IFN β -expressing cells. These data clearly indicate that both MAVS and TBK1 are limiting components in the RIG-I pathway and therefore contribute significantly to stochastic IFN β expression.

We have shown that over-expression of RIG-I or Trim25 alone only slightly increases the percentage of IFN β -producing cells, but it is possible that both must be expressed to achieve maximum levels of IFN β production. We therefore transfected RIG-I stable transfectants with a Trim25 expression plasmid, and the other way around. The cells were then induced with tetracycline, infected with SeV, and examined for IFN β mRNA expression. Control experiments using a GFP reporter indicated that under our experimental conditions approximately 70% of cells can be transfected with the second plasmid (Figure S5D). As shown in Figure 4C and 4D, a dramatic increase was observed only 6 h.p.i. when either the RIG-I or Trim25 lines were transfected with Trim25 or RIG-I, respectively. This observation was confirmed by carrying out intracellular staining and flow cytometry experiments using IFN β /YFP homozygous MEFs (Figure S6). We conclude that the combination of RIG-I and Trim25 is limiting in the RIG-I pathway.

We note that the increase of IFN β -expressing cells was not observed in uninfected cells, with the only exception being MAVS.




Figure 3. The RIG-I signaling pathway is activated in IFN β -producing cells. (A and B) Western blots showing cytoplasmic (C) versus nuclear (N) distribution of different factors present in FACS-sorted cells 8 h.p.i. (A) and 12 h.p.i. (B). (C) Western blots showing cytoplasmic distribution of signaling pathway proteins present in FACS-sorted cells 8 h.p.i. Arrows indicate MAVS protein. (D) qPCR analysis illustrating the expression levels of RIG-I, Trim25, and MDA5 genes in sorted IFN β /YFP homozygous MEF cells 8 h.p.i. IB, immunoblot. doi:10.1371/journal.pbio.1001249.g003


Thus, over-expression of these signaling components did not bypass the requirement for signaling pathway activation.

IRF7 Is a Primary Limiting Factor in Stochastic IFN Gene Expression


Expression of the IFN β gene requires an active RIG-I signaling pathway and assembly of the enhanceosome complex on the IFN β

promoter. To investigate whether individual enhanceosome components are limiting factors, we established a series of tetracycline-inducible L929 stable lines that express IRF3, IRF7, or p65 genes. Figure 5A and 5B show that, without tetracycline induction, only 10%–15% of the cells produce detectable levels of IFN β mRNA in response to virus infection. Remarkably, the percentage of IFN β -producing cells upon SeV infection increased


A



B



D



C




Figure 4. Limiting factors in stochastic IFN β gene expression. (A) Different L929 stable transfectants were induced by tetracycline (Tet) for 24 h, followed by SeV infection for 9 h. RNA ISH experiments were carried out to detect the IFN β mRNA. (B and D) Histograms showing the percentage (mean \pm standard deviation) of cells expressing IFN β from three independent ISH experiments. At least 400 cells were blindly counted and scored for each category. (C) L929 stable transfectant was transiently transfected with expression vectors encoding either *GFP* (control), *RIG-I*, or *Trim25*, then stimulated with tetracycline for 24 h. Cells were then infected with SeV for 6 h, followed by RNA ISH to detect IFN β mRNA. pt, pt-REX-DEST30.

doi:10.1371/journal.pbio.1001249.g004

to 85% when IRF7 expression was induced by tetracycline in every cell (Figure S7A). A smaller increase (55%) was observed when IRF3 was over-expressed, whereas increasing the concentration of NF- κ B had little effect, consistent with the data in Figure 3A, and previously published studies [27]. Interestingly, IRF7 over-expression also significantly increased the percentage of IFN α -producing cells after virus infection (Figure S7B and S7C). It is known that IRF7 is required for maximum induction of type I IFN genes [25], and its basal protein level is very low in most cell types except for plasmacytoid dendritic cells [26,40]. We conclude that IRF7 is a critical limiting factor that is a major contributor to stochastic expression of mouse IFN α and β genes. This conclusion is also supported by our ISH results from 4E-BP1/4E-BP2 double-knockout MEFs (Figure 5C and 5D). Previous studies have identified 4E-BPs as negative regulators of type I IFN production via translational repression of IRF7 mRNA [41]. As shown in Figure 5C and 5D, we observed a 4-fold increase of the percentage of IFN β -expressing cells in 4E-BP1/4E-BP2 double-knockout MEFs compared to wild-type MEFs, consistent with the conclusion that a limiting amount of IRF7 is a major contributor to the stochastic expression of IFN β .

We also found that type I IFN induction was exceptionally high, with much faster kinetics in cells expressing exogenous IRF7 than in control cells (Figure S7D). In the absence of tetracycline induction, low levels of IFN β , IFN α 4, and IFN α 2 mRNA were first detected 6 h, 9 h, and 12 h.p.i., respectively. When the cells were treated with tetracycline, the kinetics of IFN gene transcription changed significantly. IFN β , IFN α 4, and IFN α 2 transcripts could be detected as early as 4 h after virus infection. Even at 24 h.p.i., steady and robust transcription of these genes could still be detected. These observations are consistent with a model in which IRF3 is normally activated early for IFN gene induction. Later, higher levels of IRF7 are produced by IFN and are required for both IFN β and IFN α gene expression, but IRF7 is rapidly turned over, leading to the cessation of both IFN β and IFN α gene expression [1,25,26]. By contrast, in the presence of excess IRF7 in the tetracycline-activated cells, both IFN β and IFN α are activated earlier, and continue to be expressed because of the continuous presence of IRF7.

IRF7 Positively Regulates the RIG-I Signaling Pathway

We have shown that over-expression of IRF7 or both RIG-I and Trim25 almost completely eliminates stochastic IFN β expression (Figures 4C, 4D, and 5). To investigate the connection between these observations, we carried out microarray analysis to compare genome-wide expression profiles of L929-IRF7 stable transfectants treated with or without tetracycline. Interestingly, upon IRF7 over-expression, only two up-regulated signaling pathways were identified from the KEGG Pathway Database, and the RIG-I-like receptor signaling pathway is the most up-regulated ($p = 3.6E-06$) (Figure S8A and S8B) [42]. We did not identify signaling pathways that were similarly enriched among the down-regulated genes. Using qPCR, we confirmed that the mRNA levels of both RIG-I and Trim25 were higher in IRF7 over-expressing cells (Figure S8C). Considering the low basal expression level of IRF7, we conclude that a high level of IRF7

protein increases the percentage of IFN β -expressing cells not only by increasing its own abundance, but also by up-regulating the RIG-I signaling pathway to increase the potency of activation of the IFN β gene.


Stochastic Expression of IFN β Induced by dsRNA—poly I:C Is Due to Limiting Amounts of MDA5 and IRF7

IFN β gene expression can also be induced by transfection of the synthetic dsRNA polyriboinosinic polyribocytidylic acid (poly I:C), and this induction occurs mainly through the MDA5 signaling pathway [43]. Early studies revealed that induction of IFN β expression by dsRNA treatment is also stochastic [13,14]. We therefore asked whether stochastic IFN β gene expression induced by dsRNA is due to cell-to-cell variation in the levels of MDA5 and IRF7. Using FACS analysis, we found that poly I:C-induced IFN β expression is also stochastic (Figure 6A). When IFN β /YFP homozygous MEFs were electroporated with Cy5-labeled poly I:C, only 9% of the cells produced IFN β as detected by the presence of YFP. However, the electroporation efficiency was over 99% (Figure 6A, left panel). Interestingly, based on the Cy5 intensity, there were two populations of cells, which contained different amounts of poly I:C. When we gated these two populations out as “poly I:C-high” and “poly I:C-low”, we observed that the “poly I:C-high” population included more cells producing IFN β (Figure 6A, right panel), indicating that the amount of inducer does affect the extent of stochastic IFN β expression. However, only a small percentage of “poly I:C-high” cells expressed the IFN β gene, clearly indicating that other limiting factor(s) dominate the stochastic IFN β expression induced by poly I:C transfection. We therefore carried out experiments to identify these limiting components.


L929-MDA5 and L929-RIG-I stable transfectants were transfected with poly I:C followed by ISH to detect IFN β expression. As shown in Figure 6B and 6C, over-expression of RIG-I only slightly increased the percentage of IFN β -producing cells. By contrast over-expression of MDA5, the major cytoplasmic receptor for poly I:C, led to a substantial increase in the percentage of IFN β -producing cells (from 15% to 65%). Considering that the transfection efficiency is approximately 75% (data not shown), over-expression of MDA5 basically eliminates stochastic expression of the IFN β gene in response to poly I:C transfection. Furthermore, the results of the flow cytometry experiment also supported this conclusion. As shown in Figure 6D, after 8 h of poly I:C stimulation, we observed approximately 2.6% YFP-positive cells. Within this population, about 70% of the YFP-positive cells had higher levels of MDA5 protein (1.86% out of 2.67%). We note that the percentage of YFP-positive cells is much lower than that observed with virus infection (Figures 2D and S6).

Over-expression of IRF3 or IRF7 also increased the percentage of IFN β -producing cells in response to poly I:C (Figure S9A and S9B). As shown in Figure S8, over-expression of the IRF7 gene up-regulates MDA5 gene expression. Considering its low basal expression level, IRF7 is also an important limiting factor in stochastic IFN β expression induced by poly I:C transfection. Taken together, these data show that poly I:C-induced stochastic IFN β expression depends on the abundance of both poly I:C and


A



B



D



C




Figure 5. IRF7 is the significant limiting factor in stochastic type I IFN gene expression. (A) Different L929 stable transfectants were induced by tetracycline (Tet) for 24 h, followed by SeV infection for 9 h. RNA ISH experiments were carried out to detect IFN β mRNA. (B and D) Histograms showing the percentage (mean \pm standard deviation) of cells expressing IFN β from three independent ISH experiments. At least 400 cells were blindly counted and scored for each category. (C) RNA ISH experiments were carried out to detect IFN β mRNA in wild-type (W.T.) or 4E-BP double-knockout (DKO) MEFs infected by SeV for 9 h.
doi:10.1371/journal.pbio.1001249.g005

signaling pathway protein MDA5 as well as IRF3/IRF7, which is similar to what was found in the case of virus infection.

Variation in the Levels of RIG-I Signaling Pathway Components

We also asked whether the concentrations of proteins regulating IFN β expression are sufficiently different from cell to cell to account for the stochastic IFN β expression. Using flow cytometry, we measured the distributions of six components in the RIG-I signaling pathway for which specific antibodies are available. As shown in Figure 7A and 7B, all six proteins were log-normally distributed across the population. Quantitative immunofluorescence data for individual components show similar distributions of each factor at the single-cell level (Figure S10). Combined with our previous data, these observations suggest that naturally occurring differences in the protein levels of signaling pathway components are the primary cause of cell-to-cell variability in IFN β expression upon virus infection.

IFN-Inducible Antiviral Genes Are Not Stochastically Expressed

When IFN is secreted from virus-infected cells *in vivo*, it binds to type I IFN receptors on surrounding cells and activates a large set of genes encoding antiviral proteins (interferon-stimulated genes [ISGs]) via the Jak/STAT signal transduction pathway. We therefore carried out experiments to determine whether the induction of antiviral ISGs is also stochastic. As shown in Figure 7C, ISG15 is expressed in all cells upon treatment with IFN β . Thus, when IFN is secreted, all of the surrounding cells produce antiviral proteins. This result is also consistent with previous observations showing that the antiviral response induced by IFN is a robust feature common to all cells, and is independent of the stochastic expression of IFN receptor IFNAR [44].

Discussion

Regulation of type I IFN production is essential for the innate immune response to viral infections [45,46]. However, high levels of IFN β can be toxic [47,48]. Thus, IFN β production must be tightly regulated. This regulation appears to be both temporal and stochastic. Type I IFN genes are tightly repressed prior to virus infection, activated upon infection, and then rapidly turned off several hours later (Figure S1B and S1C). Previous studies of several cytokine genes suggest that this stochastic gene expression provides an additional mechanism of regulation whereby optimal levels of cytokine production are determined by the frequency of expressing cells rather than by protein levels per cell [18,19,49]. Thus, it is possible that stochastic expression is a primary mechanism for controlling the optimal level of IFN β production *in vivo*. In particular, we have shown that while IFN production is stochastic, the activation of the antiviral gene program by secreted IFN is not. Thus, stochastic expression of IFN would allow the regional distribution of the cytokine and activation of the surrounding cells, without producing toxic levels of IFN.

Previous studies have implicated as limiting steps enhanceosome assembly [20,21] and the assembly of an interchromosomal transcriptional hub formed through interactions between Alu

elements bearing NF- κ B sites [20]. More recently, the infecting virus, rather than intrinsic properties of the infected cell, has been implicated in this stochasm [22]. The data presented here reveal a far more complex mechanism in which cell-to-cell variations in limiting components required to support viral replication, to detect and signal the presence of viral RNA, and to activate transcription factors all contribute to the observed stochastic expression (Figure 7D). It seems likely that the key limiting factor varies between cell types, cell lines, and organisms.

The earliest step in the virus induction signaling pathway is entry of virus or dsRNA into the cell. We have shown that both inducers elicit stochastic expression, but in neither case is this due to limiting inducer (Figures S4B and 6A). We showed that both IFN β -producing and nonproducing cells were infected by SeV (Figure 2B). However, the IFN β -producing cells contained significantly higher levels of the products of viral replication and transcription. Thus, it appears that there are cell-to-cell differences in the ability to support efficient viral replication, and these differences influence the probability of IFN β gene expression. Presumably, high levels of RNA inducer in the IFN-producing cells overcome limiting amounts of RIG-I or MDA5. However, differences in viral replication alone cannot explain the observed stochasm in IFN β production. A previous study, using a cell line transfected with an IFN β -GFP reporter, concluded that stochastic IFN β expression is due entirely to heterogeneity in the infecting virus [22]. However, in that study the IFN β -GFP cell line was preselected to minimize stochastic expression of the reporter. In addition, that study involved a stably transfected gene, while the present study made use of the endogenous gene. The results presented here strongly indicate that heterogeneity of both the virus and host cells together are responsible for the stochastic expression of IFN β .

We have identified multiple limiting steps in the activation of IFN β gene expression, ranging from initial steps in virus infection and replication, to the signaling pathway, to the activation and binding of transcriptional activator proteins to the IFN β promoter. For example, over-expression of individual components in the RIG-I signaling pathway increases the percentage of IFN β -expressing cells. The largest increase was observed with IRF7, which lies at the endpoint of the RIG-I pathway, and also positively controls the expression of components in the RIG-I signaling pathway. Taken together, these data are consistent with a model in which the probability of expression of the IFN β gene in individual cells depends primarily on the activation of the RIG-I signaling pathway and the presence of sufficient numbers of IRF7 molecules to activate transcription (Figure 7D). This conclusion is consistent with the observation that both IFN β and IFN α are stochastically expressed in response to virus infection (Figure 1A and 1C). The expression of both genes requires activation of the RIG-I pathway and active IRF7 [50].

We find that limiting amounts of other RIG-I pathway components also contribute to stochastic expression of the IFN β gene, as we observed higher levels of RIG-I/Trim25 and MDA5 mRNA and protein levels in the IFN β -producing cells than in the nonproducers (Figure 3). In addition, over-expression of RIG-I and Trim25 together leads to a dramatic increase in the percentage of cells that express IFN β (Figure 4C and 4D). Similar






Figure 6. Poly I:C-induced stochastic IFN β expression depends on the amounts of poly I:C and MDA5. (A) IFN β /YFP homozygous MEF cells were electroporated with Cy5-labeled poly I:C, and FACS analysis was carried out 8 h after the electroporation to assay the strength of Cy5 and YFP. The top left panel shows untransfected MEF cells, and the bottom left panel shows the electroporated MEF cells. As indicated by arrows, the two panels to the right represent the “poly I:C high” and “poly I:C low” populations, respectively. Data shown are representative of at least three independent experiments. Numbers represent relative percentages. (B) L929-MDA5 or L929-RIG-I stable transfectants were stimulated with tetracycline (Tet) for 24 h followed by transient transfection with poly I:C. 6 h after transfection, cells were fixed, followed by RNA ISH to detect IFN β mRNA. (C) Bar plots representing the percentage (mean \pm standard deviation) of cells expressing IFN β from three independent ISH experiments performed as in (B). At least 400 cells were blindly counted and scored for each category. (D) IFN β /YFP primary MEFs were fixed 8 h after poly I:C stimulation. Intracellular staining using MDA5 antibody and FACS analysis were carried out to assay the correlation between the expression levels of IFN β and MDA5. Data shown are representative of at least three independent experiments. Numbers represent relative percentages. Iso-Ctrl, isotype control.
doi:10.1371/journal.pbio.1001249.g006

results were obtained with high levels of expression of the RIG-I signaling components MAVS and TBK1 and the transcription factors IRF3 and IRF7 (Figures 4A, 4B, and 5). Thus, it appears that many, if not all, of the components in the RIG-I signaling pathway, from the sensors of viral RNA to the essential transcription factors, can be limiting components in the virus induction pathway.

The largest increase in the percentage of IFN-producing cells was observed when IRF7 was over-expressed. IRF7 is the master regulator of type I IFN gene expression [25], and is present at low levels in all cell types except plasmacytoid dendritic cells, where it is constitutively abundant [26,40]. Our over-expression experiments show that high levels of IRF7 promote the transcription of type I IFN genes (Figure S7D), and essentially eliminate the stochastic expression of both the IFN β and α genes (Figures 5 and S7). In a previous study in human cells, both NF- κ B and IRF7 over-expression was shown to partially suppress stochastic IFN β expression [20]. Our results are consistent with this observation. However, there are two differences. First, based, at least in part, on the lack of requirement of NF- κ B in murine cells, we observed a relatively small effect of increasing NF- κ B expression. Second, we saw a greater effect of IRF7 expression in murine cells than was observed in human cells. Over-expression of IRF7 in L929 cells almost completely eliminated stochastic expression of both IFN β and α genes, while in human HeLa cells high levels of IRF7 increase the percentage of IFN β -producing cells to almost 55% [20]. Deleting the IRF7 translational repressors, 4E-BPs, also increased the IFN β -expressing MEFs by 4-fold (Figure 5C and 5D). We also showed that the RIG-I signaling pathway, and in particular RIG-I and Trim25, are up-regulated in IRF7 over-expressing cells (Figure S8). We conclude that limiting amounts of active IRF7 appear to be overcome by two mechanisms: positive auto-regulation of IRF7 expression, and IRF7-dependent up-regulation of the RIG-I signaling pathway.

We note that in addition to IFN β , several other virus-inducible genes, including TNF α , IL-6, CCL4, and CCL5, are highly expressed in the IFN β -producing cells compared to nonproducers, suggesting that many, if not all, of the virus-inducible genes are stochastically expressed. The common feature of the activation of all of these genes is that they all require the RIG-I signaling pathway [28–31]. Thus, we conclude that stochastic gene expression is primarily due to limiting components in the signaling pathway but not gene-to-gene variation in the mechanism of gene activation.

We showed that although the IFN β gene is stochastically expressed upon virus infection, the antiviral ISGs, e.g., ISG15, were equally induced in all cells (Figure 7C). However, we note that RIG-I, Trim25, and MDA5, which are also antiviral ISGs, are highly expressed in IFN β -producing cells compared to nonproducing cells (Figure 3C and 3D). We believe that the differences we observed here reflect naturally occurring cell-to-cell variability in the levels of expression of these genes prior to virus

infection, and that this variability is the primary source of stochastic IFN β gene expression. However, at later times after virus infection, we expect that the differences in the mRNA or protein levels of these genes between the YFP-positive and YFP-negative populations will be much smaller compared to those at earlier stages (8 h.p.i.). As shown in Figure S11A and S11B, our qPCR data and Western blot data support this expectation. The IFN β gene is also stochastically expressed in IFNAR-deficient MEFs, which suggests that the IFNAR levels or an IFN β feedback loop are not major factors responsible for stochastic IFN β gene expression (Figure S11C). We further measured the distributions of six components in the RIG-I signaling pathway. As shown in Figures 7A, 7B, and S10, all six proteins were log-normally distributed across the cell population, an observation that is consistent with data on other proteins [51,52]. Thus, naturally occurring differences in the protein levels and activities of individual signaling pathway components and transcription factors account for stochastic IFN β expression induced by both poly I:C induction and virus infection.

Previous studies have shown that naturally occurring differences in the levels of proteins in the apoptotic signaling pathway are the primary reasons for cell-to-cell variability in the probability of cell death [52]. Thus, the results presented here not only reveal the complexity of the regulatory mechanisms controlling stochastic IFN β gene expression, but also suggest a general mechanism used in different biological processes to establish and control stochastic gene expression. A remarkable feature of stochastic expression is that it appears to be an intrinsic property of different clonal populations of cells. For example, if a particular cell line displays a certain percentage of activated cells, that percentage differs from other cell lines, and is retained when the cells are recloned [14]. Thus, the extent of stochasm appears to be a genetic and epigenetic feature of clonal cell populations.

Materials and Methods

Cells, Reagents, and Plasmids

All cell lines, including L929, RAW 264.7, MG63, and 293T, were from the American Type Culture Center; primary MEFs were isolated using standard protocols from IFN β /YFP mice [24]. Primary human foreskin fibroblast cells were purchased from PromoCell. All cells were cultured in DMEM (Gibco) supplemented with 10% FBS (Gibco) in a 5% CO₂ incubator. Cycloheximide was purchased from Sigma-Aldrich. Human and mouse recombinant IFN proteins were purchased from PBL Interferonsource. Brefeldin A solution was purchased from eBioscience. Poly I:C was purchased from InvivoGen. Cy5-labeled poly I:C was generated using Label IT Nucleic Acid Labeling Kit (Mirus). The different expression constructs were generated by cloning the coding sequences of each gene by PCR and inserting them into the vector pt-REX-DEST30, which has the tetracycline-inducible promoter (Invitrogen).




Figure 7. Endogenous variation in the concentrations of components of the RIG-I signaling pathway. (A and B) Protein distributions in untreated primary MEFs determined by flow cytometry. a.u., arbitrary units. (C) Mouse ISG15 gene expression in MEFs 6 h after IFN β treatment, detected by ISH using a digoxigenin-labeled ISG15 probe. (D) A model depicting stochastic IFN gene expression. There is a population of ten cells with varying numbers of limiting factors in each cell. Each small, colored circle represents one of the limiting factors, and six limiting factors are

shown. Short black lines represent viral inducer. Only two cells in the population have enough of the viral inducer and all six factors to trigger transcription of the IFN β gene. Iso-Ctrl, isotype control.
doi:10.1371/journal.pbio.1001249.g007

Virus Infection and Poly I:C Transfection

Concentrated SeV stock (Cantell strain, Charles River Lab) was added to cultured cells at a concentration of 200 HAU/ml and incubated for the times indicated. Poly I:C transfection was carried out using either lipofectamine2000 (Invitrogen) or electroporation using Amaxa MEF2 Nucleofector Kit (Lonza).

RNA Preparation and PCR

Total RNA was extracted with Trizol reagent (Invitrogen). Real-time quantitative reverse transcription PCR (qRT-PCR) was conducted according to standard protocols.

Antibodies and Western Blot

Antibody against YFP was from Chemicon (Millipore) or Abcam. RIG-I, MAVS, and GAPDH antibodies were from Cell Signaling. Antibodies against p65, HDAC1, and Trim25 were from Santa Cruz Biotechnology. MDA5 and TBK1 antibodies were from Abcam and Imgenex, respectively. IFN β antibody used for FACS was from Millipore. SeV antibodies were kindly provided by Dr. Atsushi Kato (National Institute of Infectious Diseases, Japan). Nuclear/cytosol fractionation was performed using Nuclear/Cytosol Fractionation Kit (BioVision). Western blots were carried out using standard protocols.

In Situ Hybridization

Antisense RNA probes recognizing mouse IFN β or β -actin were synthesized using T7 or SP6 polymerase and digoxigenin-labeled nucleotides (Roche Applied Science). Cells were cultured on poly-D-lysine-coated 24-well plates (Fisher) and either mock- or virus-infected for the times indicated. Cells were then washed twice with PBS and fixed with 4% paraformaldehyde. Hybridization, washes, and staining were carried out as previously described [53].

Flow Cytometry

MEF cells were fixed with IC Fixation Buffer and permeabilized with Permeabilization Buffer (both from eBioscience). After incubation with appropriate antibodies, flow cytometry was done with a FACSCalibur, and data were analyzed with CellQuest software (both from Becton Dickinson).

Microarray Analysis and KEGG Pathway Enrichment Analysis

Total RNA from untreated and tetracycline-induced L929-IRF7 cells were prepared using Trizol reagent (Invitrogen) followed by purification using MEGAclear (Ambion). Biotinylated RNA probes were synthesized by two rounds of amplification using the MessageAmp II aRNA Amplification kit (Ambion). The probes were hybridized with Affymetrix Mouse Genome 430A_2.0 array chips. Affymetrix DAT files were processed using the Affymetrix Gene Chip Operating System to create CEL files. Normalized expression values were analyzed with the Bioconductor Limma package, an approach for implementing empirical Bayes linear modeling [42]. For all comparison tests, genes with an absolute fold change in transcript level exceeding 1.5 and $p < 0.05$ were selected for further analyses. The likelihood of overrepresentation of KEGG signaling pathways in the up- or down-regulated gene list relative to a background of all array genes was calculated by Fisher's exact test for statistical analysis.

Supporting Information

Figure S1 Stochastic expression of IFN β gene upon virus infection. (A) Stochastic expression of human IFN β gene in primary foreskin fibroblast cells, MG63 cells, and 293T cells 6 h.p.i., detected by ISH using a digoxigenin-labeled antisense RNA IFN β probe. Numbers on the right indicate the percentages of IFN β -expressing cells in each cell type. (B and C) Kinetics of IFN β expression in primary MEFs (B) or L929 cells (C) assayed by qRT-PCR. IFN β mRNA can be detected as early as 6 h.p.i., and maximum levels are observed at 9 and 12 h.p.i. in MEFs and L929 cells, respectively. (D) β -actin mRNA in L929 cells was detected by ISH using either digoxigenin-labeled sense or antisense RNA probe. (E) MEF cells or human MG63 cells were infected by SeV for 9 h or 6 h, respectively. ISH was carried out to detect the IFN β -expressing cells using an IFN β sense or antisense probe. (F) Human MG63 cells were infected by SeV for 9 h. IFN β protein was detected by immunocytochemistry using IFN β antibody. Similar percentages of IFN β -expressing cell were detected by either ISH or immunocytochemistry. (G) Human IFN α 8 mRNA in Namalwa cells was detected by ISH using digoxigenin-labeled probe. (TIF)

Figure S2 IFN expression in sorted MEFs and allelic expression of IFN β gene. (A) qRT-PCR analysis illustrating the expression levels of IFN genes in sorted IFN β /YFP primary MEF cells. (B) IFN β /YFP heterozygous MEFs (upper panel) and homozygous MEFs (lower panel) were infected by SeV for variable times with the presence of Brefeldin A (BFA)—which inhibits transport of proteins from endoplasmic reticulum to Golgi—in the last 4 h. Cells were fixed and stained for intracellular IFN β and YFP. If IFN β gene is monoallelically expressed, heterozygous MEF cells should have similar percentages of IFN β -positive population (from IFN β allele) and IFN β /YFP double-positive population (from IFN β -IRES-YFP allele). If IFN β gene is biallelically expressed, all, or at least most, of IFN β -expressing heterozygous cells should be both IFN β - and YFP-positive. Heterozygous MEF FACS analysis (upper panel) showed a similar percentage of IFN β -positive population (upper left panel, 5.24%) and IFN β /YFP double-positive population (upper right panel, 6.30%) at 8 h.p.i., suggesting that the IFN β gene expression was predominantly monoallelic before 8 h.p.i. During the time periods 8–12 h.p.i. and 12–16 h.p.i., the majority of IFN β -expressing cells were IFN β /YFP double-positive (upper right panel, 11.01%, and upper right panel, 3.90%, respectively), indicating that at late infection, IFN β gene expression was biallelic. As control, shown in the lower panel, IFN β -expressing homozygous MEF cells had almost no IFN β single-positive population at any given time point. Data shown are representative of at least three independent experiments. Numbers represent relative percentages. (TIF)

Figure S3 Priming of cells increases the percentage of IFN β -expressing cells. (A) Primary MEFs were primed with 250 U/ml IFN β or 250 U/ml IFN β plus 50 μ g/ml cycloheximide (CHX) for 6 h, infected by SeV, and subjected to ISH using digoxigenin-labeled IFN β RNA probe. (B) Histogram showing the percentage (mean \pm standard deviation) of cells expressing IFN β from three independent ISH experiments as in (A). (C) Human foreskin fibroblasts were primed with 250 U/ml IFN β or 250 U/ml IFN β plus 50 μ g/ml cycloheximide for 6 h, infected by SeV,

and subjected to ISH using digoxigenin-labeled IFN β RNA probe. (D) Histogram showing the percentage (mean \pm standard deviation) of cells expressing IFN β from three independent ISH experiments as in (C). (TIF)

Figure S4 Viral titer is not a limiting factor. (A) MEF cells were infected by SeV. Cells were fixed and stained for SeV using SeV antibody. Blue color shows DAPI staining (nucleus) and green color shows SeV signal. Most, if not all, cells are uniformly exposed to SeV. Scale bar, 20 μ m. (B) Percentages (mean \pm standard deviation) of IFN β -producing primary MEF cells infected by increasing amounts of SeV. At least 400 cells were counted and scored blindly for each category. In all of the experiments described in this study, we used 100–200 HAU/ml of SeV as the infecting dose. (C) qRT-PCR analysis illustrating the relative abundance of viral NP mRNA in sorted IFN β /YFP primary MEF populations 8 or 12 h.p.i. (D) L929 cells were transfected with total RNAs either from IFN β -producing or IFN β -nonproducing MEF cells sorted after being virus-infected for 8 h or 12 h. Then total RNAs were extracted from these L929 cells 8 h after transfection, and qRT-PCR experiments were carried out to detect relative abundance of IFN β mRNA in these transfected cells. (TIF)

Figure S5 Over-expression of RIG-I, MDA5, or Trim25 increases the percentage of IFN β -expressing cells. Different tetracycline-inducible L929 stable transfectants were generated. In the absence of tetracycline, expression of exogenous copies of these genes is tightly repressed in the stable transfectants, but upon the addition of 1 μ g/ml tetracycline, the stably incorporated genes are expressed at a high level. (A) Western blots showing the tetracycline-inducible expression levels of different proteins. All genes were flag tagged and proteins were detected using Flag antibody. (B) L929-RIG-I, L929-MDA5, and L929-Trim25 stable transfectants were induced by tetracycline for 24 h, followed by virus infection for 9 h. RNA ISH experiments were carried out to detect the IFN β mRNA. (C) Histogram showing the percentage (mean \pm standard deviation) of cells expressing IFN β from three independent ISH experiments. At least 400 cells were blindly counted and scored for each category. (D) L929-RIG-I stable transfectants were transfected with GFP control plasmid. The transfection efficiency was measured by GFP detection using fluorescent microscopy. (TIF)

Figure S6 Higher RIG-I and Trim25 protein levels are present in IFN β -expressing cells. IFN β /YFP homozygous MEF cells were infected with SeV for 8 h. Cells were then fixed and intracellularly stained for RIG-I and Trim25. FACS analysis was used to assay the correlation between IFN β expression (detected by YFP) and RIG-I/Trim25 expression. The top panel shows the expression of RIG-I and Trim25 in MEFs before and after virus infection. The middle panel shows the percentage of IFN β -expressing cells before and after virus infection. The bottom panel shows the RIG-I/Trim25 expression level in YFP-positive cells in the middle panel, which represent IFN β -expressing cells. Data shown are representative of at least three independent experiments. Numbers represent relative percentages. (TIF)

Figure S7 Over-expression of IRF7 eliminates stochastic IFN α gene expression. (A) L929-IRF7 stable transfectants were treated with 1 μ g/ml tetracycline for 24 h followed by immunocytochemistry using Flag antibody detecting the exogenous Flag-tagged IRF7 expression level. Increased IRF7 expres-

sion level was detected in almost every cell. Scale bar, 20 μ m. (B) The L929-IRF7 stable transfectant was induced by tetracycline for 24 h, followed by virus infection for 9 h. RNA ISH experiments were carried out to detect IFN α mRNA. (C) Histogram showing the percentage (mean \pm standard deviation) of cells expressing IFN α from three independent ISH experiments. At least 400 cells were blindly counted and scored for each category. (D) L929-IRF7 stable transfectants were treated with or without tetracycline for 24 h before SeV infection. Total RNA was extracted and semi-qRT-PCR was carried out to measure the kinetics of transcription of different type I IFN mRNAs. (TIF)

Figure S8 IRF7 up-regulates RIG-I-like receptor signaling pathway. (A) Partial list of genes whose expression in L929-IRF7 cells, as assessed by genome-wide expression profiling, was increased as a result of tetracycline induction to the level of these genes in control cells without tetracycline treatment. The names of known RIG-I-like signaling pathway genes from the KEGG Pathway Database are highlighted in grey. Asterisk indicates that the Ddx58 gene expression profile was undetectable because of the lack of a corresponding probe set on the Affymetrix Mouse Genome 430A_2.0 array chips. The fold increase of Ddx58 gene expression was determined by qRT-PCR. (B) RIG-I-like receptor signaling pathway is the most significantly up-regulated pathway identified from KEGG pathway enrichment analysis ($p = 3.6E-06$) based on L929-IRF7 microarray results. *, $p < 0.05$; ***, $p < 0.001$. (C) qRT-PCR analysis illustrating the levels of expression of IRF7, RIG-I, Trim25, and MDA5 in L929-IRF7 stable transfectants before and after induction with tetracycline. (TIF)

Figure S9 IRF3 and IRF7 are limiting factors in stochastic IFN β expression induced by poly I:C transfection. (A) L929-IRF3, L929-IRF7, or L929-p65 stable transfectants were stimulated with tetracycline for 24 h followed by transient transfection with poly I:C. 12 h after transfection, cells were fixed followed by RNA ISH to detect IFN- β mRNA. (B) Bar plots representing the percentage (mean \pm standard deviation) of cells expressing IFN β from three independent ISH experiments performed as in (A). At least 400 cells were blindly counted and scored for each category. (TIF)

Figure S10 Endogenous variation in the concentrations of components of the RIG-I signaling pathway. Primary MEF cells were fixed by 4% PFA followed by intracellular staining using appropriate antibodies recognizing different components of the signaling pathway. The intensity of immunofluorescent signal was quantified using ImageJ software. For individual factors, the highest immunofluorescence intensity was set as 1. The x -axis shows the relative immunofluorescence intensity of each factor. Each plot represents the immunofluorescence intensity calculated from 150–200 cells. (TIF)

Figure S11 Differences in the levels of RIG-I signaling pathway factors between YFP-positive and YFP-negative populations. (A) qPCR analysis illustrating the expression levels of IFN β , RIG-I, Trim25, and MDA5 genes in sorted MEF cells at 8, 12, and 24 h.p.i. (B) Western blots showing cytoplasmic distribution of signaling pathway proteins present in FACS-sorted cells at 8, 12, and 24 h.p.i. (C) IFNAR-deficient MEFs were infected by SeV. IFN β expression was detected by ISH using IFN β antisense RNA probe. (TIF)

Acknowledgments

We thank Dr. Atsushi Kato (National Institute of Infectious Diseases, Japan) for providing SeV antibodies. We thank Dr. Nahum Sonenberg (McGill University, Canada) for providing 4E-BP1/4E-BP2 double-knockout MEFs.

References

- Honda K, Takaoka A, Taniguchi T (2006) Type I interferon [corrected] gene induction by the interferon regulatory factor family of transcription factors. *Immunity* 25: 349–360.
- Takeuchi O, Akira S (2009) Innate immunity to virus infection. *Immunol Rev* 227: 75–86.
- Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, et al. (2006) Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. *Nature* 441: 101–105.
- Gack MU, Shin YC, Joo CH, Urano T, Liang C, et al. (2007) TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated antiviral activity. *Nature* 446: 916–920.
- Cui S, Eisenacher K, Kirchofer A, Brzozka K, Lammens A, et al. (2008) The C-terminal regulatory domain is the RNA 5'-triphosphate sensor of RIG-I. *Mol Cell* 29: 169–179.
- Takahashi K, Yoneyama M, Nishihori T, Hirai R, Kumeta H, et al. (2008) Nonself RNA-sensing mechanism of RIG-I helicase and activation of antiviral immune responses. *Mol Cell* 29: 428–440.
- Seth RB, Sun L, Ea CK, Chen ZJ (2005) Identification and characterization of MAVS, a mitochondrial antiviral signaling protein that activates NF-kappaB and IRF 3. *Cell* 122: 669–682.
- tenOever BR, Sharma S, Zou W, Sun Q, Grandvaux N, et al. (2004) Activation of TBK1 and IKKvarepsilon kinases by vesicular stomatitis virus infection and the role of viral ribonucleoprotein in the development of interferon antiviral immunity. *J Virol* 78: 10636–10649.
- Fitzgerald KA, McWhirter SM, Faia KL, Rowe DC, Latz E, et al. (2003) IKKepsilon and TBK1 are essential components of the IRF3 signaling pathway. *Nat Immunol* 4: 491–496.
- Sharma S, tenOever BR, Grandvaux N, Zhou GP, Lin R, et al. (2003) Triggering the interferon antiviral response through an IKK-related pathway. *Science* 300: 1148–1151.
- Kawai T, Takahashi K, Sato S, Coban C, Kumar H, et al. (2005) IPS-1, an adaptor triggering RIG-I- and Mda5-mediated type I interferon induction. *Nat Immunol* 6: 981–988.
- Ford E, Thanos D (2010) The transcriptional code of human IFN-beta gene expression. *Biochim Biophys Acta* 1799: 328–336.
- Enoch T, Zinn K, Maniatis T (1986) Activation of the human beta-interferon gene requires an interferon-inducible factor. *Mol Cell Biol* 6: 801–810.
- Zawatzky R, De Maeyer E, De Maeyer-Guignard J (1985) Identification of individual interferon-producing cells by in situ hybridization. *Proc Natl Acad Sci U S A* 82: 1136–1140.
- Hollander GA (1999) On the stochastic regulation of interleukin-2 transcription. *Semin Immunol* 11: 357–367.
- Guo L, Hu-Li J, Paul WE (2004) Probabilistic regulation of IL-4 production in Th2 cells: accessibility at the I4 locus. *Immunity* 20: 193–203.
- Guo L, Hu-Li J, Paul WE (2005) Probabilistic regulation of IL-4 production. *J Clin Immunol* 25: 573–581.
- Calado DP, Paixao T, Holmberg D, Haury M (2006) Stochastic monoallelic expression of IL-10 in T cells. *J Immunol* 177: 5358–5364.
- Kelly BL, Locksley RM (2000) Coordinate regulation of the IL-4, IL-13, and IL-5 cytokine cluster in Th2 clones revealed by allelic expression patterns. *J Immunol* 165: 2982–2986.
- Apostolou E, Thanos D (2008) Virus infection induces NF-kappaB-dependent interchromosomal associations mediating monoallelic IFN-beta gene expression. *Cell* 134: 85–96.
- Hu J, Sealfon SC, Hayot F, Jayaprakash C, Kumar M, et al. (2007) Chromosome-specific and noisy IFNB1 transcription in individual virus-infected human primary dendritic cells. *Nucleic Acids Res* 35: 5232–5241.
- Chen S, Short JA, Young DF, Killip MJ, Schneider M, et al. (2010) Heterocellular induction of interferon by negative-sense RNA viruses. *Virology* 407: 247–255.
- Balazsi G, van Oudenaarden A, Collins JJ (2011) Cellular decision making and biological noise: from microbes to mammals. *Cell* 144: 910–925.
- Scheu S, Dressing P, Locksley RM (2008) Visualization of IFNbeta production by plasmacytoid versus conventional dendritic cells under specific stimulation conditions in vivo. *Proc Natl Acad Sci U S A* 105: 20416–20421.
- Honda K, Yanai H, Negishi H, Asagiri M, Sato M, et al. (2005) IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* 434: 772–777.
- Sato M, Suemori H, Hata N, Asagiri M, Ogasawara K, et al. (2000) Distinct and essential roles of transcription factors IRF-3 and IRF-7 in response to viruses for IFN-alpha/beta gene induction. *Immunity* 13: 539–548.

Author Contributions

The author(s) have made the following declarations about their contributions: Conceived and designed the experiments: MZ HP TM. Performed the experiments: MZ. Analyzed the data: MZ JZ. Contributed reagents/materials/analysis tools: SS. Wrote the paper: MZ HP TM.

- Wang X, Hussain S, Wang EJ, Li MO, Garcia-Sastre A, et al. (2007) Lack of essential role of NF-kappa B p50, RelA, and cRel subunits in virus-induced type 1 IFN expression. *J Immunol* 178: 6770–6776.
- Kato H, Sato S, Yoneyama M, Yamamoto M, Uematsu S, et al. (2005) Cell type-specific involvement of RIG-I in antiviral response. *Immunity* 23: 19–28.
- Poock H, Bscheider M, Gross O, Finger K, Roth S, et al. (2010) Recognition of RNA virus by RIG-I results in activation of CARD9 and inflammasome signaling for interleukin 1 beta production. *Nat Immunol* 11: 63–69.
- Tamassia N, Le Moigne V, Rossato M, Donini M, McCartney S, et al. (2008) Activation of an immunoregulatory and antiviral gene expression program in poly(I:C)-transfected human neutrophils. *J Immunol* 181: 6563–6573.
- Yoshida H, Imaizumi T, Lee SJ, Tanji K, Sakaki H, et al. (2007) Retinoic acid-inducible gene-I mediates RANTES/CCL5 expression in U373MG human astrocytoma cells stimulated with double-stranded RNA. *Neurosci Res* 58: 199–206.
- Baum A, Sachidanandam R, Garcia-Sastre A (2010) Preference of RIG-I for short viral RNA molecules in infected cells revealed by next-generation sequencing. *Proc Natl Acad Sci U S A* 107: 16303–16308.
- Rehwinkel J, Tan CP, Goubau D, Schulz O, Pichlmair A, et al. (2010) RIG-I detects viral genomic RNA during negative-strand RNA virus infection. *Cell* 140: 397–408.
- Seth RB, Sun L, Chen ZJ (2006) Antiviral innate immunity pathways. *Cell Res* 16: 141–147.
- Wang J, Basagoudanavar SH, Wang X, Hopewell E, Albrecht R, et al. (2010) NF-kappa B RelA subunit is crucial for early IFN-beta expression and resistance to RNA virus replication. *J Immunol* 185: 1720–1729.
- Drahoš J, Racaniello VR (2009) Cleavage of IPS-1 in cells infected with human rhinovirus. *J Virol* 83: 11581–11587.
- Meylan E, Curran J, Hofmann K, Moradpour D, Binder M, et al. (2005) Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by hepatitis C virus. *Nature* 437: 1167–1172.
- You F, Sun H, Zhou X, Sun W, Liang S, et al. (2009) PCB2P mediates degradation of the adaptor MAVS via the HECT ubiquitin ligase AIP4. *Nat Immunol* 10: 1300–1308.
- Yang K, Shi H, Qi R, Sun S, Tang Y, et al. (2006) Hsp90 regulates activation of interferon regulatory factor 3 and TBK-1 stabilization in Sendai virus-infected cells. *Mol Biol Cell* 17: 1461–1471.
- Prakash A, Smith E, Lee CK, Levy DE (2005) Tissue-specific positive feedback requirements for production of type I interferon following virus infection. *J Biol Chem* 280: 18651–18657.
- Colina R, Costa-Mattioli M, Dowling RJ, Jaramillo M, Tai LH, et al. (2008) Translational control of the innate immune response through IRF-7. *Nature* 452: 323–328.
- Smyth GK (2005) Limma: linear models for microarray data. In: Gentleman V, Carey V, Huber W, Irizarry R, Dudoit S, eds. *Bioinformatics and computational biology solutions using R and bioconductor*. New York: Springer. pp 397–420.
- Gitlin L, Barchet W, Gilfillan S, Cella M, Beutler B, et al. (2006) Essential role of mda-5 in type I IFN responses to polyriboinosinic:polyribocytidylic acid and encephalomyocarditis picornavirus. *Proc Natl Acad Sci U S A* 103: 8459–8464.
- Levin D, Harari D, Schreiber G (2011) Stochastic receptor expression determines cell fate upon interferon treatment. *Mol Cell Biol* 31: 3252–3266.
- Garcia-Sastre A, Biron CA (2006) Type I interferons and the virus-host relationship: a lesson in detente. *Science* 312: 879–882.
- Kawai T, Akira S (2006) Innate immune recognition of viral infection. *Nat Immunol* 7: 131–137.
- Banchereau J, Pascual V (2006) Type I interferon in systemic lupus erythematosus and other autoimmune diseases. *Immunity* 25: 383–392.
- Hall JC, Rosen A (2010) Type I interferons: crucial participants in disease amplification in autoimmunity. *Nat Rev Rheumatol* 6: 40–49.
- Hu-Li J, Pannetier C, Guo L, Lohning M, Gu H, et al. (2001) Regulation of expression of IL-4 alleles: analysis using a chimeric GFP/IL-4 gene. *Immunity* 14: 1–11.
- Yoneyama M, Kikuchi M, Natsukawa T, Shinobu N, Imaizumi T, et al. (2004) The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. *Nat Immunol* 5: 730–737.
- Sigal A, Milo R, Cohen A, Geva-Zatorsky N, Klein Y, et al. (2006) Variability and memory of protein levels in human cells. *Nature* 444: 643–646.
- Spencer SL, Gaudet S, Albeck JG, Burke JM, Sorger PK (2009) Non-genetic origins of cell-to-cell variability in TRAIL-induced apoptosis. *Nature* 459: 428–432.
- Senger K, Merika M, Agalio T, Yie J, Escalante CR, et al. (2000) Gene repression by coactivator repulsion. *Mol Cell* 6: 931–937.