

A Polyhedral Description of Kernels

Qin Chen^a Xujin Chen^{b*} Wenan Zang^{c†‡}

^a Department of Mathematics, China Jiliang University
Hangzhou 310018, China

^b Institute of Applied Mathematics, Chinese Academy of Sciences
Beijing 100190, China

^c Department of Mathematics, The University of Hong Kong
Hong Kong, China

Abstract

Let G be a digraph and let $\pi(G)$ be the linear system consisting of nonnegativity, stability, and domination inequalities. We call G kernel ideal if $\pi(H)$ defines an integral polytope for each induced subgraph H of G , and call G kernel Mengerian if $\pi(H)$ is totally dual integral (TDI) for each induced subgraph H of G . In this paper we show that a digraph is kernel ideal iff it is kernel Mengerian iff it contains none of three forbidden structures; our characterization yields a polynomial-time algorithm for the minimum weighted kernel problem on kernel ideal digraphs. We also prove that it is NP -hard to find a kernel of minimum size even in a planar bipartite digraph with maximum degree at most three.

MSC 2000 subject classification. Primary: 90C10, 90C27, 90C57.

OR/MS subject classification. Primary: Programming/graphs.

Key words. Digraph, kernel, polytope, algorithm, complexity.

*Supported in part by NSF of China under Grant 10771209 and by CAS Program for Cross and Cooperative Team of Science and Technology Innovation.

†Supported in part by the Research Grants Council of Hong Kong.

‡Corresponding author. E-mail: wzang@maths.hku.hk.

1 Introduction

Digraphs considered in this paper contain no loops nor parallel arcs unless otherwise stated. Let $G = (V, A)$ be a digraph and let U be a subset of V . We call U a *stable set* of G if no two vertices in U are connected by an arc in G , call U a *dominating set* of G if for each vertex $v \notin U$, there is an arc from v to U in G , and call U a *kernel* of G if it is both stable and dominating in G . The study of kernels dates back to 1944 when von Neumann and Morgenstern [18] employed them to describe the winning positions in 2-person games; since then kernels have attracted tremendous research effort, and have been used as powerful tools for tackling many important problems arising in diverse research fields, such as logic, computational complexity, artificial intelligence, combinatorics, and coding theory. As shown by Chvátal [10], it is *NP*-complete in general to decide if a digraph has a kernel; this decision problem remains *NP*-complete even when restricted to planar digraphs with degree at most three (see Fraenkel [7]). So the focus of extensive research concerning kernels has been on special classes of digraphs.

Throughout this paper by a cycle (or a path) in a digraph we mean a directed one. We call a digraph G *even* if it contains no odd cycles, and call G *bipartite* if its underlying graph is bipartite. Von Neumann and Morgenstern [18] observed that every acyclic digraph contains a kernel, which was generalized by Richardson [15] (see Theorem 12.3.2 in [2]) as follows.

Theorem 1.1. [15] *Every even digraph has a kernel.*

Richardson [15] established this result by showing that

$$\text{every strongly connected even digraph } H \text{ is bipartite,} \quad (1.1)$$

and thus each color class is a kernel of H . Applying (1.1) to a strongly connected component H with no outgoing arcs, we can recursively find a kernel in an arbitrary even digraph G in polynomial time.

Kernels are closely related to the so-called stable matchings. As demonstrated by Irving [13], the stable matching problem on a graph with strict preference orders can be solved in polynomial time. It follows that if a digraph G is an orientation of a line graph and each clique of G is acyclic, then there is a polynomial-time algorithm for finding a kernel in G , if any. For a digraph G with a perfect underlying graph, Boros and Gurvich [4] proved that if in every clique the subgraph induced by one-way arcs is acyclic, then G has a kernel; see Aharoni and Holzman [1] for a shorter proof based on Scarf's lemma and Király and Pap [12] for another shorter proof using Sperner's lemma and for an extension to h -perfect graphs. A more general setting where Sperner's lemma applies was studied by Edmonds, Gaubert and Gurvich [8]. We point out that, despite the existence, none of these proofs leads to a polynomial-time algorithm for finding a kernel. An open problem posed by Egerváry Research Group on Combinatorial Optimization (EGRES) is to find kernels in special classes of digraphs, including the aforementioned ones.

Problem 1.1. [6] *In which classes of digraphs can we decide if a kernel exists and find one in polynomial time?*

Given a digraph G with an integral weight $w(v)$ on each vertex v , the *minimum weighted kernel problem* (MWKP) is to find a kernel in G with minimum total weight. As remarked in the Egres Open [6] and confirmed by the following theorem, the MWKP is considerably more difficult than the existence problem (recall the above Richardson's theorem).

Theorem 1.2. *Given a planar bipartite digraph G with maximum degree at most three and given a positive integer k , it is NP-complete to decide if G has a kernel with at most k vertices.*

Motivated by the following Egres open problem, we study the MWKP using polyhedral and linear programming approaches in this paper.

Problem 1.2. [6] *For which classes of digraphs can we explicitly give a linear description of the convex hull of kernels?*

Let us introduce some notations and terminology before proceeding. As usual, we use \mathbb{R} and \mathbb{Z} to denote the sets of real numbers and integers, respectively. For any two sets Ω and K , where Ω is always a set of numbers and K is always finite, we use Ω^K to denote the set of vectors $\mathbf{x} = (x(k) : k \in K)$ whose coordinates are members of Ω . Let $A\mathbf{x} \geq \mathbf{b}$ be a linear system and let P denote the polyhedron $\{\mathbf{x} : A\mathbf{x} \geq \mathbf{b}\}$. We call P *integral* if P is the convex hull of all the integral vectors contained in it. It is well known that P is integral if and only if the minimum in the LP-duality equation

$$\min\{\mathbf{w}^T \mathbf{x} : A\mathbf{x} \geq \mathbf{b}\} = \max\{\mathbf{y}^T \mathbf{b} : \mathbf{y}^T A = \mathbf{w}^T, \mathbf{y} \geq \mathbf{0}\}$$

has an integral optimal solution, for every integral vector \mathbf{w} for which the optimum is finite. If, instead, the maximum in the equation enjoys this property, then the system $A\mathbf{x} \geq \mathbf{b}$ is called *totally dual integral* (TDI) [17]. The model of TDI systems plays a crucial role in combinatorial optimization, and serves as a general framework for establishing various min-max theorems because, as shown by Edmonds and Giles [5], total dual integrality implies primal integrality: if $A\mathbf{x} \geq \mathbf{b}$ is TDI and \mathbf{b} is integral, then P is integral.

Let $G = (V, A)$ be a digraph. For each $U \subseteq V$ (resp. $U \subseteq A$), we use $G \setminus U$ to denote the digraph obtained from G by deleting all members of U , and set $G \setminus u = G \setminus U$ and $V \setminus U = V \setminus u$ (resp. $A \setminus U = A \setminus u$) if $U = \{u\}$. For each $v \in V$, we use $N_G^+(v)$ (resp. $N_G^-(v)$) to denote the set of all out-neighbors (resp. in-neighbors) of vertex v , and set $d_G^+(v) = |N_G^+(v)|$ (resp. $d_G^-(v) = |N_G^-(v)|$); we shall drop the subscript G if there is no danger of confusion. Let $\pi(G)$ denote the following linear system:

$$0 \leq x(v) \leq 1 \quad \forall v \in V, \quad (1.2)$$

$$x(u) + x(v) \leq 1 \quad \forall (u, v) \in A, \quad (1.3)$$

$$x(v) + x(N_G^+(v)) \geq 1 \quad \forall v \in V, \quad (1.4)$$

where and throughout $x(U) = \sum_{u \in U} x(u)$ for any $U \subseteq V$. Let \mathcal{C} be the collection of all cliques of G , and let $\sigma(G)$ be the linear system consisting of (1.2), (1.4), and the following inequalities:

$$x(C) \leq 1 \quad \forall C \in \mathcal{C}. \quad (1.5)$$

Observe that the upper bound in (1.2) is redundant unless v is an isolated vertex, and that

$$\text{the incidence vectors of kernels of } G \text{ are precisely integral solutions } \mathbf{x} \in \mathbb{Z}^V \text{ of } \pi(G). \quad (1.6)$$

The statement remains valid if we replace $\pi(G)$ by $\sigma(G)$. In the literature (1.2) – (1.5) are referred to as *nonnegativity*, *stability*, *domination*, and *clique* inequalities, respectively. The *kernel polytope* of G , denoted by $K(G)$, is the convex hull of incidence vectors of all kernels of G . It is clear that Problem 1.2 essentially asks for the defining system of $K(G)$.

The following is a reformulation of Rothblum's theorem [16] on stable matchings.

Theorem 1.3. [16] *Let G be an orientation of the line graph of a bipartite graph in which every clique is acyclic. Then $K(G)$ is defined by $\sigma(G)$.*

In [11], Király and Pap obtained the following strengthening of this result.

Theorem 1.4. [11] *Let G be an orientation of the line graph of a bipartite graph in which every clique is acyclic. Then $\sigma(G)$ is a TDI system.*

While these two theorems concern themselves with the system $\sigma(G)$, the present paper aims to explore integrality properties enjoyed by $\pi(G)$. Let $FK(G)$ denote the set of all solutions $\mathbf{x} \in \mathbb{R}^V$ of $\pi(G)$. We call $FK(G)$ the *fractional kernel polytope* of G . Clearly, $K(G) \subseteq FK(G)$; however, equality need not hold in general. Thus a natural question to ask is the following.

Problem 1.3. *How difficult is it to recognize all digraphs G for which $K(G) = FK(G)$ (resp. $\pi(G)$ is TDI)?*

It is worthwhile pointing out that, first, $K(G) = FK(G)$ iff $FK(G)$ is integral; second, if a digraph G happens to enjoy this primal integrality (resp. total dual integrality), then this fact cannot be certified by exhibiting “forbidden” subgraphs of G , because every digraph H is an induced subgraph of a digraph G for which $K(G) = FK(G)$ (resp. $\pi(G)$ is TDI). To see this, add a new vertex v to H and arcs (u, v) for all vertices u of H . Then the resulting digraph G has a unique kernel $\{v\}$ and hence is as desired. Even in the absence of sinks, the integrality property $K(G) = FK(G)$ is still not closed under taking subgraphs or induced subgraphs: Let G be the digraph depicted in Figure 1. Then $K(G) = FK(G)$ but neither $G \setminus v_5$ nor $G \setminus \{(v_2, v_5), (v_5, v_6)\}$ satisfies the corresponding equality. (To justify this, let \mathbf{x} be an arbitrary vector in $FK(G)$, let θ be the value of $x(v_4)$, and let δ be the value of $x(v_6)$. Using (1.2) – (1.4), it can be shown that $\theta = \delta$, and $x(v_i) = \theta$ if i is even and $1 - \theta$ otherwise. Note that both $\{v_1, v_3, v_5\}$ and $\{v_2, v_4, v_6\}$ are kernels of G . Moreover, $G \setminus v_5$ is a so-called gear, whose fractional kernel polytope is not integral as we shall show in the main theorem.) So it is unlikely to have a characterization of all digraphs G with $K(G) = FK(G)$ in terms of forbidden structures. Therefore, Problem 1.3 would be very challenging, if not intractable.




Figure 1: A digraph G with $K(G) = FK(G)$ and $K(G \setminus v_5) \neq FK(G \setminus v_5)$.

A digraph G is called *kernel ideal* if $K(H) = FK(H)$ for each induced subgraph H of G , and called *kernel Mengerian* if $\pi(H)$ is a TDI system for each induced subgraph H of G . By the aforementioned Edmonds-Giles theorem [5], every kernel Mengerian digraph is kernel ideal. The main purpose of this paper is to give a structural characterization of all kernel ideal and kernel Mengerian digraphs.

We digress to define a few terms before presenting the main result. In this paper a path P is a finite sequence $v_0 v_1 \dots v_k$, such that (v_i, v_{i+1}) is an arc for $1 \leq i \leq k-1$, and that v_0, v_1, \dots, v_k are distinct except possibly $v_0 = v_k$ (in this case P is a cycle). We follow the convention to call P a v_0 - v_k path, and call v_0 the *origin* of P and v_k the *terminus* of P ; both v_0 and v_k are referred to as the ends of P . The length of P is denoted by $|P|$. For any two vertices v_i and v_j on P with $i < j$, we use $P[v_i, v_j]$ to denote the segment of P from v_i to v_j , and set $P(v_i, v_j) = P[v_i, v_j] \setminus v_i$, $P[v_i, v_j) = P[v_i, v_j] \setminus v_j$, and $P(v_i, v_j) = P[v_i, v_j] \setminus \{v_i, v_j\}$. For notational simplicity, we also write $u \in P$ for $u \in V(P)$. Given a directed cycle C and two vertices a, b on C , we use $C[a, b]$ to denote the segment of C from a to b .

A digraph C is called a *circuit* if either its underlying graph is an undirected cycle, but C itself is not a directed cycle, or C is the digraph (on two vertices) formed by two parallel arcs. Notice that the number of sources equals the number of sinks in any circuit.

Let C be a circuit, let t_1, t_2, \dots, t_k be all the sinks of C , and let O_1, O_2, \dots, O_k be k vertex-disjoint cycles outside C . For each $i = 1, 2, \dots, k$, we perform precisely one of the following three operations with respect to t_i :

- identify t_i with a vertex on O_i ;
- add a directed path P_i from t_i to O_i ;
- split t_i into two vertices t_i^1 and t_i^2 (each of them is incident with one arc on C) and identify t_i^1 with a vertex on O_i and t_i^2 with another vertex on O_i .

The resulting digraph is called a *gear* if it is even and contains no parallel arcs. Furthermore, P_i 's are pairwise vertex-disjoint.

A *ring* is obtained from a directed cycle C by adding an s_i - t_i path P_i for $i = 1, 2$, such that

- s_1, t_1, s_2, t_2 occur on C in order when we traverse C in its direction from s_1 ;
- $s_i \neq t_i$ (but possibly $s_i = t_{3-i}$) for $i = 1, 2$;
- $P_1(s_1, t_1)$, $P_2(s_2, t_2)$, and C are pairwise vertex-disjoint;
- $P_1 \cup P_2 \cup C$ is even and contains no parallel arcs; and
- $|C[s_1, s_2]|$ is odd.

Now we are ready to state the main theorem of this paper.

Theorem 1.5. *For a digraph G , the following statements are equivalent:*

- (i) G contains no subgraph isomorphic to an odd cycle, a gear, or a ring;
- (ii) G is kernel ideal; and
- (iii) G is kernel Mengerian.

To interpret statements (ii) and (iii) in the preceding theorem, let $\mathbb{P}(G, \mathbf{w})$ stand for the linear program

$$\begin{aligned} & \text{Minimize} && \mathbf{w}^T \mathbf{x} \\ & \text{subject to} && \mathbf{x} \in FK(G), \end{aligned} \tag{1.7}$$

and let $\mathbb{D}(G, \mathbf{w})$ denote its dual. Note that if G contains no isolated vertex, then (1.2) can be replaced by $x(v) \geq 0$ for all $v \in V$ in the definition of $FK(G)$. Thus $\mathbb{D}(G, \mathbf{w})$ can be simplified

as

$$\begin{aligned} \text{Maximize} \quad & -y(A) + z(V) \\ \text{subject to} \quad & -\sum_{v \in e} y(e) + z(v) + z(N_G^-(v)) \leq w(v) \quad \forall v \in V, \end{aligned} \quad (1.8)$$

$$y(e) \geq 0 \quad \forall e \in A, \quad (1.9)$$

$$z(v) \geq 0 \quad \forall v \in V, \quad (1.10)$$

where by $v \in e$ we mean v is an end of arc e . Using these notations, we see that G is kernel ideal if and only if $\mathbb{P}(H, \mathbf{w})$ has an integral optimal solution for any induced subgraph H of G and any $\mathbf{w} \in \mathbb{Z}^{V(H)}$, and that G is kernel Mengerian if and only if both $\mathbb{P}(H, \mathbf{w})$ and $\mathbb{D}(H, \mathbf{w})$ have integral optimal solutions for any induced subgraph H of G and any $\mathbf{w} \in \mathbb{Z}^{V(H)}$ (and hence a combinatorial min-max relation follows).

Theorem 1.5 gives a structural description of all kernel ideal and kernel Mengerian digraphs in terms of forbidden structures. We point out that our characterization is actually a counterpart in the polyhedral case of the aforementioned Richardson's theorem (see Theorem 1.1), which can be rephrased as: Let G be a digraph. Then each subgraph of G has a kernel if and only if G contains no odd cycle. Interestingly, despite extensive research, a good characterization of all digraphs G such that each induced subgraph of G contains a kernel has yet to be found (see, for instance, [6, 9]). It is because kernels are usually not so well-behaved that these combinatorial objects are still surrounded by mystery. We close this section with two more open problems, which are intimately related to Theorem 1.5.

Problem 1.4. *Characterize all digraphs G such that $\sigma(H)$ defines an integral polytope for each induced subgraph H of G .*

Problem 1.5. *Characterize all digraphs G such that $\sigma(H)$ is TDI for each induced subgraph H of G .*

2 Complexity

Recall that every even digraph has a kernel; Richardson's proof [15] actually yields a polynomial-time algorithm for finding such a kernel. Let us now show that the MWKP is *NP*-hard even for some very special class of bipartite digraphs.

Proof of Theorem 1.2. Obviously, the kernel problem in our consideration is in *NP*. To prove the assertion, it suffices to reduce the planar 3-SATISFIABILITY problem (P3SAT) [14, 10] to this problem. Let $U = \{u_1, u_2, \dots, u_n\}$ be the set of variables, let $\mathcal{C} = \{C_1, C_2, \dots, C_m\}$ be the set of clauses in an arbitrary instance of the P3SAT in *CNF*, and let H be the bipartite graph with vertex set $U \cup \mathcal{C}$ such that $u_i C_j$ is an edge of H if and only if $u_i \in C_j$ or $\bar{u}_i \in C_j$ for $1 \leq i \leq n$ and $1 \leq j \leq m$. (Without loss of generality, we assume that no clause in \mathcal{C} contains both u_i and \bar{u}_i for any i .) By definition, H admits a plane embedding. For convenience, we view H as a plane graph hereafter, and use $e_i^1, e_i^2, \dots, e_i^{d_i}$ to denote all the edges incident with u_i in clockwise order in H for each i . Note that d_i equals the total number of occurrences of u_i and \bar{u}_i in \mathcal{C} .

Our objective is to construct a planar bipartite digraph G with maximum degree at most three so that there exists a kernel in G with at most $18m$ vertices if and only if \mathcal{C} is satisfiable. We can obtain the desired G from H by blowing up its vertices as described below:

- (1) Replace each vertex u_i in H by a truth-setting component T_i with vertex set $\cup_{t=1}^{d_i} (\{a_{i,t}^1, a_{i,t}^2, \dots, a_{i,t}^8\} \cup \{b_{i,t}^1, b_{i,t}^2, b_{i,t}^3, b_{i,t}^4\})$, such that T_i contains four (directed) paths $a_{i,t}^3 a_{i,t}^2 a_{i,t}^1$, $a_{i,t}^3 a_{i,t}^4 a_{i,t}^5$, $a_{i,t}^7 a_{i,t}^6 a_{i,t}^5$, $a_{i,t}^7 a_{i,t}^8 a_{i,t+1}^1$, two (directed) cycles $b_{i,t}^k b_{i,t}^{k+1} b_{i,t}^k$ for $k = 1, 3$, and two additional arcs $(a_{i,t}^1, b_{i,t}^1)$ and $(a_{i,t}^5, b_{i,t}^3)$, for each $t = 1, 2, \dots, d_i$, where d_i is the degree of u_i in H and $a_{i,d_i+1}^1 = a_{i,1}^1$. (So T_i has $12d_i$ vertices and $14d_i$ arcs in total; see Figure 2. Note that $\cup_{t=1}^{d_i} \{a_{i,t}^1, a_{i,t}^2, \dots, a_{i,t}^8\}$ induces an undirected cycle in the underlying graph of T_i .)
- (2) Replace each vertex C_j in H by a satisfaction-testing component S_j with vertex set $\{x_j, y_j^1, y_j^2, y_j^3, z_j^1, z_j^2, z_j^3\}$, such that $x_j y_j^k z_j^k$ is a path for $k = 1, 2, 3$. (So S_j has 7 vertices and 6 arcs in total; see Figure 2.)
- (3) For each edge $u_i C_j$ in H , we have $u_i \in C_j$ or $\bar{u}_i \in C_j$. Let r_j^1, r_j^2 , and r_j^3 denote the three literals in C_j throughout, and suppose $u_i C_j = e_i^t$ for some t with $1 \leq t \leq d_i$. If $u_i = r_j^k$, then add an arc from z_j^k to $a_{i,t}^2$; if $\bar{u}_i = r_j^k$, then add an arc from z_j^k to $a_{i,t}^6$.

The construction of G is completed. It is easy to see that this construction can be accomplished in polynomial time, G has $43m$ vertices and $51m$ edges (because $\sum_{i=1}^n d_i = 3m$), and G is a planar bipartite digraph with maximum degree at most three.




Figure 2: The truth-setting component T_i and satisfaction-testing component S_j .

Let us show that G has a kernel with at most $18m$ vertices if and only if \mathcal{C} is satisfiable.

Sufficiency. Suppose that $\tau : U \rightarrow \{\text{true}, \text{false}\}$ is a satisfying truth assignment for \mathcal{C} . We construct a vertex subset K of G as follows. For each variable u_i , if $\tau(u_i) = \text{true}$, then $\cup_{t=1}^{d_i} (\{a_{i,t}^2, a_{i,t}^5, a_{i,t}^8\} \cup \{b_{i,t}^1, b_{i,t}^4\}) \subseteq K$; if $\tau(u_i) = \text{false}$, then $\cup_{t=1}^{d_i} (\{a_{i,t}^1, a_{i,t}^4, a_{i,t}^6\} \cup \{b_{i,t}^2, b_{i,t}^3\}) \subseteq K$. For each clause $C_j \in \mathcal{C}$ and $k = 1, 2, 3$, if $\tau(r_j^k) = \text{true}$, then $y_j^k \in K$; if $\tau(r_j^k) = \text{false}$, then $z_j^k \in K$. This completes the construction of K . Clearly, $|K| = 18m$ (recall that $\sum_{i=1}^n d_i = 3m$), K is a stable set, and $K \cap V(T_i)$ is a kernel of T_i for $1 \leq i \leq n$. Since for each $j = 1, 2, \dots, m$, there exists at least one k with $\tau(r_j^k) = \text{true}$ and thus $y_j^k \in K$, and since (x_j, y_j^k) is an arc, we deduce that K is a kernel of G .

Necessity. Suppose G has a kernel K with at most $18m$ vertices. Then

- (4) $|K \cap \{b_{i,t}^k, b_{i,t}^{k+1}\}| = 1$ for $1 \leq i \leq n$, $1 \leq t \leq d_i$, and $k = 1, 3$.

From the structure of G , it can be seen that

- (5) for each $j = 1, 2, \dots, m$, we have $|K \cap V(S_j)| \geq 3$ and equality holds only when $y_j^{h_j} \in K$ for at least one subscript h_j with $1 \leq h_j \leq 3$.

We propose to show that

- (6) for each $i = 1, 2, \dots, n$, we have $|K \cap V(T_i)| \geq 5d_i$ and equality holds only when $K \cap V(T_i) = \bigcup_{t=1}^{d_i} (\{a_{i,t}^2, a_{i,t}^5, a_{i,t}^8\} \cup \{b_{i,t}^1, b_{i,t}^4\})$ or $K \cap V(T_i) = \bigcup_{t=1}^{d_i} (\{a_{i,t}^1, a_{i,t}^4, a_{i,t}^6\} \cup \{b_{i,t}^2, b_{i,t}^3\})$.

To this end, set $K_{i,t} = K \cap \{a_{i,t}^1, a_{i,t}^2, \dots, a_{i,t}^8\}$. Observe that

- (7) $|K_{i,t}| \geq 3$ for each $t = 1, 2, \dots, d_i$, because $K \cap V(T_i)$ is a kernel of T_i .

Combining (4) and (7), we obtain

- (8) $|K \cap V(T_i)| = 2d_i + \sum_{t=1}^{d_i} |K_{i,t}| \geq 5d_i$.

From this inequality and (5), it follows that $18m \geq |K| \geq \sum_{i=1}^n 5d_i + 3m = 18m$. Therefore

- (9) $|K \cap V(T_i)| = 5d_i$ for $1 \leq i \leq n$, and $|K \cap V(S_j)| = 3$ for $1 \leq j \leq m$.

The first equality in (9) and (8) in turn imply that

- (10) $|K_{i,t}| = 3$ for each $t = 1, 2, \dots, d_i$.

Let us consider the case when $a_{i,s}^1 \in K_{i,s}$ for some s with $1 \leq s \leq d_i$. Since $K \cap V(T_i)$ is a kernel of T_i and $K_{i,s} \subseteq K \cap V(T_i)$, it is a routine matter to check, using (10), that $K_{i,s} = \{a_{i,s}^1, a_{i,s}^4, a_{i,s}^6\}$. In view of the vertex $a_{i,s}^8$, we obtain $a_{i,s+1}^1 \in K_{i,s+1}$. Applying induction on the subscript t from s , we see that $K_{i,t} = \{a_{i,t}^1, a_{i,t}^4, a_{i,t}^6\}$ for all $t = 1, 2, \dots, d_i$. Hence $K \cap V(T_i) = \bigcup_{t=1}^{d_i} (\{a_{i,t}^1, a_{i,t}^4, a_{i,t}^6\} \cup \{b_{i,t}^2, b_{i,t}^3\})$.

So we assume that $a_{i,t}^1 \notin K_{i,t}$ for all t with $1 \leq t \leq d_i$. From (10), it follows instantly that $K_{i,t} = \{a_{i,t}^2, a_{i,t}^5, a_{i,t}^8\}$ for each t . Hence $K \cap V(T_i) = \bigcup_{t=1}^{d_i} (\{a_{i,t}^2, a_{i,t}^5, a_{i,t}^8\} \cup \{b_{i,t}^1, b_{i,t}^4\})$. Therefore (6) holds.

Let us now define a truth assignment $\tau : U \rightarrow \{true, false\}$ by setting $\tau(u_i) = true$ if $K \cap V(T_i) = \bigcup_{t=1}^{d_i} (\{a_{i,t}^2, a_{i,t}^5, a_{i,t}^8\} \cup \{b_{i,t}^1, b_{i,t}^4\})$ and setting $\tau(u_i) = false$ if $K \cap V(T_i) = \bigcup_{t=1}^{d_i} (\{a_{i,t}^1, a_{i,t}^4, a_{i,t}^6\} \cup \{b_{i,t}^2, b_{i,t}^3\})$, for $i = 1, 2, \dots, n$. It remains to verify that each clause C_j is satisfied by τ . Indeed, by (9), the subscript h_j as specified in (5) exists for each $j = 1, 2, \dots, m$, which implies that G has an arc from $z_j^{h_j}$ to $K \cap V(T_i)$ for some i . From the definition of τ , we deduce that $\tau(r_j^{h_j}) = true$. Therefore C_j is satisfied by τ for all j , completing the proof. ■

3 Obstructions

In this section we show that odd cycles, gears, and rings are all obstructions for a digraph to be kernel ideal.

Lemma 3.1. *Let G be a digraph that contains an odd cycle as a subgraph. Then G is not kernel ideal.*

Proof. Let C be an odd cycle in G and let $\hat{C} = (V, A)$ be the subgraph of G induced by all vertices in C . Consider $\mathbf{w} \in \mathbb{Z}^V$ and $\mathbf{a} \in \mathbb{Q}^V$ such that $w(v) = -1$ and $a(v) = 1/2$ for each $v \in V$. Clearly $\mathbf{a} \in FK(\hat{C})$, so $FK(\hat{C}) \neq \emptyset$. Since C is an odd cycle, each kernel of \hat{C} (if any) has size at most $(|V| - 1)/2$. Thus $\mathbf{w}^T \mathbf{a} = -|V|/2 < -(|V| - 1)/2 \leq \mathbf{w}^T \mathbf{x}$ for any incidence vector \mathbf{x} of a kernel in \hat{C} . From (1.6) we see that $FK(\hat{C}) \neq K(\hat{C})$, and hence G is not kernel ideal. ■

The remainder of this section is devoted to establishing the following statement.

Theorem 3.2. *Let G be a digraph that contains a gear or a ring as a subgraph. Then G is not kernel ideal.*

We break the proof into a series of observations. Recall the definition, a gear H is an even digraph (with no parallel arcs) obtained from the vertex-disjoint union of a circuit C and k cycles O_1, O_2, \dots, O_k by performing precisely one of the following three operations with respect to each sink t_i of C for $1 \leq i \leq k$:

- identify t_i with a vertex on O_i ;
- add a directed path P_i from t_i to O_i ;
- split t_i into two vertices t_i^1 and t_i^2 (each of them is adjacent with one arc on C) and identify t_i^1 with a vertex on O_i and t_i^2 with another vertex on O_i .

Let s_1, s_2, \dots, s_k be all the sources of C , which we call the *distinguished vertices* of H . Renaming the subscripts if necessary, we assume that $s_1, t_1, s_2, t_2, \dots, s_k, t_k$ occur on C in order if we traverse C from s_1 in the clockwise direction.

Lemma 3.3. *Let $H' = (V, A)$ be a digraph that contains a gear H as a spanning subgraph. If $d_{H'}^+(s_1) = 2$ (see the above description), then H' is not kernel ideal.*

Proof. Let Λ be the set of all subscripts i such that the above third operation has been applied with respect to t_i in the construction of H , and let Q_i be the segment of O_i from t_i^1 to t_i^2 for each $i \in \Lambda$. Then the arcs on C and on Q_i for all $i \in \Lambda$ form a circuit in H , denoted by D . Depending on the parity of $|V(D)|$, we consider two cases.

Case 1. $|V(D)|$ is odd.

In this case, define $\mathbf{w} \in \mathbb{Z}^V$ and $\mathbf{a} \in \mathbb{R}^V$ by

- $w(v) = -1$ if $v \in V(D)$ and $w(v) = 0$ otherwise;
- $a(v) = 1/2$ for each $v \in V$.

Clearly, $\mathbf{a} \in FK(H')$. Since $|V(D)|$ is odd, each kernel of H' (if any) contains at most $(|V(D)| - 1)/2$ vertices from D . Thus $\mathbf{w}^T \mathbf{a} = -|V(D)|/2 < -(|V(D)| - 1)/2 \leq \mathbf{w}^T \mathbf{x}$ for any incidence vector \mathbf{x} of a kernel in H' . From (1.6) we see that $FK(H') \neq K(H')$, and hence H' is not kernel ideal.

Case 2. $|V(D)|$ is even.

In this case, let u_1 and u_2 be the two out-neighbors of s_1 on D . Define $\mathbf{w} \in \mathbb{Z}^V$ and $\mathbf{a} \in \mathbb{R}^V$ by

- $w(s_1) = 1$, $w(u_i) = 0$ for $i = 1, 2$, $w(v) = -1$ if $v \in V(D) \setminus \{s_1, u_1, u_2\}$, and $w(v) = 0$ if $v \notin V(D)$;
- $a(s_1) = 0$ and $a(v) = 1/2$ for each $v \in V \setminus s_1$;

Clearly, $\mathbf{a} \in FK(H')$. Let \mathbf{x} be the incidence vector of an arbitrary kernel in H' (if any). If $x(u_1)$ or $x(u_2)$ is 1, then $x(V(D) \setminus \{s_1, u_1, u_2\}) \leq (|V(D)| - 4)/2$ by the stability of the kernel; if $x(u_1) = x(u_2) = 0$, then $x(s_1) = 1$ (as $d_{H'}^+(s_1) = 2$) and $x(V(D) \setminus \{s_1, u_1, u_2\}) \leq (|V(D)| - 2)/2$. In either case, we have $\mathbf{w}^T \mathbf{a} = -(|V(D)| - 3)/2 < -(|V(D)| - 4)/2 \leq \mathbf{w}^T \mathbf{x}$, which implies that $FK(H') \neq K(H')$, and hence H' is not kernel ideal. \blacksquare

Let H be a digraph obtained from a directed even cycle C by adding a new vertex v and some arcs between v and C so that no parallel arc is created. We call H a *wheel* if there are at

least three arcs from v to C . Note that there might be arcs from C to v in H . We call v the *hub* and C the *rim* of H .

Lemma 3.4. *Let $H = (V, A)$ be a wheel (see the above description). Then H is not kernel ideal.*

Proof. Let $C = u_1 u_2 \dots u_{2n} u_1$, and let $u_{i_1}, u_{i_2}, \dots, u_{i_k}$ be all the out-neighbors of v , where $1 \leq i_1 < i_2 < \dots < i_k \leq 2n$. We proceed by considering two cases.

Case 1. $|i_{j+1} - i_j|$ is odd for some j with $1 \leq j \leq k-1$.

In this case, let D be the circuit formed by $C[u_{i_j}, u_{i_{j+1}}]$ and two arcs (v, u_{i_j}) and $(v, u_{i_{j+1}})$. We define $\mathbf{w} \in \mathbb{Z}^V$ and $\mathbf{a} \in \mathbb{R}^V$ as follows:

- $w(u) = -1$ if $u \in V(D)$ and $w(u) = 0$ otherwise;
- $a(u) = 1/2$ for each $u \in V$.

Clearly, $\mathbf{a} \in FK(H)$. Since $|V(D)|$ is odd, each kernel of H (if any) contains at most $(|V(D)| - 1)/2$ vertices from D . Thus $\mathbf{w}^T \mathbf{a} = -|V(D)|/2 < -(|V(D)| - 1)/2 \leq \mathbf{w}^T \mathbf{x}$ for any incidence vector \mathbf{x} of a kernel in H . So $FK(H) \neq K(H)$, and hence H is not kernel ideal.

Case 2. $|i_{j+1} - i_j|$ is even for each j with $1 \leq j \leq k-1$.

In this case, we may assume that i_j is odd for each j (renaming subscripts if necessary). Define $\mathbf{w} \in \mathbb{Z}^V$ and $\mathbf{a} \in \mathbb{R}^V$ by

- $w(v) = k$, $w(u_{i_j}) = 0$ for $1 \leq j \leq k$, and $w(u) = -1$ if $u \in V(C) \setminus N_H^+(v)$; and
- $a(v) = 0$ and $a(u) = 1/2$ for each $u \in V \setminus v$.

Clearly, $\mathbf{a} \in FK(H)$. Let \mathbf{x} be the incidence vector of an arbitrary kernel U in H . If $x(u_{i_j}) = 1$ for some j with $1 \leq j \leq k$, then $U = \{u_1, u_3, \dots, u_{2n-1}\}$ by the hypothesis of the present case. So $\mathbf{w}^T \mathbf{x} = -n + k$. If $x(u_{i_j}) = 0$ for each j with $1 \leq j \leq k$, then $v \in U$ and $U \cap V(C) \subseteq \{u_2, u_4, \dots, u_{2n}\}$. So $\mathbf{w}^T \mathbf{x} \geq -n + k$. In either case, we have $\mathbf{w}^T \mathbf{x} \geq -n + k$. Hence $\mathbf{w}^T \mathbf{a} = -(2n - k)/2 = -n + k/2 < \mathbf{w}^T \mathbf{x}$, which implies that $FK(H) \neq K(H)$, and hence H is not kernel ideal. \blacksquare

Recall the definition, an odd ring H is obtained from a directed cycle C by adding an s_i - t_i path P_i for $i = 1, 2$, such that

- s_1, t_1, s_2, t_2 occur on C in order when we traverse C in its direction from s_1 ;
- $s_i \neq t_i$ (but possibly $s_i = t_{3-i}$) for $i = 1, 2$;
- $P_1(s_1, t_1)$, $P_2(s_2, t_2)$, and C are pairwise vertex-disjoint;
- $P_1 \cup P_2 \cup C$ is even and contains no parallel arcs; and
- $|C[s_1, s_2]|$ is odd.

Lemma 3.5. *Let $H = (V, A)$ be a ring. Then H is not kernel ideal.*

Proof. Since H contains no odd cycles, $|C[s_i, t_i]|$ and $|P_i|$ have the same parity for $i = 1, 2$. So there is a one-to-one correspondence between the kernels of H and those of C , and hence H has precisely two kernels X_1 and X_2 which form the unique bipartition of H , where $s_i \in X_i$ for $i = 1, 2$. Let $\mathbf{x} \in \mathbb{R}^V$ be defined as follows:

- $x(v) = 1/2$ for any vertex v outside $C(s_1, s_2] \cup P_1(s_1, t_1]$;
- $x(v) = 3/4$ for any vertex v in $V(C(s_1, s_2] \cup P_1(s_1, t_1]) \cap X_1$;
- $x(v) = 1/4$ for any vertex v in $V(C(s_1, s_2] \cup P_1(s_1, t_1]) \cap X_2$.

Clearly, $x \in FK(H)$; however, it cannot be expressed a convex combination of the incidence vectors of X_1 and X_2 , which implies that $FK(H) \neq K(H)$, so H is not kernel ideal. ■

Lemma 3.6. *Let $H' = (V, A')$ be a digraph obtained from a ring $H = (V, A)$ by adding an arc that is not parallel to any arc in H . Then H' contains an odd cycle, or a gear, or a ring with at most $|V| - 1$ vertices.*

Proof. We assume that H' contains no odd cycle, otherwise we are done. Since H is strongly connected, so is H' and hence it is bipartite by (1.1).

Recall the definition, H is obtained from a cycle C by adding an s_i - t_i path P_i for $i = 1, 2$ with the properties as described above the preceding lemma. Set $P_3 = C[s_1, t_1]$ and $P_4 = C[s_2, t_2]$. Obviously, P_1 and P_3 (similarly, P_2 and P_4) are in the same position in H . Let $e = (u, v)$ be the arc in $A' \setminus A$. There are now five cases, corresponding to possible locations of u and v .

Case 1. $u \in C[t_1, s_2]$ and $v \in C[t_2, s_1]$.

If $u = s_2$, let i be a subscript in $\{2, 4\}$ for which $P_i(s_2, t_2)$ contains at least one vertex, and let Σ be obtained from H' by deleting all vertices in $P_i(s_2, t_2)$, then Σ is a ring with at most $|V| - 1$ vertices. If $u \neq s_2$, then $C[s_2, u] \cup P_2 \cup \{(u, v)\}$ is a gear.

Case 2. $\{u, v\} \subseteq P_1$.

If $v \in P_1(u, t_1]$, let Σ be obtained from H by replacing $P_1[u, v]$ with (u, v) , then Σ is a ring with at most $|V| - 1$ vertices. So we suppose $v \in P_1[s_1, u)$. Furthermore, $\{u, v\} \neq \{t_1, s_1\}$, for otherwise Case 1 occurs. Let Σ be obtained from H' by deleting all vertices in $P_1(u, t_1) \cup C(s_1, s_2)$ if $u \neq t_1$, and let Σ be $P_1 \cup P_3 \cup \{(u, v)\}$ if $u = t_1$. Then Σ is a gear in either subcase.

Case 3. $u \in P_1$ and $v \in P_3$.

In this case we may assume that $u \notin \{s_1, t_1\}$, for otherwise Case 2 occurs (with P_3 in place of P_1). Thus $C \cup P_1[u, t_1] \cup \{(u, v)\}$ is a gear.

Case 4. $u \in P_1(s_1, t_1)$ and $v \in P_2$.

Let C' be obtained from C by replacing P_4 with P_2 . Then $C' \cup P_1[u, t_1] \cup \{(u, v)\}$ is a gear.

Case 5. $\{u, v\} \subseteq C[t_1, s_2]$.

If $v \in C(u, s_2]$, let Σ be obtained from H by replacing $C[u, v]$ with (u, v) , then Σ is a ring with at most $|V| - 1$ vertices. If $v \in C[t_1, u)$, then $P_1 \cup C[s_1, u] \cup \{(u, v)\}$ is a gear.

Each of the remaining cases is a mirror image of one case listed above. ■

Proof of Theorem 3.2. Let H be a gear or a ring in G with the smallest number of vertices, and let \hat{H} be the subgraph of G induced by all vertices in H . We may assume that \hat{H} is kernel ideal, otherwise we are done. By Lemma 3.5, \hat{H} is not a ring. Thus Lemma 3.6 and Lemma 3.1 allow us to further assume that H is a gear. Recall the structural description of H above Lemma 3.3; using this lemma, we obtain $d_{\hat{H}}^+(s_1) \geq 3$. So \hat{H} has an arc (s_1, v) outside the circuit C . Depending on the structure of H , we distinguish between two cases.

Case 1. H has precisely one distinguished vertex.

By assumption, $H \cup \{(s_1, v)\}$ contains no gear with fewer vertices than H . Consequently, H is obtained from the cycle O_1 by adding two arcs (s_1, t_1^1) and (s_1, t_1^2) . As \hat{H} is not a wheel with hub s_1 and rim O_1 by Lemma 3.4, there exists an arc (p, q) outside O_1 yet connecting two

vertices on O_1 . For convenience, set $t_1^3 = v$. Swapping t_1^1 and t_1^2 if necessary, we assume that t_1^1, t_1^2, t_1^3 occur on O_1 in order if we traverse O_1 from t_1^1 in the clockwise direction.

Consider the subcase when $\{p, q\} \subseteq O_1[t_1^1, t_1^2]$. Let Σ be obtained from H by replacing $O_1[p, q]$ with the arc (p, q) if $q \in O_1(p, t_1^2]$, let Σ be $O_1[t_1^1, t_1^2] \cup \{(p, q), (s_1, t_1^1), (s_1, t_1^2)\}$ if $p = t_1^2$, and let Σ be $O_1[t_1^3, p] \cup \{(p, q), (s_1, t_1^1), (s_1, t_1^3)\}$ if $p \neq t_1^2$ and $q \in O_1[t_1^1, p)$. Then Σ is a gear in G with fewer vertices than H , a contradiction.

So symmetry allows us to assume that one of p and q is on $O_1(t_1^1, t_1^2)$ and the other is on $O_1(t_1^2, t_1^3)$. Let Σ be obtained from $H \cup \{(s_1, t_1^3)\}$ by deleting all vertices on $O_1(p, q)$ if $p \in O_1(t_1^1, t_1^2)$, and let Σ be $O_1[t_1^1, p] \cup \{(p, q), (s_1, t_1^1), (s_1, t_1^2)\}$ otherwise. Then Σ is a gear in G with fewer vertices than H , again a contradiction.

Case 2. H has at least two distinguished vertices.

Recall the construction of H and the three operations performed with respect to each sink t_i , set $P_i = \emptyset$ if the first and the third operations are applied, and set $t_i^1 = t_i^2 = t_i$ if the first and second operations are applied. Renaming the subscripts if necessary, we assume that $s_1, t_1^1, t_1^2, s_2, t_2^1, t_2^2, \dots, s_k, t_k^1, t_k^2$ occur on C in order if we traverse C from s_1 in the clockwise direction. Moreover, let $Q_i = t_i$ if the first and second operations are applied and let Q_i be the segment of O_i from t_i^1 to t_i^2 if the third operations are applied. Then the arcs on C and on all these Q_i form a circuit in H , denoted by D . Given two vertices p and q on D , we use $D[p, q]$ to denote the segment of D from p to q in the clockwise direction.

Let us now construct a subgraph Σ of $H \cup \{(s_1, v)\}$ as follows.

When $v \in D[s_i, t_i^1]$ for some i with $1 \leq i \leq k$, let H' be the digraph obtained from $H \cup \{(s_1, v)\}$ by deleting all vertices on $D(s_1, v)$, and let Σ be the component of H' containing s_1 .

When $v \in D[t_i^2, s_{i+1})$ for some i with $1 \leq i \leq k$, where $s_{k+1} = s_1$, let H' be the digraph obtained from $H \cup \{(s_1, v)\}$ by deleting all vertices on $D(v, s_1)$, and let Σ be the component of H' containing s_1 .

When $v \in P_i \cup O_i \setminus \{t_i^1, t_i^2\}$ for some i with $1 \leq i \leq k$, let R stand for one of the segments $D(s_1, t_i^1)$ and $D(t_i^2, s_1)$ that contains a distinguished vertex s_j with $2 \leq j \leq k$, let H' be the digraph obtained from $H \cup \{(s_1, v)\}$ by deleting all vertices on R , and let Σ be the component of H' containing s_1 .

It is easy to see that Σ is a gear in G with fewer vertices than H in every subcase; this contradiction completes the proof of the present lemma. \blacksquare

4 Structures

Let Q, P_1, P_2 be three directed paths such that the ends s_i, t_i of P_i are both on Q for $i = 1, 2$ and that $P_1(s_1, t_1)$, $P_2(s_2, t_2)$, and Q are pairwise vertex-disjoint. We say that P_1 and P_2 *cross* with respect to Q if s_1, t_1, s_2, t_2 are distinct and each segment of Q (possibly Q is a cycle) between s_1 and t_1 contains precisely one of s_2 and t_2 .

For convenience, we call a digraph G *permissible* hereafter if G contains no subgraph isomorphic to an odd cycle, a gear, or a ring. With this notion, Theorem 1.5 amounts to saying that a digraph is kernel ideal (also kernel Mengerian) if and only if it is permissible. The following theorem aims to give a construction of strongly connected components of permissible digraphs, and exhibit some important properties enjoyed by them, where a cycle C in G is called *induced*

if no arc outside C in G connects two vertices on C . Roughly speaking, every strongly connected permissible digraph admits an ear decomposition (see, for instance, [3]) such that, except the initial cycle ear, each ear (cycle or path) is a child of a unique parent ear. Furthermore, each such ear is subject to strict restrictions on the locations of its ends.

Theorem 4.1. *Let $G = (V, A)$ be a strongly connected permissible digraph, where $|V| \geq 2$. Then G has a rooted tree-like structure (see Figure 3 for an illustration); that is, there exists a rooted tree T , such that*

- (i) *the root r of T corresponds to an (arbitrarily given) induced cycle P_r of G ;*
- (ii) *each $v \in V(T) \setminus r$ corresponds to an s_v - t_v path P_v (possibly $s_v = t_v$), such that if u is the parent of v in T , then both s_v and t_v are on P_u , and at least one of s_v and t_v is on $P_u(s_u, t_u)$ if $u \neq r$;*
- (iii) *P_r and $P_v(s_v, t_v)$ for all $v \in V(T) \setminus r$ are pairwise vertex-disjoint, and G is the union of P_v for all $v \in V(T)$;*
- (iv) *for any $v \in V(T) \setminus r$ and any child x of v , if $t_x \in P_v(s_x, t_v]$, then $s_x = s_v$;*
- (v) *for any $v \in V(T) \setminus r$ and any child x of v , if $t_x \in P_v[s_v, s_x)$, then $t_x \neq s_v$. Also, $s_x \neq t_v$ if $s_v \neq t_v$;*
- (vi) *for any $v \in V(T)$ and any two children x, y of v , paths P_x and P_y do not cross with respect to P_v ;*
- (vii) *for any $v \in V(T) \setminus r$ and any two children x, y of v , if $s_x = s_v$, then $s_y \in P_v[s_v, t_x)$;*
- (viii) *for any $v \in V(T) \setminus r$ and any two children x, y of v , if $t_x \in P_v[s_v, s_x)$ and $t_y \in P_v[s_v, s_y]$, then $s_y \notin P_v[t_x, s_x)$;*
- (ix) *for any two children x, y of r , if $s_x \neq t_x$ and s_x, t_x, t_y, s_y occur on P_r in order when we traverse P_r in its direction from s_x , then $s_y = s_x$; and*
- (x) *for any two children x, y of r , if $s_x \neq t_x$ and s_x, t_y, s_y, t_x occur on P_r in order when we traverse P_r in its direction from s_x , then precisely one of the following three cases occurs:*
 - *both s_y and t_y are on $P_r(s_x, t_x)$;*
 - *$s_x = s_y = t_y$;*
 - *$s_y = t_x, t_y = s_x$, and $|P_r[s_x, s_y]|$ is even.*

The following two lemmas will be used repeatedly in the proof of this theorem.

Lemma 4.2. *Let H be obtained from a directed cycle C by adding an s_1 - t_1 path P_1 (possibly $s_1 = t_1$) and an s_2 - t_2 path P_2 , such that*

- *$P_1(s_1, t_1)$, $P_2(s_2, t_2)$, and C are pairwise vertex-disjoint;*
- *$P_1 \cup P_2 \cup C$ is even and contains no parallel arcs; and*
- *s_1 and t_1 are on C , and one of s_2 and t_2 is on $P_1(s_1, t_1)$, the other is in $C \setminus \{s_1, t_1\}$.*

Then H contains a gear.

Proof. Symmetry allows us to assume that s_2 is on $P_1(s_1, t_1)$ if $s_1 = t_1$. Let F be the digraph obtained from H by first deleting all arcs on $P_1[s_1, s_2]$ if s_2 is on $P_1(s_1, t_1)$, by first deleting all arcs on $C[s_1, s_2]$ if s_2 is on $C(s_1, t_1)$, and by first deleting all arcs on $C[s_2, s_1]$ if s_2 is on $C(t_1, s_1)$, and then deleting all resulting isolated vertices (if any). Then F is a gear in each case. ■




Figure 3: A rooted tree-like structure.

Lemma 4.3. *Let H be obtained from a directed cycle C by adding an s_i - t_i path P_i for $i = 1, 2$, where s_i and t_i are two distinct vertices on C , such that*

- $P_1(s_1, t_1)$, $P_2(s_2, t_2)$, and C are pairwise vertex-disjoint;
- $P_1 \cup P_2 \cup C$ is even and contains no parallel arcs; and
- P_1 and P_2 cross with respect to C .

Then H contains a gear.

Proof. Swapping the subscripts if necessary, we may assume that s_1, s_2, t_1, t_2 occur on C in order if we traverse C in its direction. Thus the digraph obtained from H by first deleting all arcs on $C[s_1, s_2]$ and then deleting all resulting isolated vertices (if any) is a gear. ■

Proof of Theorem 4.1. We shall first construct a rooted tree T and paths P_v for all $v \in V(T)$ with properties (i)-(iii), and then demonstrate that they enjoy properties (iv)-(x) as well.

In the initialization step, we take an arbitrary induced cycle P_r in G (which is even), and construct a rooted tree T consisting of the root r only, which corresponds to P_r . At a general step, suppose we have constructed a rooted tree T and paths P_v for all $v \in V(T)$ with the properties as specified in (i), (ii), and (iii). If G is already the union of P_v for all $v \in V(T)$, then we are done. Otherwise, we augment T by adding a new vertex v and construct P_v with the desired properties.

Let us make some observations about the current T and P_v for $v \in V(T)$ before describing the construction procedure. Throughout we use R_v to denote the path from r to each vertex v in T .

(1) For each $v \in V(T)$, the union $\cup_{z \in V(R_v)} P_z$ is strongly connected and hence has a cycle containing P_v .

To justify this, we apply induction on $|R_v|$. If $|R_v| = 0$; that is, $v = r$, then (1) is trivial because P_r is a cycle. So we proceed to the induction step. Let u be the parent of v in T . By induction hypothesis, $\cup_{z \in V(R_u)} P_z$ is strongly connected. Since P_v is a path with ends in $\cup_{z \in V(R_u)} P_z$, it is a routine matter to check that $\cup_{z \in V(R_v)} P_z$ is also strongly connected. Thus (1) holds.

(2) For each $v \in V(T)$, if G has a path Q whose one end t is on $P_v(s_v, t_v)$, the other end s is on P_u but outside P_v for some ancestor u of v in T , and vertices in $V(Q) \setminus \{s, t\}$ are all outside $\cup_{z \in V(R_v)} P_z$, then G contains a gear.

We prove (2) by induction on $|V(T)|$. Since the statement follows instantly from Lemma 4.2 when $|V(T)| = 2$, we proceed to the induction step. Let a be the parent of v . We may assume that

(3) $s \notin P_a$ and $s_v \neq t_v$.

Otherwise, $s \in P_a$ or $s_v = t_v$. By (1), P_a is contained in a cycle C in $\cup_{z \in V(R_a)} P_z$. If $s \in P_a$, then $s \in C$; set $L = \emptyset$ in this case. If $s_v = t_v$ while $s \notin P_a$ then, since $\cup_{z \in V(R_a)} P_z$ is strongly connected and since s_v is on $P_a(s_a, t_a)$ (recall property (ii)), there exists a directed path L from C to s in $\cup_{z \in V(R_a)} P_z$, such that $L \cup Q$ is a directed path in G which has only two ends in $C \cup P_v$ and has one end in $C \setminus s_v$. Applying Lemma 4.2 with respect to P_v , $L \cup Q$, and C , we deduce that G contains a gear, so (3) holds.

(4) s_v is on $P_a[t_v, t_a]$.

Otherwise, s_v is on $P_a[s_a, t_v)$. Let $P_{v'}$ denote the path obtained from P_a by replacing $P_a[s_v, t_v]$ with P_v , let $R_{v'}$ be the path obtained from R_v by contracting the edge va into a single vertex v' , and let T' be the rooted tree consisting of $R_{v'}$ only. By (3), we have $s \notin P_{v'}$. From the induction hypothesis on T' , it follows that G contains a gear, so (4) is justified.

(5) t is the origin of Q .

Otherwise, t is the terminus of Q . By (4), $P_v \cup P_a$ has a cycle C containing P_v . Since $\cup_{z \in V(R_a)} P_z$ is strongly connected, it contains a directed path L from s to C . Thus Q , L , and C form a gear in G . This proves (5).

Let Q' stand for the path $P_v[s_v, t] \cup Q$. Then Q' is a directed path from s_v to s whose internal vertices are all outside $\cup_{z \in V(R_a)} P_z$. If $s_v \notin \{s_a, t_a\}$, then (3) and the induction hypothesis with respect to the path Q' and rooted tree $T' = R_a$ imply that G contains a gear. So we may assume that $s_v \in \{s_a, t_a\}$. In view of (3) and (4), we have $s_v = t_a$. Hence t_v is on $P_a(s_a, t_a)$ by property (ii). Since $\cup_{z \in V(R_a)} P_z$ is strongly connected, it has a cycle C containing P_a and a directed path L from s to C , such that $Q \cup L$ is a directed path in G which has only one end in C . Thus $P_v[t, t_v]$, $Q \cup L$, and C form a gear in G , completing the proof of (2).

(6) Suppose G has a path Q whose ends s and t (possibly $s = t$) are the only vertices of Q in $\cup_{z \in V(T)} P_z$. Let v be a vertex in T such that at least one of s and t is on $P_v(s_v, t_v)$ and that no descendant of v in T has this property, where $P_v(s_v, t_v) = P_r$ if $v = r$. Then both s and t are on P_v .

To justify this, assume the contrary: t is on $P_v(s_v, t_v)$ while s is outside P_v (renaming the ends of Q if necessary). By (2), we obtain

(7) s is not on P_z for any ancestor z of v in T .

Observe that

(8) s is not on P_z for any descendant z of v in T either.

Otherwise, from the hypothesis on v , we deduce that $s \in \{s_z, t_z\}$. So s is on P_a , where a is the parent of z in T . Once again by the hypothesis on v , we have $s \in \{s_a, t_a\}$ if $a \neq v$. Let us proceed in this way and eventually we see that s is on P_v , this contradiction justifies (8).

From (7), (8), and property (ii), we deduce that T has a vertex u such that s is on $P_u(s_u, t_u)$ and that neither vertex in $\{u, v\}$ is a descendant of the other. Symmetry allows us to assume

that t is the terminus of Q . Let C be a directed cycle containing P_v in $\cup_{z \in V(R_v)} P_z$ and let a be the parent of v in T . Since $\cup_{z \in V(R_a \cup R_u)} P_z$ is also strongly connected, it contains a directed path L from s to C . Thus Q , L , and C form a gear in G . Therefore (6) is established.

Now we are ready to describe the construction procedure. Suppose G is not the union of P_v for all vertices v in the current T . Then the strong connectedness of G guarantees the existence of a path Q as specified in (6). Let v be as defined in (6). We augment T by adding a vertex z and an edge vz and set $P_z = Q$. The process is repeated until G is the union of P_v for all $v \in V(T)$.

It remains to show that the rooted tree T and paths P_v enjoy all the desired properties. In our proofs of (iv)-(viii), we reserve the symbol u for the parent of the vertex v specified in the statements, and the symbol C_a for a cycle containing P_a in $\cup_{z \in V(R_a)} P_z$ (see (1)) for each vertex a on T .

(iv) If $s_x \neq s_v$, then $P_x \cup P_v[s_x, t_v] \cup C_u$ would be a gear; this contradiction justifies (iv).

(v) Recall (ii), if one of s_x and t_x is in $\{s_v, t_v\}$, then the other is on $P_v(s_v, t_v)$. Set $H = P_x \cup P_v[s_x, t_v] \cup C_u$ if $t_x = s_v$, and set $H = P_x \cup P_v \cup C_u[s_v, t_v]$ if $s_x = t_v \neq s_v$. Then H is a gear in either case and hence (v) holds.

(vi) Applying Lemma 4.3 to $C_v \cup P_x \cup P_y$, we deduce instantly that P_x and P_y do not cross with respect to P_v .

(vii) Assume the contrary: $s_y \notin P_v[s_v, t_x]$. By (iv), $t_y \notin P_v(s_v, t_v)$. So $t_y \in P_v(s_v, s_y]$ (see (v)). Set $H = P_v[s_v, t_x] \cup P_x \cup P_y$ if $t_y \in P_v(s_v, t_x)$, and set $H = P_v[s_v, s_y] \cup P_x \cup P_y$ if $t_y \in P_v[t_x, s_y]$. Notice that, by (vi), if $t_y \in P_v(s_v, t_x)$, then $s_y = t_x$, and that if $s_v = t_v$, then $s_y \neq t_v$ by the assumption. Consequently, H is a gear in either case, a contradiction. So (vii) is justified.

(viii) Assume the contrary: $s_y \in P_v[t_x, s_x]$. Note that if $s_y \neq t_x$, then $t_y \in P_v[t_x, s_y]$ by (vi). Set $H = P_x \cup C_v[s_x, s_y] \cup P_y$ if $s_y \neq t_x$, and set $H = P_x \cup C_v[s_x, t_x] \cup P_y$ if $s_y = t_x$. By (v), we have $t_x \neq s_v$. Besides, (v) implies $t_y \neq s_v$ when $t_x = s_y \neq t_y$. It follows that H is a gear in either case; this contradiction proves (viii).

(ix) If $s_y \neq s_x$, then $P_r[s_x, s_y] \cup P_x \cup P_y$ would be a gear, a contradiction.

(x) If $t_x = s_y$ but $s_x \neq t_y$, then $H = P_x \cup P_y \cup P_r[s_x, s_y]$ would be a gear. If $s_x = t_y$, $t_x = s_y$, but $P_r[s_x, s_y]$ is odd, then $P_r \cup P_x \cup P_y$ would be a ring. Using symmetry, it is easy to see that precisely one of the three stated cases can occur, completing the proof. ■

Given a digraph G , we use G^s to denote the digraph obtained from G by contracting each strongly connected component of G into a single vertex, such that the number of arcs between any two vertices in G^s equals the number of arcs between the corresponding strongly connected components of G . We call G^s the *skeleton* of G . Clearly, G^s contains no directed cycles but it may contain parallel arcs. Recall that a digraph is called *weakly connected* if its underlying graph is connected.

Theorem 4.4. *Let G be a weakly connected digraph with no gear. Suppose each sink of G^s corresponds to a strongly connected component of G with at least two vertices. Then the underlying graph of G^s is a tree. In particular, there is at most one arc between any two strongly connected components of G .*

Proof. Since G^s is acyclic, we have

(1) G^s contains a path from each vertex u to some sink of G^s .

Suppose on the contrary that the underlying graph of G^s is not a tree. Then G^s contains at least one circuit; let C be such a circuit with the smallest number of sinks (possibly C is formed by two parallel arcs). Notice that

(2) there is no path P from a sink of C to some other vertex on C in G^s , such that the internal vertices of P are all outside C .

Otherwise, it is a routine matter to check that $C \cup P$ contains either a directed cycle or a circuit with fewer sinks than C , this contradiction justifies (2).

Let t_1, t_2, \dots, t_k be all the sinks of C . By (1), G^s contains a path P_i from t_i to some sink u_i of G^s , for each $i = 1, 2, \dots, k$.

(3) C and $P_i \setminus t_i$, $i = 1, 2, \dots, k$, are pairwise vertex-disjoint.

Indeed, from (2) we see that $P_i \setminus t_i$ is vertex-disjoint from C for $1 \leq i \leq k$. Also, P_i and P_j are vertex-disjoint whenever $i \neq j$, for otherwise let x be a common vertex of P_i and P_j . Then $C \cup P_i[t_i, x] \cup P_j[t_j, x]$ would contain a circuit with fewer sinks than C , a contradiction. So (3) holds.

We propose to show that a gear in G can be obtained from $C \cup (\cup_{i=1}^k P_i)$ by blowing up its vertices.

To this end, let Ω_u denote the strongly connected component of G corresponding to each vertex u in G^s . For each vertex v of C (resp. each degree-two vertex v in $\cup_{i=1}^k P_i$), let e_v^1 and e_v^2 denote the two arcs of C (resp. of $\cup_{i=1}^k P_i$) incident with v , where e_v^1 enters v if v is not a source of C . We also view e_v^1 and e_v^2 as two arcs in G ; in this situation, we use v^j to denote the end of e_v^j in Ω_v for $j = 1, 2$.

For each vertex v on C that is not a sink of C and for each degree-two vertex v on $\cup_{i=1}^k P_i$, let Q_v be a path from v^1 to v^2 in Ω_v . Let Q be the union of Q_v 's for all vertices v in $C \cup (\cup_{i=1}^k P_i)$.

For each sink t_i on C with $t_i = u_i$ (that is, P_i consists of a single vertex), let R_i^j be a path in Ω_{t_i} from t_i^j to t_i^{2-j} for $j = 1, 2$, such that at least one of R_i^1 and R_i^2 has positive length (such paths exist because Ω_{t_i} contains at least two vertices by hypothesis). Let O_i be the directed cycle in $R_i^1 \cup R_i^2$ containing t_i^1 and set $Q_i = O_i \cup R_i^2$.

For each sink t_i on C with $t_i \neq u_i$ (that is, P_i contains at least two vertices), we assume for simplicity that t_i (resp. u_i) is the end of the starting (resp. ending) arc of P_i contained in Ω_{t_i} (resp. Ω_{u_i}). Let R_i^1 be a path in Ω_{t_i} from t_i^1 to t_i , and let R_i^2 be a path in Ω_{t_i} from t_i^2 to R_i^1 . Let O_i be a cycle passing through u_i in Ω_{u_i} and set $Q_i = O_i \cup R_i^1 \cup R_i^2$.

From (3) it follows instantly that the arcs of G in $C \cup Q \cup (\cup_{i=1}^k P_i \cup Q_i)$ form a gear in G , this contradiction implies that the underlying graph of G^s must be a tree. ■

5 Reductions

In this section we introduce a series of reduction operations, under which the integrality properties enjoyed by the fractional kernel polytope and its defining system are preserved. Our reductions are all based on the assumption that every permissible digraph with fewer vertices or with fewer arcs than the input digraph G is kernel ideal and kernel Mengerian (see Theorem 1.5).

For any real number α , we shall use $[\alpha]^+$ as a shorthand for $\max\{0, \alpha\}$. We shall also use $\phi_v(\mathbf{y}, \mathbf{z})$ to denote the left-hand side of (1.8) throughout.

Reduction 1

Instance: A digraph $G = (V, A)$ with a sink a and a weight function $\mathbf{w} \in \mathbb{Z}^V$.

Description: Let $G' = (V', A')$ be the digraph obtained from G by deleting all vertices in $\{a\} \cup N_G^-(a)$ and let \mathbf{w}' be the restriction of \mathbf{w} to $\mathbb{Z}^{V'}$. Suppose \mathbf{x}' and $(\mathbf{y}', \mathbf{z}')$ are integral optimal solutions to $\mathbb{P}(G', \mathbf{w}')$ and $\mathbb{D}(G', \mathbf{w}')$, respectively. Let $e_i = (b_i, a)$ for $1 \leq i \leq k$ be all the arcs incident with a , and set $\alpha_i = z'(N_G^-(b_i) \cap V')$ for $1 \leq i \leq k$, where $z'(\emptyset) = 0$. Define

- $x^*(a) = 1$ and $x^*(b_i) = 0$ for $1 \leq i \leq k$;
- $y^*(e_1) = [-w(a)]^+ + [-w(b_1)]^+ + \alpha_1$, $y^*(e_i) = [-w(b_i)]^+ + \alpha_i$ for $2 \leq i \leq k$, and $y^*(e) = 0$ for all $e \notin A' \cup \{e_1, e_2, \dots, e_k\}$;
- $z^*(a) = w(a) + \sum_{i=1}^k y^*(e_i)$ and $z^*(b_i) = 0$ for $1 \leq i \leq k$; and
- $x^*(v) = x'(v)$ for all $v \in V'$, $y^*(e) = y'(e)$ for all $e \in A'$, and $z^*(v) = z'(v)$ for all $v \in V'$.

Lemma 5.1. *For Reduction 1, the following statements hold:*

- (i) \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively; and
- (ii) the optimal value of $\mathbb{P}(G, \mathbf{w})$ equals $w(a)$ plus that of $\mathbb{P}(G', \mathbf{w}')$.

Proof. Clearly, \mathbf{x}^* satisfies (1.2) – (1.4). Since $w(a) + [-w(a)]^+ = [w(a)]^+ \geq 0$, we have $z^*(a) \geq 0$. It is then easy to see that $(\mathbf{y}^*, \mathbf{z}^*)$ satisfies (1.8) – (1.10). So \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral feasible solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively. By the above definition, we have $\mathbf{w}^T \mathbf{x}^* = w(a) + (\mathbf{w}')^T \mathbf{x}' = w(a) - y'(A') + z'(V') = z^*(a) - \sum_{i=1}^k y^*(e_i) - y'(A') + z'(V') = -y^*(A) + z^*(V)$. From the LP-duality theorem, we thus deduce that \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively. Hence both (i) and (ii) are established. ■

Reduction 2

Instance: A permissible digraph $G = (V, A)$ with a weight function $\mathbf{w} \in \mathbb{Z}^V$, and three distinct vertices a, b, c of G , such that a and c are nonadjacent and have no common in-neighbor, $N_G^+(a) = \{b\}$, and $N_G^+(b) = \{c\}$ (arc (c, b) may exist).

Description: Let $G' = (V', A')$ be the digraph obtained from G by first deleting the arc (a, b) and then identifying a with c (we still use c to denote the resulting vertex), and let $w'(c) = w(a) + w(c)$ and $w'(v) = w(v)$ if $v \in V' \setminus c$. Suppose \mathbf{x}' and $(\mathbf{y}', \mathbf{z}')$ are integral optimal solutions to $\mathbb{P}(G', \mathbf{w}')$ and $\mathbb{D}(G', \mathbf{w}')$, respectively. Set $\alpha = \sum_{v \in N_G^-(a)} y'((v, a))$, $\beta = z'(N_G^-(a))$, and $\gamma = w(a) + \alpha - \beta$.

- $x^*(a) = x'(c)$, and $x^*(v) = x'(v)$ for every $v \in V \setminus a$;
- $y^*((a, b)) = [-\gamma]^+$, $y^*((b, c)) = y'((b, c)) + [\gamma]^+$, and $y^*(e) = y'(e)$ for every $e \in A \setminus \{(a, b), (b, c)\}$; and
- $z^*(a) = [\gamma]^+$, $z^*(b) = z'(b) + [-\gamma]^+$, and $z^*(v) = z'(v)$ for every $v \in V \setminus \{a, b\}$.

Lemma 5.2. *For Reduction 2, the following statements hold:*

- (i) G' is also permissible;

- (ii) \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively; and
(iii) the optimal value of $\mathbb{P}(G, \mathbf{w})$ equals that of $\mathbb{P}(G', \mathbf{w}')$.

Proof. Suppose on the contrary that G' contains a subgraph Σ which is isomorphic to an odd cycle, or a gear, or a ring. It is then a routine matter to check that the subgraph of G induced by all arcs in $\Sigma \cup \{(a, b), (b, c)\}$ would contain an odd cycle, or a gear, or a ring, contradicting the hypothesis that G is permissible. So (i) holds.

To prove the remaining two statements, observe that

- $x^*(a) + x^*(b) = x^*(b) + x^*(c) = x'(b) + x'(c) \leq 1$;
- $\phi_a(\mathbf{y}^*, \mathbf{z}^*) = -\sum_{a \in e} y^*(e) + z^*(a) + z^*(N_G^-(a)) = -\alpha - y^*((a, b)) + [\gamma]^+ + \beta = -\alpha - [-\gamma]^+ + [\gamma]^+ + \beta = -\alpha + \gamma + \beta = w(a)$;
- $\phi_b(\mathbf{y}^*, \mathbf{z}^*) = -\sum_{b \in e} y^*(e) + z^*(b) + z^*(N_G^-(b)) = (-y^*((a, b)) - \sum_{b \in e} y'(e) - [\gamma]^+) + (z'(b) + [-\gamma]^+) + z^*(a) + z'(N_{G'}^-(b)) = -\sum_{b \in e} y'(e) + z'(b) + z'(N_{G'}^-(b)) \leq w'(b) = w(b)$; and
- $\phi_c(\mathbf{y}^*, \mathbf{z}^*) = -\sum_{c \in e} y^*(e) + z^*(c) + z^*(N_G^-(c)) \leq (-\sum_{c \in e} y'(e) + \alpha - [\gamma]^+) + z'(c) + z'(N_{G'}^-(c)) - \beta + [-\gamma]^+ \leq w'(c) + \alpha - \beta - \gamma = w(a) + w(c) + \alpha - \beta - \gamma = w(c)$.

Combining the above observations, we conclude that \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral feasible solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively. By the above definition, we have $\mathbf{w}^T \mathbf{x}^* = (\mathbf{w}')^T \mathbf{x}' = -y'(A') + z'(V') = -y^*(A) + y^*((a, b)) + [\gamma]^+ + z^*(V) - z^*(a) - [-\gamma]^+ = -y^*(A) + z^*(V)$. From the LP-duality theorem, we further deduce that \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively. Hence both (ii) and (iii) follow. \blacksquare

Reduction 3

Instance: A permissible digraph $G = (V, A)$ with a weight function $\mathbf{w} \in \mathbb{Z}^V$, and an induced cycle $abcd a$ of length four in G with $N_G^+(b) = \{c\}$ and $N_G^+(d) = \{a\}$. (We remark that $N_G^-(a) \cap N_G^-(c) = \emptyset$, otherwise such a common neighbor together with $abcd a$ would form a gear.)

Description: Let $G' = (V', A')$ be the digraph obtained from G by identifying a with c (we still use a denote the resulting vertex), and let $w'(a) = w(a) + w(c)$ and $w'(v) = w(v)$ if $v \in V' \setminus a$. Suppose \mathbf{x}' and $(\mathbf{y}', \mathbf{z}')$ are integral optimal solutions to $\mathbb{P}(G', \mathbf{w}')$ and $\mathbb{D}(G', \mathbf{w}')$, respectively. We may assume that $z'(a) = 0$ (otherwise, replace $z'(b)$ by $z'(b) + z'(a)$ and replace $z'(a)$ by 0). As $\phi_a(\mathbf{y}', \mathbf{z}') \leq w(a) + w(c)$, the above remark implies that one of the inequalities $-\sum_{a \in e} y'(e) + z'(N_G^-(a)) \leq w(a)$ and $-\sum_{c \in e} y'(e) + z'(N_G^-(c)) \leq w(c)$ holds; symmetry allows us to assume the former. Set $\alpha = [-\sum_{c \in e} y'(e) + z'(N_G^-(c)) - w(c)]^+$. Define

- $x^*(a) = x^*(c) = x'(a)$, and $x^*(v) = x'(v)$ for each $v \in V \setminus \{a, c\}$;
- $y^*((c, d)) = y'((c, d)) + \alpha$, and $y^*(e) = y'(e)$ for each $e \in A \setminus (c, d)$;
- $z^*(a) = z^*(c) = 0$, $z^*(d) = z'(d) + \alpha$, and $z^*(v) = z'(v)$ for each $v \in V \setminus \{a, c, d\}$.

Lemma 5.3. *For Reduction 3, the following statements hold:*

- (i) G' is also permissible;
(ii) \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively; and
(iii) the optimal value of $\mathbb{P}(G, \mathbf{w})$ equals that of $\mathbb{P}(G', \mathbf{w}')$.

Proof. Suppose on the contrary that G' contains a subgraph Σ which is isomorphic to an odd cycle, or a gear, or a ring. It is then a routine matter to check that the subgraph of G induced by all arcs in $\Sigma \cup \{(a, b), (b, c), (c, d), (d, a)\}$ would contain an odd cycle, or a gear, or a ring, contradicting the hypothesis that G is permissible. So (i) holds.

Clearly \mathbf{x}^* is an integral feasible solution to $\mathbb{P}(G, \mathbf{w})$. Let us now show that $\phi_v(\mathbf{y}^*, \mathbf{z}^*) \leq w(v)$ for all $v \in V$. Since $N_G^+(b) = \{c\}$ and $N_G^+(d) = \{a\}$, we have $\phi_v(\mathbf{y}^*, \mathbf{z}^*) = \phi_v(\mathbf{y}', \mathbf{z}') \leq w(v)$ for each $v \in V \setminus \{a, c, d\}$. From the above definition, we see that

- $\phi_d(\mathbf{y}^*, \mathbf{z}^*) = \phi_d(\mathbf{y}', \mathbf{z}') \leq w(d)$;
- $\phi_c(\mathbf{y}^*, \mathbf{z}^*) = -\sum_{e \in c} y'(e) + z'(N_G^-(c)) - \alpha = [-\sum_{e \in c} y'(e) + z'(N_G^-(c)) - w(c)] - \alpha + w(c) \leq w(c)$; and
- $\phi_a(\mathbf{y}^*, \mathbf{z}^*) = -\sum_{e \in a} y'(e) + z'(N_G^-(a)) + \alpha \leq \max\{w(a), -\sum_{e \in a} y'(e) + z'(N_G^-(a)) + (-\sum_{e \in c} y'(e) + z'(N_G^-(c)) - w(c))\} = \max\{w(a), \phi_a(\mathbf{y}', \mathbf{z}') - w(c)\} = w(a)$.

Combining the above observations, we conclude that $(\mathbf{y}^*, \mathbf{z}^*)$ is an integral feasible solution to $\mathbb{D}(G, \mathbf{w})$. Since $\mathbf{w}^T \mathbf{x}^* = (\mathbf{w}')^T \mathbf{x}' = -y'(A) + z'(V) = -y^*(A) + z^*(V)$, from the LP-duality theorem, we deduce that \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively. Hence both (ii) and (iii) are established. \blacksquare

Reduction 4

Instance: A permissible digraph $G = (V, A)$ with a weight function $\mathbf{w} \in \mathbb{Z}^V$, and a path abc of G with $(b, a) \in A$ and $N_G^+(a) = \{b\}$ (arc (c, b) may exist).

Lemma 5.4. $\mathbb{P}(G, \mathbf{w})$ has an integral optimal solution.

Proof. Let \mathbf{x} be an optimal solution to $\mathbb{P}(G, \mathbf{w})$. We propose to construct an integral optimal solution \mathbf{x}^* to $\mathbb{P}(G, \mathbf{w})$, starting from \mathbf{x} .

By (1.3) and (1.4), we have $x(a) + x(b) = 1$, which implies $\mathbf{x} \in FK(G \setminus (b, c))$. Since $G \setminus (b, c)$ is kernel ideal (recall the assumption at the beginning of this section), $\mathbf{x} = \sum_{i=1}^k c_i \mathbf{x}_i$, where \mathbf{x}_i is the incidence vector of a kernel U_i in $G \setminus (b, c)$, $c_i > 0$, for $1 \leq i \leq k$, and $\sum_{i=1}^k c_i = 1$. Renaming subscripts if necessary, we may assume that $\mathbf{w}^T \mathbf{x}_1 \leq \mathbf{w}^T \mathbf{x}_i$ for all $2 \leq i \leq k$. Thus $\{b, c\} \subseteq U_1$, for otherwise $\mathbf{x}^* = \mathbf{x}_1$ is as desired. It follows that $x_1(b) = x_1(c) = 1$.

Since $x(b) + x(c) \leq 1$ and $c_1 x_1(v) \leq x(v)$ for $v = b, c$, we obtain $c_1 \leq \min\{x(b), x(c)\} \leq 1/2$. Let α be the constant in $[c_1, 1/2]$ such that $\frac{1-\alpha}{1-c_1}(x(b) + x(c) - 2c_1) + 2\alpha = 1$, and set $\mathbf{x}' = \alpha \mathbf{x}_1 + \frac{1-\alpha}{1-c_1} \sum_{i=2}^k c_i \mathbf{x}_i$. Clearly, $\mathbf{x}' \in K(G \setminus (b, c))$ and $\mathbf{w}^T \mathbf{x}' \leq \mathbf{w}^T \mathbf{x}$. Since $x'(b) + x'(c) = \frac{1-\alpha}{1-c_1}(x(b) + x(c) - 2c_1) + 2\alpha = 1$, from the definition we deduce that $\mathbf{x}' \in FK(G \setminus (b, a))$. As $G \setminus (b, a)$ is kernel ideal, we further have $\mathbf{x}' \in K(G \setminus (b, a))$. Therefore there exists the incidence vector \mathbf{x}^* of a kernel U of $G \setminus (b, a)$ such that $\mathbf{w}^T \mathbf{x}^* \leq \mathbf{w}^T \mathbf{x}'$; this \mathbf{x}^* is as desired because U is also a kernel of G and $\mathbf{w}^T \mathbf{x}^* \leq \mathbf{w}^T \mathbf{x}$. \blacksquare

In the remainder of this reduction, we reserve the symbol \mathbf{x} for an integral optimal solution to $\mathbb{P}(G, \mathbf{w})$ and the pair (\mathbf{y}, \mathbf{z}) for an optimal solution to $\mathbb{P}(G, \mathbf{w})$. Set $\delta = y((b, c))$.

Description: Let $G' = (V, A \setminus (b, c))$, and let $w'(v) = w(v) + \lceil \delta \rceil$ if $v \in \{b, c\}$ and $w'(v) = w(v)$ if $v \in V \setminus \{b, c\}$. Suppose \mathbf{x}' and $(\mathbf{y}', \mathbf{z}')$ are integral optimal solutions to $\mathbb{P}(G', \mathbf{w}')$ and $\mathbb{D}(G', \mathbf{w}')$, respectively. In particular, set $\mathbf{x}' = \mathbf{x}$ if \mathbf{x} is also an optimal solution to $\mathbb{P}(G', \mathbf{w}')$. Since

$N_G^+(a) = \{b\}$, we may assume that $z'(b) = 0$ (otherwise we replace $z'(a)$ by $z'(a) + z'(b)$ and replace $z'(b)$ by 0). Define

- $\mathbf{x}^* = \mathbf{x}'$;
- $y^*((b, c)) = \lceil \delta \rceil$ and $y^*(e) = y'(e)$ for all $e \in A \setminus (b, c)$; and
- $\mathbf{z}^* = \mathbf{z}'$.

Lemma 5.5. *For Reduction 4, the following statements hold:*

- (i) G' is also permissible;
- (ii) \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively; and
- (iii) the optimal value of $\mathbb{P}(G, \mathbf{w})$ equals that of $\mathbb{P}(G', \mathbf{w}')$ minus $\lceil \delta \rceil$.

Proof. Statement (i) is trivial. Define $w''(v) = w(v) + \delta$ if $v \in \{b, c\}$ and $w''(v) = w(v)$ if $v \in V \setminus \{b, c\}$. Clearly, $\mathbf{w}' = \mathbf{w}''$ if δ is integral. Observe that \mathbf{x} and $(\mathbf{y}|_{A \setminus (b, c)}, \mathbf{z})$ are feasible solutions to $\mathbb{P}(G', \mathbf{w}'')$ and $\mathbb{D}(G', \mathbf{w}'')$, respectively, and satisfy the complementary slackness condition. So

(1) \mathbf{x} is also an optimal solution to $\mathbb{P}(G', \mathbf{w}'')$ and hence $(\mathbf{w}'')^T \mathbf{x}' \geq (\mathbf{w}'')^T \mathbf{x}$.

Let us first consider the case when \mathbf{x} is an optimal solution to $\mathbb{P}(G', \mathbf{w}')$. Now $\mathbf{x}^* = \mathbf{x}$ and $(\mathbf{w}')^T \mathbf{x} = -y'(A \setminus (b, c)) + z'(V)$. So $\phi_b(\mathbf{y}^*, \mathbf{z}^*) = \phi_b(\mathbf{y}', \mathbf{z}') - y^*((b, c)) \leq w'(b) - \lceil \delta \rceil = w(b)$. As $z'(b) = 0$, we have $\phi_c(\mathbf{y}^*, \mathbf{z}^*) = \phi_c(\mathbf{y}', \mathbf{z}') - y^*((b, c)) + z'(b) \leq w'(c) - \lceil \delta \rceil = w(c)$. From the feasibility of $(\mathbf{y}', \mathbf{z}')$ we deduce that $(\mathbf{y}^*, \mathbf{z}^*)$ is a feasible solution to $\mathbb{D}(G, \mathbf{w})$, with value $-y^*(A) + z^*(V) = -y^*((b, c)) - y'(A \setminus (b, c)) + z'(V) = (\mathbf{w}')^T \mathbf{x} - \lceil \delta \rceil$. Again, by the complementary slackness condition, $\delta(x(b) + x(c) - 1) = 0$, so either $x(b) + x(c) = 1$ or $\delta = x(b) = x(c) = 0$ (as \mathbf{x} is integral), and hence the equality $\mathbf{w}^T \mathbf{x} = (\mathbf{w}')^T \mathbf{x} - \lceil \delta \rceil$ holds in either subcase. Therefore $-y^*(A) + z^*(V) = \mathbf{w}^T \mathbf{x}$. From the LP-duality theorem, we deduce that $(\mathbf{y}^*, \mathbf{z}^*)$ is an integral optimal solution to $\mathbb{D}(G, \mathbf{w})$.

So we assume that \mathbf{x} is not an optimal solution to $\mathbb{P}(G', \mathbf{w}')$. Thus $\mathbf{w}' \neq \mathbf{w}''$ by (1). It follows that δ is not integral. By the complementary slackness condition, we get

(2) $x(b) + x(c) = 1$.

Let ω^* denote the optimal value of $\mathbb{P}(G, \mathbf{w})$ (which is an integer as \mathbf{x} is integral). From (2) we see that

(3) $(\mathbf{w}'')^T \mathbf{x} = \omega^* + \delta$ and $(\mathbf{w}')^T \mathbf{x} = \omega^* + \lceil \delta \rceil$.

Since \mathbf{x} is not an optimal solution to $\mathbb{P}(G', \mathbf{w}')$, we have $(\mathbf{w}')^T \mathbf{x}' < (\mathbf{w}')^T \mathbf{x} = \omega^* + \lceil \delta \rceil$. So

(4) $(\mathbf{w}')^T \mathbf{x}' \leq \omega^* + \lceil \delta \rceil - 1$.

From the definitions of \mathbf{w}' and \mathbf{w}'' , we see that $(\mathbf{w}')^T \mathbf{x}' \geq (\mathbf{w}'')^T \mathbf{x}'$. Using (1) and (3), we thus obtain $(\mathbf{w}')^T \mathbf{x}' \geq (\mathbf{w}'')^T \mathbf{x} = \omega^* + \delta$, contradicting (4). \blacksquare

Reduction 5

Instance: A permissible digraph $G = (V, A)$ with a weight function $\mathbf{w} \in \mathbb{Z}^V$, and three distinct vertices a, b, c of G , such that $N_G^+(a) = \{b\}$, $N_G^+(b) = \{a\}$, and $(c, b) \in A$, and that $G \setminus (c, b)$ is the union of two disjoint graphs $G_1 = (V_1, A_1)$ and $G \setminus V_1$, with $\{a, b\} \subsetneq V_1$ and $c \notin V_1$.

Description: Let $G_2 = (V_2, A_2)$ denote $G \setminus (V_1 \setminus \{a, b\})$, let α be the minimum weight of a kernel in $(G_2 \setminus \{a, b\}, \mathbf{w}|_{V_2 \setminus \{a, b\}})$, and let β be the minimum weight of a kernel in $(G_2 \setminus \{a, b, c\}, \mathbf{w}|_{V_2 \setminus \{a, b, c\}})$ if $V_2 \neq \{a, b, c\}$ and let $\beta = 0$ otherwise. Define $w_1(a) = w(a) + \alpha$, $w_1(b) = w(b) + \beta$,

and $w_1(v) = w(v)$ if $v \in V_1 \setminus \{a, b\}$, and define $w_2(a) = -\alpha$, $w_2(b) = -\beta$, and $w_2(v) = v$ if $v \in V_2 \setminus \{a, b\}$. Observe that the minimum weight of a kernel in (G_2, \mathbf{w}_2) is zero, and that each of a and b is contained in a minimum weighted kernel of (G_2, \mathbf{w}_2) . Suppose \mathbf{x}_i and $(\mathbf{y}_i, \mathbf{z}_i)$ are integral optimal solutions to $\mathbb{P}(G_i, \mathbf{w}_i)$ and $\mathbb{D}(G_i, \mathbf{w}_i)$, respectively, for $i = 1, 2$, such that $x_1(v) = x_2(v)$ for $v \in \{a, b\}$. (Given \mathbf{x}_1 , the existence of such an \mathbf{x}_2 is guaranteed by the preceding observation.) Define

- $x^*(v) = x_i(v)$ if $v \in V_i$ for $i = 1, 2$;
- $y^*(e) = y_1(e) + y_2(e)$ if $e \in \{(a, b), (b, a)\}$, and $y^*(e) = y_i(e)$ if $e \in A_i \setminus \{(a, b), (b, a)\}$ for $i = 1, 2$; and
- $z^*(v) = z_1(v) + z_2(v)$ if $v \in \{a, b\}$, and $z^*(v) = z_i(v)$ if $v \in V_i \setminus \{a, b\}$ for $i = 1, 2$.

Lemma 5.6. *For Reduction 5, the following statements hold:*

- (i) \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively; and
- (ii) the optimal value of $\mathbb{P}(G, \mathbf{w})$ equals that of $\mathbb{P}(G_1, \mathbf{w}_1)$.

Proof. (i) From its definition, we see that \mathbf{x}^* is a feasible solution to $\mathbb{P}(G, \mathbf{w})$. To establish the dual feasibility of $(\mathbf{y}^*, \mathbf{z}^*)$, it suffices to verify that $\phi_v(\mathbf{y}^*, \mathbf{z}^*) \leq w(v)$ for $v \in \{a, b\}$. Indeed, by definition, $\phi_v(\mathbf{y}^*, \mathbf{z}^*) = -\sum_{e \in e \in A} y^*(e) + z^*(a) + z^*(b) + z^*(N_G^-(v) \setminus \{a, b\}) = \sum_{i=1}^2 [-\sum_{v \in e \in A_i \setminus \{(a, b), (b, a)\}} y_i(e) - y_i((a, b)) - y_i((b, a)) + z_i(a) + z_i(b) + z_i(N_{G_i}^-(v) \setminus \{a, b\})] = \sum_{i=1}^2 [-\sum_{v \in e \in A_i} y_i(e) + z_i(v) + z_i(N_{G_i}^-(v))] \leq w_1(v) + w_2(v) = w(v)$, so $(\mathbf{y}^*, \mathbf{z}^*)$ is a feasible solution to $\mathbb{D}(G, \mathbf{w})$. As $w_1(v) + w_2(v) = w(v)$ and $x_1(v) = x_2(v)$ for $v \in \{a, b\}$, we have $\mathbf{w}^T \mathbf{x}^* = \sum_{i=1}^2 \mathbf{w}_i^T \mathbf{x}_i = \sum_{i=1}^2 [-y_i(A_i) + z_i(V_i)] = -y^*(A) + z^*(V)$. By the LP-duality theorem, \mathbf{x}^* and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral optimal solutions to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively.

(ii) Since the minimum weight of a kernel in (G_2, \mathbf{w}_2) is zero (see description above), $\mathbf{w}_2^T \mathbf{x}_2 = 0$. So $\mathbf{w}^T \mathbf{x}^* = \sum_{i=1}^2 \mathbf{w}_i^T \mathbf{x}_i = \mathbf{w}_1^T \mathbf{x}_1$. Hence the optimal value of $\mathbb{P}(G, \mathbf{w})$ equals that of $\mathbb{P}(G_1, \mathbf{w}_1)$. \blacksquare

Reduction 6

Instance: A permissible digraph $G = (V, A)$ with a weight function $\mathbf{w} \in \mathbb{Z}^V$, and three distinct vertices a, b, c of G , such that $N_G^+(a) = N_G^-(a) = \{b\}$, $N_G^+(b) = \{a\}$, and $N_G^-(b) = \{a, c\}$.

Throughout this reduction, $G' = (V', A')$ stands for the digraph arising from $G \setminus a$ by adding an arc (b, c) . Clearly, G' is permissible and hence is both kernel ideal and kernel Mengerian by the assumption at the beginning of this section. Let ω^* stand for the common optimal value of $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$. For each subgraph H of G , we use $\Gamma(H)$ to denote the set of all kernels in H , and use $\chi_U \in \{0, 1\}^{V(H)}$ to denote the incidence vector of each U in $\Gamma(H)$.

Lemma 5.7. $\mathbb{P}(G, \mathbf{w})$ has an integral optimal solution.

Proof. Let \mathbf{x} be an optimal solution to $\mathbb{P}(G, \mathbf{w})$ such that

- (1) $x(b)$ is as large as possible.

By (1.3) and (1.4), we have

- (2) $x(a) + x(b) = 1$.

Suppose for a contradiction that \mathbf{x} is not integral. Let us make some further observations about \mathbf{x} .

$$(3) \ x(b) + x(c) < 1.$$

Otherwise, $x(b) + x(c) = 1$ by (1.3). Thus $x(a) = x(c)$ by (2) and $\mathbf{x}|_{V'} \in FK(G')$. Let $w'(c) = w(a) + w(c)$ and $w'(v) = w(v)$ for every $v \in V' \setminus c$. As G' is kernel ideal, $\mathbb{P}(G', \mathbf{w}')$ admits an integral optimal solution \mathbf{x}' , with value $(\mathbf{w}')^T \mathbf{x}' \leq (\mathbf{w}')^T \mathbf{x}|_{V'} = \mathbf{w}^T \mathbf{x} - w(a)x(a) - w(c)x(c) + w'(c)x(c) = \mathbf{w}^T \mathbf{x} = \omega^*$. Define $\mathbf{x}^* \in \mathbb{Z}_+^V$ by $x^*(a) = x'(c)$ and $x^*(v) = x'(v)$ for every $v \in V'$. Clearly, \mathbf{x}^* is a feasible solution to $\mathbb{P}(G, \mathbf{w})$ with value $\mathbf{w}^T \mathbf{x}^* = (\mathbf{w}')^T \mathbf{x}' \leq \omega^*$. So \mathbf{x}^* is an integral optimal solution to $\mathbb{P}(G, \mathbf{w})$. Hence we may assume that (3) holds.

$$(4) \ 0 < x(a), x(b) < 1.$$

Otherwise, $x(u) = 1$ for some $u \in \{a, b\}$. Let $e \in \{(a, b), (b, a)\}$ be the arc leaving u . Then $\mathbf{x} \in FK(G \setminus e)$. Since $G \setminus e$ is kernel ideal, $\mathbb{P}(G \setminus e, \mathbf{w})$ has an integral optimal solution \mathbf{x}^* , for which $\mathbf{w}^T \mathbf{x}^* \leq \mathbf{w}^T \mathbf{x} = \omega^*$. Clearly, $\mathbf{x}^* \in K(G)$. So \mathbf{x}^* is an optimal integral solution to $\mathbb{P}(G, \mathbf{w})$. Hence we may assume (4).

$$(5) \ w(b) > w(a).$$

To justify this, let $\tilde{\mathbf{x}}$ be defined by $\tilde{x}(a) = x(c)$, $\tilde{x}(b) = 1 - x(c)$, and $\tilde{x}(v) = x(v)$ for each $v \in V \setminus \{a, b\}$. Then $\tilde{\mathbf{x}}$ is a feasible solution to $\mathbb{P}(G, \mathbf{x})$. From (1) and (3), we deduce that $\mathbf{w}^T \tilde{\mathbf{x}} > \mathbf{w}^T \mathbf{x}$, which implies $w(a)x(c) + w(b)(1 - x(c)) = w(a)\tilde{x}(a) + w(b)\tilde{x}(b) > w(a)x(a) + w(b)x(b)$, so $(1 - x(b) - x(c))w(b) > (x(a) - x(c))w(a)$ and hence $w(b) > w(a)$ by (2) and (3).

Let (\mathbf{y}, \mathbf{z}) be an optimal solution to $\mathbb{D}(G, \mathbf{w})$ with minimum $z(c)$. By the complementary slackness condition and (4), we obtain

$$(6) \ \phi_v(\mathbf{y}, \mathbf{z}) = w(v) \text{ for } v \in \{a, b\}.$$

$$(7) \ y((c, b)) = 0, z(c) = w(b) - w(a) > 0, \text{ and } x(c) + x(N_G^+(c)) = 1.$$

To justify this, note that $\phi_b(\mathbf{y}, \mathbf{z}) = \phi_a(\mathbf{y}, \mathbf{z}) - y((c, b)) + z(c)$. So $-y((c, b)) + z(c) = w(b) - w(a) > 0$ by (6) and (5). Hence $y((c, b)) = 0$ and $z(c) = w(b) - w(a) > 0$, for otherwise, let $(\bar{\mathbf{y}}, \bar{\mathbf{z}})$ be obtained from (\mathbf{y}, \mathbf{z}) by replacing $z(c)$ with $z(c) - y((c, b))$ and $y((c, b))$ with 0. Then $(\bar{\mathbf{y}}, \bar{\mathbf{z}})$ is also an optimal solution to $\mathbb{D}(G, \mathbf{w})$ with $\bar{z}(c) < z(c)$, contradicting the minimality assumption on (\mathbf{y}, \mathbf{z}) . Since $z(c) > 0$, from the complementary slackness condition we deduce that $x(c) + x(N_G^+(c)) = 1$. So (7) is established.

Let $\mathbf{x}' \in \mathbb{R}_+^{V'}$ be defined by $x'(b) = 1 - x(c)$ and $x'(v) = x(v)$ if $v \in V' \setminus b$. Then $\mathbf{x}' \in FK(G')$. Since G' is kernel ideal, $\mathbf{x}' = \sum_{U \in \Gamma(G')} r_U \chi_U$, where $\sum_{U \in \Gamma(G')} r_U = 1$ and $r_U \geq 0$ for all $U \in \Gamma(G')$. Set $\mathcal{D} = \{U \in \Gamma(G') : r_U > 0\}$. Clearly, \mathcal{D} is the disjoint union of the following three sets:

- $\mathcal{A} = \{U \in \mathcal{D} : b \in U, U \setminus b \in \Gamma(G' \setminus b)\};$
- $\mathcal{B} = \{U \in \mathcal{D} : b \in U, U \setminus b \notin \Gamma(G' \setminus b)\};$ and
- $\mathcal{C} = \{U \in \mathcal{D} : c \in U\}.$

We propose to show that

$$(8) \ \mathcal{A} \neq \emptyset \neq \mathcal{B}.$$

Indeed, by (7), we have $x(b) + x(c) + x(N_{G'}^+(c) \setminus b) = x(c) + x(N_G^+(c)) = 1$. Using (3), we obtain $x(N_{G'}^+(c) \setminus b) > 0$. So $x'(N_{G'}^+(c) \setminus b) > 0$ and hence $\mathcal{A} \neq \emptyset$. Suppose on the contrary that $\mathcal{B} = \emptyset$. Then $\mathbf{x}' = \sum_{U \in \mathcal{A}} r_U \chi_U + \sum_{U \in \mathcal{C}} r_U \chi_U$. Define $\mathbf{x}^* = \sum_{U \in \mathcal{A}} r_U \chi_{\{a\} \cup (U \setminus b)} + \sum_{U \in \mathcal{C}} r_U \chi_{\{a\} \cup U}$. From the definitions of \mathcal{A} and \mathcal{C} , we see that $\mathbf{x}^* \in FK(G)$. Note that $x^*(a) = x'(b) + x'(c) = 1 - x(c) + x(c) = 1$, $x^*(b) = 0$, and $x^*(c) = x'(c) = x(c)$. So $\mathbf{w}^T \mathbf{x}^* - \mathbf{w}^T \mathbf{x} = w(a)x^*(a) - w(a)x(a) - w(b)x(b) < w(a) - w(a)x(a) - w(a)x(b) = w(a) - w(a) = 0$, where the inequality follows from (5) and the first equality from (2). Thus $\mathbf{w}^T \mathbf{x}^* < \mathbf{w}^T \mathbf{x} = \omega^*$; this contradiction implies that $\mathcal{B} \neq \emptyset$.

Take A in \mathcal{A} and B in \mathcal{B} . From the definitions of \mathcal{A} and \mathcal{B} , we see that

(9) $b \in A \cap B$, $c \notin A \cup B$, and $A \cap (N_{G'}^+(c) \setminus b) \neq \emptyset = B \cap (N_{G'}^+(c) \setminus b)$.

Let $\theta = |A \cap (N_{G'}^+(c) \setminus b)|$. Then $\theta \geq 1$ by (9). Set $\alpha = 1$ if $w(A \setminus b) + \theta w(a) \geq w(B \setminus b) + \theta w(b)$ and $\alpha = 2$ otherwise. In view of (2)-(4), we can find a constant ϵ such that $0 < \epsilon < \min\{r_A, r_B\}$ and $0 < \theta\epsilon \leq \min\{x(b), 1 - x(b) - x(c)\} = \min\{1 - x(a), x(a) - x(c)\}$. Let $\mathbf{x}'' = \mathbf{x}' + \epsilon \cdot [(-1)^\alpha \chi_A + (-1)^{3-\alpha} \chi_B]$. It is clear from $\epsilon < \min\{r_A, r_B\}$ that \mathbf{x}'' remains to be a convex combination of incidence vectors of kernels of G' in \mathcal{D} . So $\mathbf{x}'' \in FK(G')$. Define $\mathbf{x}^* \in \mathbb{R}^V$ by $x^*(a) = x(a) + (-1)^\alpha \theta\epsilon$, $x^*(b) = x(b) + (-1)^{3-\alpha} \theta\epsilon$, and $x^*(v) = x''(v)$ for all $v \in V \setminus \{a, b\}$.

(10) $\mathbf{x}^* \in FK(G)$.

To justify this, note that $0 \leq x^*(a), x^*(b) \leq 1$ by the choice of ϵ , and that $x^*(c) = x''(c) = x'(c) = x(c)$ because $c \notin A \cup B$ by (9). Moreover, $x^*(a) + x^*(b) = x(a) + x(b) = 1$ by (2), and $x^*(b) + x^*(c) = x(b) + x(c) + (-1)^{3-\alpha} \theta\epsilon \leq 1$ by the choice of ϵ . Since $x''(N_{G'}^+(c) \setminus b) = x'(N_{G'}^+(c) \setminus b) + (-1)^\alpha \theta\epsilon$ by (9) and the definition of θ , we have $x^*(c) + x^*(N_G^+(c)) = x^*(b) + x^*(c) + x^*(N_G^+(c) \setminus b) = x(b) + (-1)^{3-\alpha} \theta\epsilon + x''(c) + x''(N_{G'}^+(c) \setminus b) = x(b) + (-1)^{3-\alpha} \theta\epsilon + x'(c) + x'(N_{G'}^+(c) \setminus b) + (-1)^\alpha \theta\epsilon = x(b) + x'(c) + x'(N_{G'}^+(c) \setminus b) = x(b) + x(c) + x(N_G^+(c) \setminus b) = x(c) + x(N_G^+(c)) = 1$ by (7). So (10) holds.

(11) $\mathbf{w}^T \mathbf{x}^* = \mathbf{w}^T \mathbf{x} + (-1)^\alpha \epsilon \cdot [w(A \setminus b) + \theta w(a) - w(B \setminus b) - \theta w(b)]$.

By direct computation, we obtain $\mathbf{w}^T \mathbf{x}^* = w(a)x^*(a) + w(b)x^*(b) + \sum_{v \in V \setminus \{a, b\}} w(v)x''(v) = w(a)[x(a) + (-1)^\alpha \theta\epsilon] + w(b)[x(b) + (-1)^{3-\alpha} \theta\epsilon] + [\sum_{v \in V \setminus \{a, b\}} w(v)x'(v) + (-1)^\alpha w(A \setminus b)\epsilon + (-1)^{3-\alpha} w(B \setminus b)\epsilon] = [w(a)x(a) + w(b)x(b) + \sum_{v \in V \setminus \{a, b\}} w(v)x'(v)] + (-1)^\alpha \epsilon \cdot [w(A \setminus b) + \theta w(a) - w(B \setminus b) - \theta w(b)] = \mathbf{w}^T \mathbf{x} + (-1)^\alpha \epsilon \cdot [w(A \setminus b) + \theta w(a) - w(B \setminus b) - \theta w(b)]$. So (11) is justified.

(12) $\alpha = 1$ and \mathbf{x}^* is also an optimal solution to $\mathbb{P}(G, \mathbf{w})$.

Suppose the contrary: $\alpha = 2$. From the definition of α , we see that $w(A \setminus b) + \theta w(a) < w(B \setminus b) + \theta w(b)$. Thus $\mathbf{w}^T \mathbf{x}^* < \mathbf{w}^T \mathbf{x} = \omega^*$ by (11), a contradiction (see (10)). So $\alpha = 1$ and hence $w(A \setminus b) + \theta w(a) \geq w(B \setminus b) + \theta w(b)$ by the definition of α . It follows from (11) that $\mathbf{w}^T \mathbf{x}^* \leq \mathbf{w}^T \mathbf{x}$. Therefore, by (10), \mathbf{x}^* is also an optimal solution to $\mathbb{P}(G, \mathbf{w})$.

From (12) we conclude that $x^*(b) = x(b) + \theta\epsilon > x(b)$. This contradiction to (1) proves the present lemma. \blacksquare

By Lemma 5.7, the common optimal value ω^* of $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$ is an integer. It remains to prove that $\mathbb{D}(G, \mathbf{w})$ admits an integral solution of value ω^* . To this end, we assume, throughout the remainder of this reduction, that

$$\text{each of } a \text{ and } b \text{ is contained in a kernel of } (G, \mathbf{w}) \text{ with total weight } \omega^*. \quad (5.1)$$

Otherwise, let $\kappa(u)$ be the minimum total weight of a kernel containing a vertex u in (G, \mathbf{w}) , and let $\tilde{\mathbf{w}}$ be obtained from \mathbf{w} by replacing $w(v)$ with $w(v) - (\kappa(v) - \omega^*)$ for $v = a$ and b . Since $\tilde{\mathbf{w}} \leq \mathbf{w}$, from (1.8)-(1.10) we see that every feasible solution to $\mathbb{D}(G, \tilde{\mathbf{w}})$ remains feasible to $\mathbb{D}(G, \mathbf{w})$. Hence we may assume that (5.1) holds. We reserve the symbol \mathbf{x}_v for an integral optimal solution to $\mathbb{P}(G, \mathbf{w})$ with $x_v(v) = 1$ for $v \in \{a, b\}$. It is clear that

$$x_b(a) = x_b(c) = 0 \text{ and } \mathbf{x}_b|_{V'} \in K(G'). \quad (5.2)$$

The reduction given below depends on whether $w(a) \geq w(b)$ or not.

Case 1. $w(a) \geq w(b)$.

Description: Let $\mathbf{w}' \in \mathbb{Z}^{V'}$ be defined by $w'(c) = w(a) + w(c)$ and $w'(v) = w(v)$ if $v \in V' \setminus c$. Suppose $(\mathbf{y}', \mathbf{z}')$ is an integral optimal solution to $\mathbb{D}(G', \mathbf{w}')$. We may assume that $y'((c, b)) = 0$ and $z'(c) = 0$ (otherwise, replace $y'((b, c))$ with $y'((b, c)) + y'((c, b))$, $y'((c, b))$ with 0, $z'(b)$ with $z'(b) + z'(c)$, and $z'(c)$ with 0). Let $\alpha = w(a) + y'((b, c)) - z'(b)$. Define

- $y^*((a, b)) = 0$, $y^*((b, a)) = y'((b, c))$, $y^*((c, b)) = \alpha$, and $y^*(e) = y'(e)$ for each $e \in A' \setminus \{(b, c), (c, b)\}$; and
- $z^*(a) = \alpha$ and $z^*(v) = z'(v)$ for each $v \in V \setminus a$.

Lemma 5.8. *For Reduction 6 in Case 1, the following statements hold:*

- (i) $(\mathbf{y}^*, \mathbf{z}^*)$ is an integral optimal solution to $\mathbb{D}(G, \mathbf{w})$; and
- (ii) the optimal value of $\mathbb{D}(G, \mathbf{w})$ equals that of $\mathbb{D}(G', \mathbf{w}')$.

Proof. Let us first show that

(1) the optimal value of $\mathbb{P}(G', \mathbf{w}')$ is ω^* . So $-y'(A') + z'(V') = \omega^*$ by duality.

Indeed, for any $\mathbf{x}'' \in K(G')$, let $x(a) = x''(c)$ and $x(v) = x''(v)$ if $v \in V'$. Then $\mathbf{x} \in K(G)$ and $\mathbf{w}^T \mathbf{x} = (\mathbf{w}')^T \mathbf{x}''$. Thus ω^* is a lower bound on the optimal value of $\mathbb{P}(G', \mathbf{w}')$. From (5.2) we see that $\mathbf{x}' = \mathbf{x}_b|_{V'}$ is feasible solution to $\mathbb{P}(G', \mathbf{w}')$. Since $(\mathbf{w}')^T \mathbf{x}' = \mathbf{w}^T \mathbf{x}_b = \omega^*$, statement (1) holds.

Observe that $-y^*(A) + z^*(V) = (-y'(A' \setminus \{(b, c), (c, b)\}) - y^*((a, b)) - y^*((b, a)) - y^*((c, b))) + (z'(V') + z^*(a)) = (-y'(A') - \alpha) + (z'(V') + \alpha) = -y'(A') + z'(V')$. Using (1), we obtain

(2) $-y^*(A) + z^*(V) = \omega^*$.

(3) $(\mathbf{y}^*, \mathbf{z}^*)$ is a feasible solution to $\mathbb{D}(G, \mathbf{w})$.

To justify this, note that $y'((c, b)) = z'(c) = 0$. Since $\phi_b(\mathbf{y}', \mathbf{z}') \leq w'(b) = w(b)$, we have $-y'((b, c)) + z'(b) \leq w(b) \leq w(a)$ by the assumption on Case 1. So $\alpha = w(a) + y'((b, c)) - z'(b) \geq 0$ and hence $\mathbf{y}^* \geq \mathbf{0}$ and $\mathbf{z}^* \geq \mathbf{0}$. Moreover,

- $\phi_a(\mathbf{y}^*, \mathbf{z}^*) = -y^*((a, b)) - y^*((b, a)) + z^*(a) + z^*(b) = -y'((b, c)) + \alpha + z'(b) = w(a)$;
- $\phi_b(\mathbf{y}^*, \mathbf{z}^*) = \phi_b(\mathbf{y}', \mathbf{z}') + y'((b, c)) + y'((c, b)) - y^*((a, b)) - y^*((b, a)) - y^*((c, b)) + z^*(a) = \phi_b(\mathbf{y}', \mathbf{z}') - y^*((c, b)) + z^*(a) = \phi_b(\mathbf{y}', \mathbf{z}') - \alpha + \alpha \leq w'(b) = w(b)$;
- $\phi_c(\mathbf{y}^*, \mathbf{z}^*) = \phi_c(\mathbf{y}', \mathbf{z}') + y'((b, c)) + y'((c, b)) - z'(b) - y^*((c, b)) \leq w'(c) + y'((b, c)) - z'(b) - \alpha = w(a) + w(c) + y'((b, c)) - z'(b) - \alpha = w(c)$; and
- $\phi_v(\mathbf{y}^*, \mathbf{z}^*) = \phi_v(\mathbf{y}', \mathbf{z}') \leq w'(v) = w(v)$ for every $v \in V \setminus \{a, b, c\}$.

Combining the above observations, we conclude that $(\mathbf{y}^*, \mathbf{z}^*)$ is a feasible solution to $\mathbb{D}(G, \mathbf{w})$.

Since ω^* is the optimal value of $\mathbb{D}(G, \mathbf{w})$, (i) and (ii) follow instantly from (1)-(3). \blacksquare

Case 2. $w(a) < w(b)$.

Description: Set $\alpha = w(b) - w(a)$. Let $\mathbf{w}' \in \mathbb{Z}^{V'}$ be defined by $w'(b) = 0$, $w'(v) = w(v) - \alpha$ if $v \in \{c\} \cup N_G^+(c) \setminus b$, and $w'(v) = w(v)$ if $v \in V' \setminus (\{c\} \cup N_G^+(c))$. Suppose $(\mathbf{y}', \mathbf{z}')$ is an integral optimal solution to $\mathbb{D}(G', \mathbf{w}')$. We may assume that $y'((c, b)) = 0$ (otherwise, replace $y'((c, b))$ with $y'(c, b) + y'(b, c)$ and $y'((c, b))$ with 0), and assume that $z'(b) = z'(c) = 0$ (otherwise, replace $y'((b, c))$ with $y'((b, c)) - z'(b) - z'(c)$, and replace both $z'(b)$ and $z'(c)$ with 0. Then $\phi_b(\mathbf{y}', \mathbf{z}') = -y'((b, c)) + z'(b) + z'(c) \leq w'(b) = 0$ and $\mathbf{y}' \geq \mathbf{0}$, $\mathbf{z}' \geq \mathbf{0}$ remain valid). Define

- $y^*((a, b)) = [-w(a)]^+$, $y^*((b, a)) = 0$, and $y^*(e) = y'(e)$ if $e \in A \setminus \{(a, b), (b, a)\}$;
- $z^*(a) = [w(a)]^+$, $z^*(c) = \alpha$, and $z^*(v) = z'(v)$ if $v \in V \setminus \{a, c\}$.

Lemma 5.9. *For Reduction 6 in Case 2, the following statements hold:*

- (i) $(\mathbf{y}^*, \mathbf{z}^*)$ is an integral optimal solution to $\mathbb{D}(G, \mathbf{w})$; and
- (ii) the optimal value of $\mathbb{D}(G, \mathbf{w})$ equals that of $\mathbb{D}(G', \mathbf{w}')$ plus $w(b)$.

Proof. Recall the definition of x_v for $v \in \{a, b\}$. Since $x_v(v) = 1$, by the complementary slackness condition we have

- (1) $\phi_v(\mathbf{y}, \mathbf{z}) = w(v)$ for each $v \in \{a, b\}$.

Let (\mathbf{y}, \mathbf{z}) be an optimal solution to $\mathbb{D}(G, \mathbf{w})$ with minimum $z(c)$. Then

- (2) $y((c, b)) = 0$ and $z(c) = \alpha > 0$.

To justify this, note that $w(b) = \phi_b(\mathbf{y}, \mathbf{z}) = \phi_a(\mathbf{y}, \mathbf{z}) - y((c, b)) + z(c) = w(a) - y((c, b)) + z(c)$ by (1). Hence $-y((c, b)) + z(c) = \alpha > 0$, which implies $y((c, b)) = 0$ and $z(c) = \alpha$, for otherwise, let $(\bar{\mathbf{y}}, \bar{\mathbf{z}})$ be obtained from (\mathbf{y}, \mathbf{z}) by replacing $y((c, b))$ with 0 and replacing $z(c)$ with $-y((c, b)) + z(c)$. Then $(\bar{\mathbf{y}}, \bar{\mathbf{z}})$ is also an optimal solution to $\mathbb{D}(G, \mathbf{w})$ with $\bar{z}(c) < z(c)$, contradicting the assumption on (\mathbf{y}, \mathbf{z}) . So (2) is established.

By (2) and the complementary slackness condition, $x_a(c) + x_a(N_G^+(c)) = 1$. It follows from $x_a(b) = 0$ that

- (3) $x_a(u) = 1$ and $x_a(\{c\} \cup N_G^+(c) \setminus u) = 0$ for some vertex $u \in \{c\} \cup N_G^+(c) \setminus b$.

We propose to show that

- (4) the optimal value of $\mathbb{D}(G', \mathbf{w}')$ is $\omega^* - w(b)$. So $-y'(A') + z'(V') = \omega^* - w(b)$.

To justify this, define $(\mathbf{y}'', \mathbf{z}'')$ by $y''((b, c)) = y''((c, b)) = 0$ and $y''(e) = y(e)$ for each $e \in A' \setminus \{(b, c), (c, b)\}$, and $z''(b) = z''(c) = 0$ and $z''(v) = z(v)$ for each $v \in V' \setminus \{b, c\}$. Observe that

- $\phi_b(\mathbf{y}'', \mathbf{z}'') = -y''((b, c)) - y''((c, b)) + z''(b) + z''(c) = 0 = w'(b)$;
- $\phi_c(\mathbf{y}'', \mathbf{z}'') = \phi_c(\mathbf{y}, \mathbf{z}) + y((c, b)) - z(c) - y''((b, c)) - y''((c, b)) + z''(c) \leq w(c) - \alpha = w'(c)$,

where the inequality follows from (2);

- $\phi_v(\mathbf{y}'', \mathbf{z}'') = \phi_v(\mathbf{y}, \mathbf{z}) - z(c) + z''(c) \leq w(v) - \alpha = w'(v)$ for each $v \in N_G^+(c) \setminus b$; and
- $\phi_v(\mathbf{y}'', \mathbf{z}'') = \phi_v(\mathbf{y}, \mathbf{z}) \leq w(v) = w'(v)$ for each $v \in V' \setminus (\{c\} \cup N_G^+(c))$.

Hence $(\mathbf{y}'', \mathbf{z}'')$ is a feasible solution to $\mathbb{D}(G', \mathbf{w}')$ with value $-y''(A') + z''(V') = -y(A) + z(V) - \phi_b(\mathbf{y}, \mathbf{z}) + \phi_b(\mathbf{y}'', \mathbf{z}'') = \omega^* - w(b)$ by (1).

Note that $\mathbf{x}_a|_{V' \setminus b} \in K(G' \setminus b)$. Let $x''(b) = 1$ if $x_a(c) = 0$, $x''(b) = 0$ if $x_a(c) = 1$, and $x''(v) = x_a(v)$ if $v \in V' \setminus b$. Clearly, \mathbf{x}'' is a feasible solution to $\mathbb{P}(G', \mathbf{w}')$. In view of (3), we have

$$\begin{aligned}
 (\mathbf{w}')^T \mathbf{x}'' &= \sum_{v \in V' \setminus (\{c\} \cup N_G^+(c))} w(v)x_a(v) + \sum_{v \in (\{c\} \cup N_G^+(c)) \setminus b} (w(v) - \alpha)x_a(v) \\
 &= \sum_{v \in V' \setminus (\{c\} \cup N_G^+(c))} w(v)x_a(v) + (w(u) - \alpha) \\
 &= \sum_{v \in V' \setminus b} w(v)x_a(v) - \alpha = \sum_{v \in V'} w(v)x_a(v) - \alpha = \omega^* - w(a) - \alpha = \omega^* - w(b).
 \end{aligned}$$

Since $-y''(A') + z''(V') = (\mathbf{w}')^T \mathbf{x}'' = \omega^* - w(b)$, we obtain (4) using the LP-duality theorem.

- (5) $(\mathbf{y}^*, \mathbf{z}^*)$ is a feasible solution to $\mathbb{D}(G, \mathbf{w})$.

To justify this, observe that

- $\phi_a(\mathbf{y}^*, \mathbf{z}^*) = -y^*((a, b)) - y^*((b, a)) + z^*(a) + z^*(b) = -[-w(a)]^+ + [w(a)]^+ + z'(b) = w(a)$;
- $\phi_b(\mathbf{y}^*, \mathbf{z}^*) = \phi_a(\mathbf{y}^*, \mathbf{z}^*) - y^*((c, b)) + z^*(c) \leq w(a) + \alpha = w(b)$;

- $\phi_c(\mathbf{y}^*, \mathbf{z}^*) = \phi_c(\mathbf{y}', \mathbf{z}') + y'((b, c)) - z'(c) + z^*(c) \leq w'(c) + \alpha = w(c)$;
- $\phi_v(\mathbf{y}^*, \mathbf{z}^*) = \phi_v(\mathbf{y}', \mathbf{z}') - z'(c) + z^*(c) \leq w'(v) + \alpha = w(v)$ for each $v \in N_G^+(c) \setminus b$; and
- $\phi_v(\mathbf{y}^*, \mathbf{z}^*) = \phi_v(\mathbf{y}', \mathbf{z}') \leq w'(v) = w(v)$ for each $v \in V' \setminus (\{c\} \cup N_G^+(c))$.

Therefore (5) holds.

$$(6) \quad -y^*(A) + z^*(V) = \omega^*.$$

Indeed, since $z'(c) = 0$ and $y'((b, c)) = 0$, we obtain $-y'(A' \setminus (b, c)) + z'(V' \setminus c) = \omega^* - w(b)$ by (4). It follows that $-y^*(A) + z^*(V) = -y'(A' \setminus (b, c)) - y^*((a, b)) - y^*((b, a)) + z'(V' \setminus c) + z^*(a) + z^*(c) = \omega^* - w(b) - [-w(a)]^+ + [w(a)]^+ + \alpha = \omega^* - w(b) + w(a) + \alpha = \omega^*$, as desired.

Since ω^* is the optimal value of $\mathbb{D}(G, \mathbf{w})$, (i) and (ii) follow instantly from (4)-(6). \blacksquare

6 Integralities

Given the structural description and reduction operations presented in the previous sections, we are ready to establish the main result now.

Proof of Theorem 1.5. Implication (iii) \Rightarrow (ii) follows directly from the Edmonds-Giles theorem [5] stated in Section 1. Implication (ii) \Rightarrow (i) is given by Lemma 3.1 and Theorem 3.2. It remains to show implication (i) \Rightarrow (iii).

Let $G = (V, A)$ be a permissible digraph. To prove that G is kernel Mengerian; that is, $\mathbb{D}(G, \mathbf{w})$ has an integral optimal solution for any $\mathbf{w} \in \mathbb{Z}^V$, we apply induction on $|V| + |A|$. The statement holds trivially when $|V| + |A| \leq 4$. So we proceed to the induction step, and assume that every permissible digraph $G' = (V', A')$, with $|V'| + |A'| < |V| + |A|$, is kernel Mengerian. By induction hypothesis, we may further assume that

- the underlying graph of G is connected; and
- none of Reductions 1-6 in Section 5 is applicable to G (see Lemmas 5.1, 5.2, 5.4, and 5.5).

Throughout this section we reserve symbol H for a strongly connected component of G with no outgoing arcs (that is, G contains no arc from $V(H)$ to $V \setminus V(H)$).

Claim 6.1. *The following statements hold for H :*

- (i) H contains at least two vertices;
- (ii) H is bipartite; and
- (iii) no two vertices in H have a common in-neighbor outside H .

Since Reduction 1 does not apply to G , we have (i). Statement (ii) follows directly from (1.1). To prove statement (iii), suppose on the contrary that some vertices a and b in H have a common in-neighbor c outside H . Let Q_a (resp. Q_b) be a directed path from a to b (resp. from b to a) in H . Then $Q_a \cup Q_b \cup \{(c, a), (c, b)\}$ would contain a gear, a contradiction. Thus Claim 6.1 is established.

Claim 6.2. *H is obtained from an induced even cycle $C = v_1 v_2 \dots v_{2t} v_1$, with $t \geq 3$, by replacing each even-indexed vertex v_{2i} , for $1 \leq i \leq t$, with a stable set S_{2i} of size at least two, with $v_{2i} \in S_{2i}$, such that*

- there is an arc from v_{2i-1} to each vertex in S_{2i} ;
- there is an arc from each vertex in S_{2i} to v_{2i+1} ; and

• $\{v_1, v_3, \dots, v_{2t-1}\}, S_2, S_4, \dots, S_{2t}$ are pairwise disjoint,
 where $v_{2t+1} = v_1$. (So H has $2 \sum_{i=1}^t |S_{2i}|$ arcs in total.)

To justify this, recall that H has a tree-like structure defined by a tree T rooted at a vertex r ; see Theorem 4.1 for its structural description and undefined notations involved in our proof. Observe that

(1) T has at least two vertices.

Otherwise, H is an induced cycle by Theorem 4.1. Since H has no outgoing arcs, it cannot contain four or more vertices, for otherwise Reduction 2 applies to three consecutive vertices on H (in view of Claim 6.1(iii)). Since G contains no odd cycle, H has only two vertices a and b , where b has a neighbor c outside H . Now $N_G^+(a) = \{b\}$, $N_G^+(b) = \{a\}$, and $(c, b) \in A$ is a cut arc of G by Theorem 4.4. So either Reduction 5 or Reduction 6 applies to $\{a, b, c\}$; this contradiction proves (1).

For each vertex v in T , let R_v be the path from r to v ; we call $|R_v|$ the *level* of v in T . Let us consider an arbitrary vertex pair (v, x) in T , such that

- (2) $x (\neq r)$ is a leaf of T and v is its parent;
- (3) x has the highest level among all leaves of T ; and
- (4) each internal vertex (if any) of $P_{v,x}$, the subpath of P_v between s_x and t_x , has degree two in H .

The existence of such a pair is guaranteed by (1) and Theorem 4.1(vi).

(5) Each internal vertex of P_x (if any) has degree two in H . Moreover, $|P_x| \leq 3$, with equality only when $(s_x, t_x) \in A$ and $|N_H^+(t_x)| \geq 2$.

The first half of this statement follows instantly from Theorem 4.1(ii) and (iii). Let abt_x be a subpath of P_x if $|P_x| \geq 3$, and let c be an out-neighbor of t_x . Note that $s_x \neq t_x$ if $|P_x| = 3$ because H contains no odd cycle. Since Reduction 2 does not apply to $\{a, b, t_x\}$ or to $\{b, t_x, c\}$ (see Claim 6.1(iii)), we get the second half of (5).

(6) If $|P_x| = 3$, then $v \neq r$.

Assume on the contrary that $v = r$. By (5), we have $(s_x, t_x) \in A$ and $|N_H^+(t_x)| \geq 2$. As P_r is an induced cycle, it contains (s_x, t_x) . Let u be an out-neighbor of t_x outside $P_r \cup P_x$. From Theorem 4.1(ii) and (iii), we see that (t_x, u) is contained in P_y for some child y other than x of r in T . Since $s_y = t_x$ and $|P_r[s_x, s_y]|$ is odd, $P_r \cup P_x \cup P_y$ is a ring; this contradiction justifies (6).

(7) If $P_{v,x} = P_v[t_x, s_x]$, then $|P_{v,x}| = 0$ or $v = r$ or $|N_H^+(t_x)| = 1$.

Assume the contrary: $|P_{v,x}| \geq 1$, $v \neq r$, and t_x has an out-neighbor $u \notin P_{v,x}$. By Theorem 4.1(v), we have $t_x \neq s_v$. By Theorem 4.1(ii) and (iii), arc (t_x, u) is contained in P_y for some child y of v other than x in T . Since $s_y = t_x$, by Theorem 4.1(viii), we obtain $t_y \in P_v(s_y, t_v]$, which together with Theorem 4.1(iv) implies $s_y = s_v$, a contradiction. So (7) holds.

(8) $s_y \neq t_y$ for every child y of v in T .

Suppose on the contrary that P_y is a cycle. Since y is a leaf of T with the highest level, we have $|P_y| = 2$ by (5) (with (v, y) in place of (v, x)). Let a and b be the two vertices in P_y , where $b = s_y$, and let c be an out-neighbor of b on P_v . Then Reduction 4 applies to $\{a, b, c\}$; this contradiction establishes (8).

(9) We may assume that $|P_x| \geq 2$.

By (8), we have $|P_{v,x}| \geq 1$. Let us first consider the case when $P_{v,x} = P_v[s_x, t_x]$. Since G contains no parallel arcs, $|P_x| + |P_{v,x}| \geq 3$. By (4) and (5), each internal vertex of $P_{v,x}$ and that of P_x has degree two in H . Swapping $P_{v,x}$ and P_x if necessary, we may assume that $|P_x| \geq 2$.

It remains to consider the case when $P_{v,x} = P_v[t_x, s_x]$. Assume the contrary: $|P_x| = 1$. Thus $(s_x, t_x) \in A$. Since H contains no odd cycle, $|P_{v,x}|$ is odd. As Reduction 2 does not apply to any three consecutive vertices on $P_v(t_x, s_x]$ (see (4)), we have $|P_{v,x}| \leq 3$, with equality only when $(t_x, s_x) \in A$. Thus H contains both (s_x, t_x) and (t_x, s_x) no matter whether $|P_{v,x}| = 3$ or 1.

By Theorem 4.1(v), we have $t_x \neq s_v$. Let D be a cycle containing P_v in $\cup_{z \in V(R_v)} P_z$ (see (1) in the proof of Theorem 4.1), and let c be the out-neighbor of s_x on D . Clearly, $c \neq t_x$. If $|P_{v,x}| = 3$, then $D \cup \{(s_x, t_x), (t_x, s_x)\}$ would be a ring. So we assume $|P_{v,x}| = 1$. Since Reduction 4 does not apply to $\{t_x, s_x, c\}$, we have $|N_H^+(t_x)| \geq 2$ and hence $v = r$ by (7). Thus P_r is not an induced cycle as one of (s_x, t_x) and (t_x, s_x) is outside P_r , contradicting Theorem 4.1(i). So (9) holds.

(10) $|P_x| = 2$.

Otherwise, from (5), (6), and (9) we see that $|P_x| = 3$, $v \neq r$, $(s_x, t_x) \in A$, and $|N_H^+(t_x)| \geq 2$. By (7), we have $t_x \in P_v(s_x, t_v]$. Thus Theorem 4.1(iv) enforces $s_x = s_v$. Let u be an out-neighbor of t_x such that arc (t_x, u) is outside P_v . In view of Theorem 4.1(ii) and (iii), t_x is an internal vertex of P_v and (t_x, u) belongs to P_y for some child y of v in T . Thus the existence of both P_x and P_y contradicts Theorem 4.1(vii), and hence (10) is justified.

(11) $|P_{v,x}| = 2$ and $P_{v,x} = P_v[s_x, t_x]$. Moreover, if $v \neq r$, then $s_x = s_v$ and t_x is an internal vertex of P_v .

By (8), we have $|P_{v,x}| \geq 1$. By Claim 6.1(ii), $|P_{v,x}|$ has the same parity as $|P_x|$ and hence is even by (10). Since Reduction 2 does not apply to any three consecutive vertices on $P_{v,x}$ (see Claim 6.1(iii) and (4)), we have $|P_{v,x}| = 2$. As Reduction 3 does not apply to the four vertices in $P_x \cup P_{v,x}$, we further obtain $P_{v,x} = P_v[s_x, t_x]$. If $v \neq r$, then $s_x = s_v$ by Theorem 4.1(iv) and hence t_x is an internal vertex of P_v by Theorem 4.1(ii). Thus (11) follows.

(12) For each in-neighbor u of t_x outside P_v , we have $N_H^-(u) = \{s_x\}$ and $N_H^+(u) = \{t_x\}$.

By (11) and Theorem 4.1(ii), arc (u, t_x) is contained in P_y for some child y of v in T . If $v \neq r$, then $s_y = s_x = s_v$ by (11) and Theorem 4.1(iv) and (vii). If $v = r$, we also have $s_y = s_x$ by Theorem 4.1(ix). Since y has the same level as x in T , by (5), (10), and (11) (with (v, y) in place of (v, x)), we have $N_H^-(u) = \{s_x\}$ and $N_H^+(u) = \{t_x\}$. This proves (12).

(13) $|N_H^+(t_x)| \geq 2$.

Suppose on the contrary that t_x has only one out-neighbor, say a . Then $a \in P_v$ by (11). If $N_H^+(a) = \{b\}$ and $b \neq s_x$, then Reduction 2 applies to $\{t_x, a, b\}$ by (12), a contradiction. If $N_H^+(a) = \{s_x\}$, then Reduction 3 applies to the four vertices on $P_v[s_x, a]$, again a contradiction. So $|N_H^+(a)| \geq 2$. Let R_v be the path from r to v in T and let D be a shortest cycle in $\cup_{q \in V(R_v)} P_q$ containing P_v (see (1) in the proof of Theorem 4.1). Then D contains precisely one out-neighbor of a , say b . Let c be an out-neighbor of a outside D . Since H is strongly connected, it contains a path Q from a to a vertex on D , passing through (a, c) , such that all internal vertices of Q are outside D . Thus $P_v[s_x, a] \cup P_x \cup Q$ would be a cycle if Q is a cycle and $D \cup P_x \cup Q$ would be a ring otherwise; this contradiction establishes (13).

(14) $v = r$ and hence T is a star centered at v .

Suppose on the contrary that $v \neq r$. By (11), $s_x = s_v$ and t_x is an internal vertex of P_v . By (13), t_x has an out-neighbor u such that $\text{arc}(t_x, u)$ is outside P_v . By Theorem 4.1(ii), $\text{arc}(t_x, u)$ is contained in P_y for some child y of v in T , which contradicts Theorem 4.1(vii). So (14) is true.

(15) Let y be a child of r in T such that $s_y = t_x$ and, subject to this, $|P_r[s_y, t_y]|$ is as small as possible. Then each internal vertex (if any) of $P_r[s_y, t_y]$ has degree two in H .

Suppose on the contrary that some internal vertex of $P_r[s_y, t_y]$ has degree at least three in H . By (14), r has a child z in T such that at least one end of P_z is on $P_r[s_y, t_y]$. By Theorem 4.1(vi), both s_z and t_z are on $P_r[s_y, t_y]$. By (8), Theorem 4.1(ix), and (11) (with y in place of x), we obtain $t_z \in P_r(s_z, t_y]$. By Theorem 4.1(ix), we further have $s_z = s_y$. Thus $|P_r[s_z, t_z]| < |P_r[s_y, t_y]|$, contradicting the hypothesis on y . Therefore (15) holds.

By (13), the vertex y specified in (15) is available. From (5), (10), and (11) (with y in place of x), we see that $|P_y| = |P_r[s_y, t_y]| = 2$ and that both the internal vertex of P_y and that of $P_r[s_y, t_y]$ have degree two. The process can be repeated by replacing x with y . Let X be the set of all children x of r in T , such that each internal vertex of $P_r[s_x, t_x]$ has degree two in H (see (4) and (11)). From the above observation, we deduce that $P_r \cup (\cup_{x \in X} P_x)$ is obtained from $P_r = v_1 v_2 \dots v_{2t} v_1$ by replacing each even-indexed vertex v_{2i} with a stable set S_{2i} of size at least two such that

- there is an arc from v_{2i-1} to each vertex in S_{2i} ;
- there is an arc from each vertex in S_{2i} to v_{2i+1} ; and
- $\{v_1, v_3, \dots, v_{2t-1}\}, S_2, S_4, \dots, S_{2t}$ are pairwise disjoint,

where $v_{2t+1} = v_1$. For convenience, we view v_{2i} as a vertex in S_{2i} for $1 \leq i \leq t$. Observe that $t \geq 3$, for otherwise Reduction 3 would apply to the four vertices on P_r .

Let y be an arbitrary child of r in T . Since the terminus t_y of P_y is some v_{2i-1} , by (12) we have $y \in X$, and hence $H = P_r \cup (\cup_{x \in X} P_x)$. Therefore Claim 6.2 is established.

We shall rely heavily on the structural description of H given in Claim 6.2. Let $B = \{(v_1, u) : u \in S_2 \setminus v_2\}$, let $G' = (V, A')$ be the digraph $G \setminus B$, and let $H' = H \setminus B$. Set $K_1 = \{v_1, v_3, \dots, v_{2t-1}\}$ and $K_2 = V(H) \setminus K_1$. Observe that

(16) K_1 and K_2 are the only two kernels in each of H and H' . Thus the restriction of any kernel of G to H is either K_1 or K_2 .

In the remainder of our proof, we reserve the symbol \mathbf{x}' for an integral optimal solution to $\mathbb{P}(G', \mathbf{w})$ and the pair $(\mathbf{y}', \mathbf{z}')$ for an integral optimal solution to $\mathbb{D}(G', \mathbf{w})$ such that

(17) $z'(v_1)$ is as small as possible.

It follows instantly from (16) that

(18) \mathbf{x}' is also a feasible solution to $\mathbb{P}(G, \mathbf{w})$.

Claim 6.3. $H = G$.

Assume the contrary: $H \neq G$. Let us define $\bar{\mathbf{w}} \in \mathbb{Z}^{V(H)}$ as

- $\bar{w}(v) = w(v) + \sum_{u \in N_G^-(v) \setminus V(H)} y'((u, v)) - z'(N_G^-(v) \setminus V(H))$ for each $v \in V(H)$.

Set $\bar{\mathbf{x}} = \mathbf{x}'|_{V(H)}$. Using optimality of \mathbf{x}' and $(\mathbf{y}', \mathbf{z}')$, it is easy to check that $\bar{\mathbf{x}}$ and $(\mathbf{y}'|_{A(H')}, \mathbf{z}'|_{V(H)})$ are feasible solutions to $\mathbb{P}(H', \bar{\mathbf{w}})$ and $\mathbb{D}(H', \bar{\mathbf{w}})$, respectively, and satisfy the complementary slackness condition (because H has no outgoing arcs). So

(19) $\bar{\mathbf{x}}$ is an integral optimal solution to $\mathbb{P}(H', \bar{\mathbf{w}})$ and hence also an integral optimal solution to $\mathbb{P}(H, \bar{\mathbf{w}})$ by (16). Furthermore, $\bar{\mathbf{w}}^T \bar{\mathbf{x}} = -y'(A(H')) + z'(V(H))$.

Let $(\bar{\mathbf{y}}, \bar{\mathbf{z}})$ be an integral optimal solution to $\mathbb{D}(H, \bar{\mathbf{w}})$. By (19), we have

$$(20) \quad \bar{\mathbf{w}}^T \bar{\mathbf{x}} = -\bar{y}(A(H)) + \bar{z}(V(H)).$$

Now let us define $\mathbf{y}^* \in \mathbb{Z}^A$ and $\mathbf{z}^* \in \mathbb{Z}^V$ as

- $\mathbf{y}^*|_{A(H)} = \bar{\mathbf{y}}$ and $\mathbf{y}^*|_{A \setminus A(H)} = \mathbf{y}'|_{A \setminus A(H)}$;
- $\mathbf{z}^*|_{V(H)} = \bar{\mathbf{z}}$ and $\mathbf{z}^*|_{V \setminus V(H)} = \mathbf{z}'|_{V \setminus V(H)}$.

Since H has no outgoing arcs, the definition of $\bar{\mathbf{w}}$ guarantees that $(\mathbf{y}^*, \mathbf{z}^*)$ is a feasible solution to $\mathbb{D}(G, \mathbf{w})$. Moreover,

$$\begin{aligned} -y^*(A) + z^*(V) &= -\bar{y}(A(H)) + \bar{z}(V(H)) - y'(A \setminus A(H)) + z'(V \setminus V(H)) \\ &= \bar{\mathbf{w}}^T \bar{\mathbf{x}} - y'(A \setminus A(H)) + z'(V \setminus V(H)) \\ &= \bar{\mathbf{w}}^T \bar{\mathbf{x}} + y'(A(H')) - z'(V(H)) - y'(A') + z'(V) \\ &= -y'(A') + z'(V) \\ &= \mathbf{w}^T \mathbf{x}', \end{aligned}$$

where the second equality follows from (20) and the fourth one from (19). Thus, by (18) and the LP-duality theorem, \mathbf{x}' and $(\mathbf{y}^*, \mathbf{z}^*)$ are integral optimal solution to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively. Therefore we may assume that $H = G$, otherwise the desired statement of Theorem 1.5 has been established.

Claim 6.4. $z'(v_1) \geq 1$.

Otherwise, $z'(v_1) = 0$. Let us define $\mathbf{y}^* \in \mathbb{Z}^A$ by

- $\mathbf{y}^*|_{A'} = \mathbf{y}'|_{A'}$ and $y^*((v_1, u)) = 0$ for each $(v_1, u) \in B$.

Then $(\mathbf{y}^*, \mathbf{z}')$ is a feasible solution to $\mathbb{D}(G, \mathbf{w})$ with value $\mathbf{w}^T \mathbf{x}'$. Thus, by (18) and the LP-duality theorem, \mathbf{x}' and $(\mathbf{y}^*, \mathbf{z}')$ are integral optimal solution to $\mathbb{P}(G, \mathbf{w})$ and $\mathbb{D}(G, \mathbf{w})$, respectively. So we may assume that $z'(v_1) \geq 1$.

Claim 6.5. $\phi_u(\mathbf{y}', \mathbf{z}') = w(u)$ for all $u \in K_2$.

Assume the contrary: $\phi_u(\mathbf{y}', \mathbf{z}') < w(u)$ for some $u \in K_2$, say $u \in S_{2i}$ for some i with $1 \leq i \leq t$ (recall Claim 6.2). Set $\epsilon = \min\{w(u) - \phi_u(\mathbf{y}', \mathbf{z}'), z'(v_1)\}$. From Claim 6.4, we see that ϵ is a positive integer. Let us define $\mathbf{y} \in \mathbb{Z}^{A'}$ and $\mathbf{z} \in \mathbb{Z}^V$ as

- $y(e) = y'(e) + \epsilon$ if $e = (v_{2j-1}, v_{2j})$ for $i+1 \leq j \leq t$, and $y(e) = y'(e)$ otherwise;
- $z(v_1) = z'(v_1) - \epsilon$, $z(v_{2j}) = z'(v_{2j}) + \epsilon$ for $i+1 \leq j \leq t$, $z(u) = z'(u) + \epsilon$, and $z(v) = z'(v)$ otherwise.

Using Claims 6.2 and 6.3, it is a routine matter to check that (\mathbf{y}, \mathbf{z}) is a feasible solution to $\mathbb{D}(G', \mathbf{w})$. Since $-y(A') + z(V) = -(y'(A') + (t-i)\epsilon) + z'(V) - \epsilon + (t-i+1)\epsilon = -y'(A') + z'(V)$, pair (\mathbf{y}, \mathbf{z}) is also an integral optimal solution to $\mathbb{D}(G', \mathbf{w})$. As $z(v_1) < z'(v_1)$, the existence of (\mathbf{y}, \mathbf{z}) contradicts the choice of $(\mathbf{y}', \mathbf{z}')$ (see (17)). Thus Claim 6.5 is justified.

Claim 6.6. $z'(u) = 0$ for all $u \in K_1 \setminus v_1$.

Assume the contrary: $z'(v_{2i+1}) > 0$ for some i with $1 \leq i \leq t-1$. Set $\epsilon = \min\{z'(v_{2i+1}), z'(v_1)\}$. By Claim 6.4, ϵ is a positive integer. Let us define $\mathbf{y} \in \mathbb{Z}^{A'}$ and $\mathbf{z} \in \mathbb{Z}^V$ as

- $y(e) = y'(e) + \epsilon$ if $e = (v_{2j+1}, v_{2j+2})$ for $1 \leq j \leq t-1$ and $j \neq i$, and $y(e) = y'(e)$ otherwise;
- $z(v_1) = z'(v_1) - \epsilon$, $z(v_{2i+1}) = z'(v_{2i+1}) - \epsilon$, $z(v_{2j}) = z'(v_{2j}) + \epsilon$ for $1 \leq j \leq t$, and $z(v) = z'(v)$ otherwise.

From Claims 6.2 and 6.3, it is easy to see that (\mathbf{y}, \mathbf{z}) is a feasible solution to $\mathbb{D}(G', \mathbf{w})$. Since $-y(A') + z(V) = -(y'(A') + (t-2)\epsilon) + z'(V) - 2\epsilon + t\epsilon = -y'(A') + z'(V)$, pair (\mathbf{y}, \mathbf{z}) is also an integral optimal solution to $\mathbb{D}(G', \mathbf{w})$. As $z(v_1) < z'(v_1)$, the existence of (\mathbf{y}, \mathbf{z}) contradicts the choice of $(\mathbf{y}', \mathbf{z}')$ (see (17)). This proves Claim 6.6.

Claim 6.7. *Let $\mathbf{x}^* \in \mathbb{Z}^V$ be defined by $x^*(u) = 1$ if $u \in K_2$ and 0 otherwise. Then \mathbf{x}^* is an integral optimal solution to $\mathbb{P}(G, \mathbf{w})$.*

To justify this, note first that $G = H$ by Claim 6.3. So, using (16), \mathbf{x}^* is a feasible solution to $\mathbb{P}(G, \mathbf{w})$. By Claim 6.5, we have $w(K_2) = \sum_{u \in K_2} \phi_u(\mathbf{y}', \mathbf{z}') = -y'(A') + z'(V)$, where the second equality holds because $z'(u) = 0$ for all $u \in K_1 \setminus v_1$ by Claim 6.6, and $z'(v_1)$ appears in none of $\phi_u(\mathbf{y}', \mathbf{z}')$ with $u \neq v_2$ (recall that $G' = G \setminus B$). From the LP-duality theorem, we thus conclude that \mathbf{x}^* is an integral optimal solution to $\mathbb{P}(G', \mathbf{w})$, and hence also an integral optimal solution to $\mathbb{P}(G, \mathbf{w})$ by (16).

Let $(\mathbf{y}^*, \mathbf{z}^*)$ be an optimal solution to $\mathbb{D}(G, \mathbf{w})$. Observe that

$$(21) \quad z^*(u) = 0 \text{ for all } u \in K_1.$$

Indeed, by definition, $x^*(u) = 1$ for each $u \in K_2$. So $x^*(u) + x^*(N_G^+(u)) \geq 2$ for each $u \in K_1$. Thus (21) follows instantly from the optimality established in Claim 6.7 and the complementary slackness condition.

Claim 6.8. *$\mathbb{D}(G, \mathbf{w})$ has an integral optimal solution.*

To justify this, let \mathbb{D}' denote the linear program obtained from $\mathbb{D}(G, \mathbf{w})$ by deleting the variables $z(u)$ for all $u \in K_1$. It follows from (21) that the optimal value of \mathbb{D}' is at least $y^*(A) + z^*(V)$. By Claims 6.2 and 6.3, G is bipartite. So the incidence matrix M_1 of the underlying graph of G is totally unimodular. Let M_2 be the matrix formed by the columns of M_1 corresponding to arcs leaving K_2 . Observe that for each $u \in K_2$ and the unique arc e leaving u , the variables $z(u)$ and $y(e)$ appear, with opposite coefficients, in exactly the same inequalities of \mathbb{D}' . Clearly, the constraint matrix of \mathbb{D}' is $(-M_2, M_2)$, which is totally unimodular. Hence \mathbb{D}' has an integral optimal solution $(\mathbf{y}', \mathbf{z}')$ with value $y'(A) + z'(K_2) \geq y^*(A) + z^*(V)$. Let $\mathbf{y} \in \mathbb{Z}^A$ and $\mathbf{z} \in \mathbb{Z}^V$ be defined by $\mathbf{y} = \mathbf{y}'$, $z(u) = z'(u)$ if $u \in K_2$, and $z(u) = 0$ if $u \in K_1$. Then (\mathbf{y}, \mathbf{z}) is an integral solution to $\mathbb{D}(G, \mathbf{w})$ with value $y(A) + z(V) \geq y^*(A) + z^*(V)$. Therefore it is also optimal.

This completes the proof of Claim 6.8 and hence of Theorem 1.5. ■

Acknowledgment. We are indebted to an anonymous referee for discovering an error in our original proof and a shorter proof of Claim 6.8. We greatly appreciate his/her insightful skepticism and invaluable comments and suggestions.

References

- [1] R. Aharoni and R. Holzman, Fractional kernels in digraphs, *J. Combin. Theory Ser. B* **73** (1998), 1-6.
- [2] J. Bang-Jensen and G. Gutin, *Digraphs: Theory, Algorithms and Applications*, Springer, 2001.
- [3] J.A. Bondy and U.S.R. Murty, *Graph Theory*, Springer, 2008.
- [4] E. Boros and V. Gurvich, Perfect graphs are kernel-solvable, *Discrete Math.* **159** (1996), 35-55.
- [5] J. Edmonds and R. Giles, A min-max relation for submodular functions on graphs, *Ann. Discrete Math.* **1** (1977), 185-204.
- [6] Egres Open (Categories: *Stable Matchings and Kernels* and *Integer Polyhedra*), Egerváry Research Group on Combinatorial Optimization (EGRES), Eötvös Loránd University, Hungary. (See website <http://lemon.cs.elte.hu/egres/open/>)
- [7] A. Fraenkel, Planar kernel and Grundy with $d \leq 3$, $d_{\text{out}} \leq 2$, $d_{\text{in}} \leq 2$ are NP-complete, *Discrete Appl. Math.* **3** (1981), 257-262.
- [8] J. Edmonds, S. Gaubert and V. Gurvich, Sperner oiks, *Electron. Notes Discrete Math.* **36** (2010), 1273-1280.
- [9] H. Galeana-Sánchez and V. Neumann-Lara, Extending kernel perfect digraphs to kernel perfect critical digraphs, *Discrete Math.* **94** (1991), 181-187.
- [10] M. Garey and D. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness*, Freeman and Co., New York, 1979.
- [11] T. Király and J. Pap, Total dual integrality of Rothblum's description of the stable-marriage polyhedron, *Math. Oper. Res.* **33** (2008), 283-290.
- [12] T. Király and J. Pap, A note on kernels and Sperner's lemma, *Discrete Appl. Math.* **157** (2009), 3327-3331.
- [13] R. Irving, An efficient algorithm for the stable roommates problems, *J. Algorithms* **6** (1985), 577-595.
- [14] D. Lichtenstein, Planar formulae and their uses, *SIAM J. Comput.* **11** (1982), 329-343.
- [15] M. Richardson, Solutions of irreflexive relations, *Ann. of Math. (2)* **58** (1953), 573-590.
- [16] U. Rothblum, Characterization of stable matchings as extreme points of a polytope, *Math. Programming* **54** (1992), 57-67.
- [17] A. Schrijver, *Theory of Linear and Integer Programming*, John Wiley & Sons, New York, 1986.
- [18] J. von Neumann and O. Morgenstern, *Theory of Games and Economic Behavior*, Princeton University Press, Princeton, NJ, 1944.