

## CORRESPONDENCE

(The Editors do not hold themselves responsible for opinions expressed by correspondents)

### Post-radiotherapy morphological changes of parotid gland are dose- and radiotherapy technique-dependent

The Editor,

We would like to thank the author for providing the additional information on pre- and post-therapeutic assessment of submandibular glands in CT [1]. It is of great interest to know the time dependent changes of the submandibular glands following radiochemotherapy [1]. Regarding this, we would like to provide further information from our recent studies [2, 3]. In investigating the sonographic appearances of parotid glands in two groups of patients (nasopharyngeal carcinoma (NPC) patients received intensity-modulated radiotherapy (IMRT) with parotid sparing, and NPC patients received conventional radiotherapy without parotid sparing) and healthy subjects, we found that parotid glands in patients treated with IMRT had a sonographic appearance similar to that in healthy subjects (*i.e.* usually hyperechoic, homogeneous and marginally seen intra-parotid ducts), while parotid glands in patients treated with conventional RT tended to be hypoechoic, heterogeneous and clearly seen intra-parotid ducts [2]. The result of this study shows that the sonographic appearance of the parotid gland in post-radiotherapy patients was dependent on the radiotherapy techniques; this may be related to the different radiation doses delivered to the parotid glands.

In another study, we compared the parotid gland volume of a group of NPC patients before and 2–3 years after radiotherapy using CT [3]. The results of the study showed that the parotid gland volume decreased after radiotherapy and the degree of parotid volume shrinkage was dose-dependent [3].

The assessment of salivary glands after radiotherapy is important [4]. Radiologists and sonographers should be aware of the dose- and radiotherapy technique-dependent

variations of parotid glands when examining patients with previous head and neck radiotherapy.

<sup>1</sup>M T C YING, PhD,  
<sup>1</sup>S C H CHENG, BSc,  
<sup>1</sup>V W C WU, PhD AND  
<sup>2</sup>D L W KWONG, MD

<sup>1</sup>Department of Health Technology and Informatics  
The Hong Kong Polytechnic University Hung Hom  
Kowloon  
Hong Kong SAR  
China

<sup>2</sup>Department of Clinical Oncology  
Queen Mary Hospital  
The University of Hong Kong Pokfulam  
Hong Kong SAR  
China

(Received 28 June 2011, Accepted 5 July 2011)

DOI: 10.1259/bjr/30087983

### References

1. Gossner J. Post-radiogenic density changes on CT of the salivary gland are time-dependent. *Br J Radiol* 2011;84.
2. Cheng SCH, Ying MTC, Kwong DLW, Wu VWC. Sonographic appearance of parotid glands in patients treated with intensity-modulated radiotherapy or conventional radiotherapy for nasopharyngeal carcinoma. *Ultrasound Med Biol* 2011;37:220–30.
3. Wu VWC, Ying MTC, Kwong DLW. Evaluation of radiation-induced changes to parotid glands following conventional radiotherapy in patients with nasopharyngeal carcinoma. *Br J Radiol* 2011;84:843–849.
4. Cheng SCH, Wu VWC, Kwong DLW, Ying MTC. Assessment of post-radiotherapy salivary glands. *Br J Radiol* 2011;84: 393–402.

---

Address correspondence to: Dr Michael Ying, Department of Health Technology and Informatics, The Hong Kong Polytechnic University, Yuk Choi Road, Hung Hom, Kowloon, Hong Kong SAR, China. E-mail: htmying@polyu.edu.hk