

DMID—11-369 revised

1
2
3 Changes in nasopharyngeal carriage and serotype distribution of antibiotic-
4
5 resistant *Streptococcus pneumoniae* before and after introduction of 7-valent
6
7 pneumococcal conjugate vaccine, Hong Kong
8
9

10
11
12
13 Pak-Leung Ho^{1*}, Susan S. Chiu², Maggie Y. Chan¹, Irene Ang¹, Kin-Hung Chow¹, Yu-Lung
14
15 Lau²
16
17
18
19
20

21 ¹*Department of Microbiology and* ²*Department of Paediatrics and Adolescent Medicine, The*
22
23 *University of Hong Kong, Pokfulam Road, Hong Kong Special Administrative Region,*
24
25 *CHINA*
26
27
28
29
30

31 Keywords: *Streptococcus pneumoniae*, drug resistance, serotype, prevalence
32
33
34
35
36
37

38 *corresponding author: Division of Infectious Diseases, Department of Microbiology and
39
40 Centre of Infection, The University of Hong Kong, Queen Mary hospital, Pokfulam Road,
41
42 Pokfulam, Hong Kong SAR, CHINA. Tel.: +852-2855 4897; fax: +852-2855 1241.
43
44

45 *E-mail address:* plho@hkucc.hku.hk (P. L. Ho).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

This study assessed the changes in serotype distribution and antibiotic resistance of *Streptococcus pneumoniae* isolates in children before and after introduction of the 7-valent pneumococcal conjugate vaccine (PCV7) in Hong Kong. Nasopharyngeal specimens were collected from 1978 and 2221 children (ages, 2 to 6 years) attending day care centers or kindergartens in period 1 (1999-2000) and period 2 (2009-2010), respectively. Carriage of PCV7 serotypes decreased from 12.8% to 8.6% ($P<0.01$). The relative contribution of PCV7 serotypes 14 and 18C had decreased while that for non-PCV7 serotypes 19A, 6A, 6C, 23A and 15B had increased. In period 2, PCV7 penetration rate (at least one dose) for children aged 2, 3, 4 and 5 years were 43%, 35.7%, 26.7% and 20.4%, respectively. In multivariate analysis, PCV7 use was the only independent variable associated with fewer PCV7 serotypes carriage (OR 0.5, $P=0.001$). In period 2, high rates of dual penicillin/erythromycin nonsusceptibility were found in serotypes 6B (77.3%), 14 (100%), 19F (100%), 23F (78%), 19A (75%), 6A (87.8%), 6C (59.3%) and 23A (78.9%).

169 words

1. Introduction

Streptococcus pneumoniae is a major cause of childhood infections (Ho et al., 2006b; Ho et al., 2006a; Ho et al., 2007). In western countries, the introduction of the 7-valent pneumococcal conjugate vaccine (PCV-7) has led to drastic decline in the incidence of invasive pneumococcal disease (IPD) caused by serotypes included in the vaccine (Dagan, 2009a; Pilishvili et al., 2010). The vaccine also protects against carriage of those serotypes in the vaccinated children (Hanquet et al., 2010; Huang et al., 2009). Consequently, there is reduced transmission of these serotypes by vaccinated children; resulting an extension of protection to unvaccinated population (i.e. herd immunity) (Centers for Disease Control and Prevention, 2005). However, these benefits have been partly offset by an increase in the incidence of diseases caused by serotypes not included in the vaccine; the magnitude of such replacement remained low relative to the reduction in PCV7 diseases (Pilishvili et al., 2010). In February 2010, the Food and Drug Administration in the United States approved a 13-valent pneumococcal vaccine (PCV13) which provides coverage for the seven serotypes common to PCV7 and six additional serotypes (1, 3, 5, 6A, 7F and 19A) (Centers for Disease Control and Prevention, 2010b). In the United States, PCV13 has been recommended to replace PCV7 in the childhood immunization program from February 2010 onward (Centers for Disease Control and Prevention, 2010b). The policy change is expected to further reduce IPD in the United States (Centers for Disease Control and Prevention, 2010a).

In Hong Kong, PCV7 has been available in the market since October 2005. The 10-valent pneumococcal conjugate vaccine (PCV10) and PCV13 were later marketed in August 2009 and May 2010, respectively (Ho et al., 2011). In Asia, Hong Kong is one of the first cities to introduce PCV7 into the childhood immunization program and this was implemented since September 2009 (Centre for Health Protection, 2011a). The immunization schedule consists of a standard three-dose primary series (at 2, 4 and 6 months of age) and a booster

1 dose at 12-15 months. Before its introduction into the childhood immunization program,
2 PCV7 was available as a self-financed item and its uptake in the vaccine target population
3 had been low (Ho et al., 2011). In our locality, the effect of the introduction of PCV7 to the
4 market and before its full implementation through the childhood immunization program upon
5 IPD was recently reported by our group (Ho et al., 2011). In children <5 years, our data
6 showed that the proportion of PCV7 serotypes had declined from 89.5% in 1995-2001 to
7 65.7% in 2007-2009 (Ho et al., 2011). The current study was conducted to assess the changes
8 in serotype distribution and antibiotic resistance of nasopharyngeal isolates prior to and
9 during the early introduction of PCV7 in Hong Kong.
10
11
12
13
14
15
16
17
18
19
20
21
22
23

24 **2. Materials and method**

25 *2.1 Study design and data collection*

26
27
28
29 Nasopharyngeal specimens were collected from children between 2 and 6 years of age
30 who attend day care centers (DCC) or kindergartens (KG) in Hong Kong during September
31 2009 to April 2010 (period 2). A similarly designed study was conducted during December
32 1999 and June 2000 (period 1) (Chiu et al., 2001). The same recruitment process and
33 sampling strategy was used in this and the previous study (Chiu et al., 2001). In brief, Hong
34 Kong is divided into 18 school districts, and the sample size of each district was calculated
35 according to the number of DCC and KG places in each district. Ages of children who attend
36 DCC and KG range from 2 to 6 years and 3 to 6 years, respectively. Normally, all children
37 attend 5 days per week for 7 to 9 hours a day in DCC and for 3 to 4 hours a day in a KG. Out
38 of the total of 147,516 DCC and KG places, DCC accounted for 30% while KG accounted for
39 70%. This ratio was used to calculate the number of children to be recruited from DCC and
40 KG from each district to make the targeted study population of about 2000. Parental consent
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 was obtained by the research staff. The protocol is approved by the Institutional Review
2 Board at the Hong Kong West Cluster/University of Hong Kong.
3

4
5 As in the previous study (Chiu et al., 2001), a standardized questionnaire was used to
6
7 obtain the following information from parents of participating children: household size,
8
9 overcrowding (≤ 5.5 m/person in accordance with the guideline of the Hong Kong Housing
10
11 Authority), number and age of siblings, participant demographics, medical history, presence
12
13 of symptoms of upper respiratory tract infection (URTI) symptoms at the time of sampling,
14
15 recent use of antibiotics (past 3 months), vaccination history, physician visits (past 3 months)
16
17 and prior hospitalization (past 1 year).
18
19
20
21
22
23

24 *2.2 Specimen collection and microbiological methods*

25

26
27 A calcium alginate-tipped swab on a flexible aluminum wire (TRANSWAB per nasal;
28
29 Medical Wire and Equipment Co. Ltd, Corsham, Wilts, England) was used. The specimens
30
31 were brought back to the laboratory immediately for processing. For selective isolation of *S.*
32
33 *pneumoniae*, swabs were inoculated onto 5% horse blood agar supplemented with gentamicin
34
35 (2 $\mu\text{g/ml}$) and incubated in 5% CO₂ for 16 to 24 hours. All isolates were identified by colony
36
37 morphology, Gram stain, optochin susceptibility, and bile solubility. Isolates with atypical
38
39 optocin/bile solubility test results were confirmed by a slide co-agglutination test (Phadebact
40
41 Pneumococcus Test, Remel).
42
43
44

45
46 Susceptibility of the isolates to antibiotics was determined by Etest (AB Biodisk,
47
48 Solna, Sweden) or the disc diffusion method. The D-test was used to detect inducible
49
50 resistance to clindamycin. Quality control strains were included on each day of testing. All
51
52 results were interpreted according to the Clinical Laboratory Standards Institute (Clinical and
53
54 Laboratory Standard Institute,). Pneumococcal isolates were serotyped by PCR assays and/or
55
56 the quellung reactions. The isolates were initially tested by a sequential multiplex PCR
57
58
59
60
61
62
63
64
65

1 approach. Strains that could not be serotyped by the PCR assays and those that required
2 further testing to the serotype level were then tested by the quellung method with sera of
3 various reactivities from the Statens Serum Institut (Copenhagen, Denmark) (Ho et al., 2004;
4 Pai et al., 2006). The serotypes distribution for the isolates collected in period 1 has been
5 previously reported but the method do not allow some subtypes to be distinguished (Ho et al.,
6 2004). Hence, all the isolates (n=163) that could not be identified the serotype level and those
7 that were designated an untypeable at that time were retrieved and tested again by PCR
8 methods (Pai et al., 2006). PCR assays were used to identify the newly described 6C and 6D
9 serotypes (Ho et al., 2010). All serogroup 6 isolates, including those collected in period 1
10 were tested by PCR assays that could distinguish the 4 subtypes (6A-6D) of serogroup 6
11 isolates. All isolates were tested in a uniform manner.

28 2.3 Statistical analysis

31 Data from the two similarly designed studies was pooled and compared. The sampling
32 periods, 1999-2000 (period 1) and 2009-2010 (period 2), were intended to indicate time
33 windows before and after the registration and marketing of PCV7 in Hong Kong, respectively.
34 The serotype coverage for the 7-valent, 10-valent and 13-valent PCVs for carriage was
35 calculated as the proportion of all isolates included in the vaccine formulations, without
36 taking into account the potential serogroup cross-protection. In view of the demonstrated
37 efficacy of PCV7 for serotype 6A, additional coverage due to this cross protection was also
38 calculated. The Chi square or Student's *t* test was used to compare participant demographics,
39 pneumococcal carriage rates and serotype frequencies between the two collection periods.
40 Potential risk factors for carriage of PCV7 were studied by univariate analysis. Variables that
41 were significant in the univariate analysis were further analyzed by logistic regression using
42 the forward-conditional method. The correlation between PCV7 penetration and reduction in
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 carriage of PCV7 serotypes by age groups was analyzed by the Correlate Bivariate analysis.
2 A P value of <0.05 was considered to indicate statistical significance. All statistic analysis
3 was performed by the SPSS statistic package.
4
5
6
7
8
9

10 **3. Results**

11 *3.1 Demographics*

12 There were 1978 and 2211 child participants in period 1 and period 2, respectively.
13
14 As shown in **Table 1**, there are some differences in the characteristics of the children
15 recruited in the two periods. The average age for period 2 children was significantly younger
16 than that for period 1 children. Four variables were less common among period 2 children:
17 recent antibiotic use, overcrowding, URTI at time of sampling and physician visits. However,
18 about twice as many children had history of prior hospitalization in period 2 than in period 1.
19 The two groups of children had similar household size and number of siblings.
20
21
22
23
24
25
26
27
28
29
30

31 As PCV7 has only been available in Hong Kong after October 2005, no children had
32 been vaccinated with PCV7 in period 1. Among children recruited in period 2, the PCV7
33 vaccine penetration (at least one dose) was 28.1% (622/2211). PCV7 vaccination was
34 significantly more likely in younger children than in older children. The age-stratified PCV7
35 penetration rate was 20.4% (149/731) among children aged 5 years, 26.7% (203/759) among
36 children aged 4 years, 35.7% (196/549) among children aged 3 years and 43% (74/172)
37 among children aged 2 years (P <0.001, chi square for trend). The age at first vaccination
38 could be obtained from 366 children. Among these children, 20.5% (75/366) were vaccinated
39 at age ≤1 year, 49.7% (182/366) at age 2 years and 29.8% (109/366) at age 3-5 years. At the
40 time of sampling, 80.7% (502/622) of the vaccinated children could be considered age
41 appropriately vaccinated for PCV7. In period 2, 26.1% (578/2211) children had received
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 influenza vaccination in past year. Information about influenza vaccination was not obtained
2 for period 1.
3
4
5
6

7 3.2 Serotype distribution 8

9 A total of 730 isolates were included in the study, comprising 383 isolates collected in
10 period 1 and 347 isolates in period 2. PCV7 serotypes included 443 isolates, PCV13-
11 nonPCV7 serotypes included 90 isolates and non-PCV13 serotypes included 207 isolates.
12 **Table 2** showed the serotype distribution of the isolates according to these PCV categories for
13 the two collection periods. In both collection periods, the three most prevalent serotypes were
14 6B, 19F and 23F. The prevalence of seven serotypes had significantly changed during the two
15 time periods. The two serotypes that had become less common in period 2 were 14 (8.9% vs.
16 2.6%, $P<0.01$) and 18C (2.1% vs. 0.6%, $P=0.049$). The five serotypes that had significantly
17 increased in prevalence were 19A (0% vs. 2.3%, $P<0.001$), 6A (3.4% vs. 11.8%, $P<0.001$),
18 6C (1.8% vs. 7.8%, $P<0.001$), 23A (1.6% vs. 5.5%, $P=0.003$) and 15B (0.3% vs. 2.6%,
19 $P=0.007$).
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 3.3 Changes in pneumococcal conjugate vaccine coverage over time 40

41 **Table 2** also showed that the serotype coverage of PCV7 had significantly decreased
42 from 66.1% in period 1 to 54.8% in period 2 ($P = 0.001$). The reduction in the serotype
43 coverage of PCV7 was greater among isolates obtained from younger children. The serotype
44 coverage of PCV7 in period 1/period 2 for children of different ages was as follows: 2 year
45 (94.1%/41.3%), 3 years (69.5%/56.2%), 4 years (69.1%/57.4%) and 5 years (59.8%/54%),
46 respectively. Due to the absence of serotypes 1, 5 and 7F, serotype coverage of PCV7 and
47 PCV10 was the same. The serotype coverage of PCV13 was similar in period 1 and period 2,
48 being 72.6% and 70.6% for all ages, respectively. In period 1, the difference in serotype
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 coverage of PCV7 (66.1%) and PCV13 (72.6%) approached but did not reach statistical
2 significance ($P=0.05$). In period 2, serotype coverage of PCV13 was 15.9% higher than that
3 of PCV7 ($P<0.001$, 95% confidence interval for difference, 8.7 – 22.8%). However, if
4 complete cross protection against 6A was assumed, the difference in serotype coverage of
5 PCV13 and PCV7 became insignificant (4%, 95% CI -2.9% to 10.9%).
6
7
8
9
10

11 3.4 Changes in pneumococcal carriage over time 12 13 14 15

16 Overall, the nasopharyngeal carriage rate in period 2 was 15.7% (347/2221) which
17 was lower than that in period 1 (19.4%, 383/1978, $P < 0.01$). However, the carriage rate of
18 penicillin-nonsusceptible (Pen-NS) *S. pneumoniae* in the two periods remained unchanged
19 (11.3% vs. 11.1%, $P = 0.8$). Table 3 showed that the carriage rate of PCV7 isolates had
20 significantly decreased from 12.8% (253/1978) in period 1 to 8.6% (190/2211) in period 2 (P
21 <0.001). The decline in PCV7 carriage was most remarkable among children aged 2 years
22 (from 27.1% to 7.0%) and 3 years (from 22.7% to 9.1%). The changes among children aged 4
23 years and 5 years were only modest. After stratification by age, the magnitude of reduction in
24 PCV7 carriage was found to correlate positively with the PCV7 penetration rate (correlate
25 bivariate analysis, Pearson correlation = 0.99, $P = 0.01$). The carriage rate for PCV13-
26 nonPCV10 types showed a nearly 100% increase from 1.3% (25/1978) to 2.5% (55/2211) (P
27 < 0.001). Increases in PCV13-nonPCV10 serotypes were greatest in children aged 2 years but
28 the difference was not statistically significant. The prevalence of non-PCV13 carriage remain
29 unchanged over the two time periods (5.3% [105/1978] vs. 4.6% [102/2211], $P = 0.5$).
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 To further understand how availability of PCV7 might have affected pneumococcal
52 carriage, the data for children enrolled in period 2 was subjected to univariate and
53 multivariate analyses. Univariate analysis showed that overcrowding had positive correlation
54 with PCV7 carriage (odds ratio [OR] 1.8, 95% CI 1.03-3.2; $P=0.04$) while a history of PCV7
55
56
57
58
59
60
61
62
63
64
65

1 vaccination had negative correlation (OR 0.5, 95% CI 0.3-0.8; $P=0.001$). The PCV7 carriage
2 rate in children who were age appropriately vaccinated for PCV7 was significantly lower
3 than those who were not (4.8% vs. 9.7%, respectively, $P < 0.001$). No statistically significant
4 association between PCV7 carriage and the other child characteristics (recent antibiotic use,
5 prior hospitalization, respiratory symptoms at time of sampling, having young siblings aged
6 ≤ 5 years and recent physician visit) were found. In the logistic regression, a history of PCV7
7 vaccination was the only variable significantly associated with a lower chance of PCV7
8 carriage (OR 0.5, 95% CI 0.3-0.7; $P=0.001$). On the other hand, the carriage rates for PCV13-
9 nonPCV10 (2.2% vs. 2.6%, $P = 0.6$, respectively) and non-PCV13 (5.2% vs. 4.4%, $P = 0.5$,
10 respectively) isolates were similar among children who were and were not age appropriately
11 vaccinated for PCV7.
12
13
14
15
16
17
18
19
20
21
22
23
24
25

26 In period 1, a history of recent use of antibiotics was significantly associated with
27 carriage of penicillin-nonsusceptible *S. pneumoniae* (Odds ratio [OR] 1.9, 95% confidence
28 interval [CI], 1.2-2.9, $P=0.003$) but not with erythromycin-nonsusceptible *S. pneumoniae*
29 ($P=0.2$). In period 2, on the contrary, recent use of antimicrobial agents was significantly
30 associated erythromycin-nonsusceptible *S. pneumoniae* (OR 4.7, 95% CI, 1.7-14.0) but not
31 with penicillin-nonsusceptible *S. pneumoniae* ($P=0.2$).
32
33
34
35
36
37
38
39
40
41
42
43

44 3.5 Changes in antimicrobial susceptibility over time

45 Table 4 summarized the serotype-specific antimicrobial susceptibility rates of the
46 isolates for the two collection periods. Overall, it showed that the rates of penicillin
47 nonsusceptibility (meningitis breakpoint, 58.2% vs. 70.6%, for period 1 vs. period 2,
48 respectively, $P<0.001$), erythromycin nonsusceptibility (77.0% vs. 86.7%, for period 1 vs.
49 period 2, respectively, $P<0.001$), and dual penicillin/erythromycin nonsusceptibility (52.5%
50 to 70.3%, for period 1 vs. period 2, respectively, $P<0.001$) had significantly increased over
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 time. There was no significant change in nonsusceptibility rate for levofloxacin (0% vs. 0.3%)
2 over time. The capsular serotype of the only isolate with resistance to levofloxacin (MIC 16
3 $\mu\text{g/ml}$) was 19F.
4
5

6
7 Stratification by serotype showed that the penicillin and dual penicillin/erythromycin
8 nonsusceptibility rates for serotypes 19F, 14, 6A and 23A had significantly increased over the
9 two time periods. The changes for serotype 6A were most dramatic. The erythromycin
10 nonsusceptibility rates for these serotypes were similar in the two time periods. In period 1,
11 five serotypes (6B, 19F, 23F, 14 and 6C) were often nonsusceptible to both penicillin and
12 erythromycin with rates of 42.9% to 86.5%. In period 2, these serotypes continued to exhibit
13 high rates of dual penicillin/erythromycin nonsusceptibility (59.3% to 100%). Besides these,
14 an additional three serotypes (19A, 6A and 23A) were found to exhibit high rates of dual
15 penicillin/erythromycin nonsusceptibility (75% to 87.8%). No serotype 19A was found in
16 period 1. Six (75%) of the eight serotype 19A isolates found in period 2 were nonsusceptible
17 to both penicillin and erythromycin.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 PCV7 covered significantly fewer dual penicillin/erythromycin-nonsusceptible
35 isolates collected in period 2 than in period 1 (66% vs. 93.5%, respectively, $P<0.001$).
36 Although PCV13 covered more isolates, the serotype coverage for dual
37 penicillin/erythromycin-nonsusceptible isolates in period 2 was still lower than that for period
38 1 (83.1% vs. 94.0%, respectively, $P<0.001$).
39
40
41
42
43
44
45

46 Figure 1 showed a scatter plot of the serotype-specific prevalence with the penicillin-
47 nonsusceptibility rates. Curve fitting and linear regression revealed that the two variables had
48 significant associations for both time periods. The correlation coefficient was higher in period
49 1 ($R\text{-square} = 0.74$, $P<0.001$) than in period 2 ($R\text{-square} = 0.43$, $P<0.001$). Weaker but still
50 statistically significant correlations were obtained for erythromycin nonsusceptibility. The
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 finding showed that antimicrobial resistance could be an important determinant of the
2 serotype prevalence.
3
4
5
6

7 **4. Discussion**

8
9 This study assessed the changes in incidence of pneumococcal carriage, antimicrobial
10 resistance rates and serotype distribution of nasopharyngeal isolates prior to and during early
11 introduction of PCV7 in Hong Kong. The carriage of PCV7 serotypes was found to have
12 declined significantly, especially among younger children in whom the vaccine penetration is
13 higher. This was partially off-set by a slight increase in the non-PCV7 serotypes. The net
14 effect was that children were less likely to carry pneumococci in the latter period. This
15 occurred at a time when the vaccine uptake was still low (20.4% to 43%) among children
16 aged 2 to 5 years. Consequently, fewer isolates recovered in period 2 were included in the
17 PCV7 formulation than in period 1. These observations were in agreement with experiences
18 in places where PCV7 was made readily available earlier (Dagan, 2009b; Huang et al., 2009;
19 Sa-Leao et al., 2009). However, not all the seven PCV7 serotypes had decreased **and overall**
20 **carriage of types included in PCV7 was still high.** The types that had declined significantly
21 were serotype 14 and 18C. This may be explained by variations in the vaccine efficacy for
22 the different serotypes (Whitney et al., 2006). In the Kaiser PCV7 efficacy study, the
23 geometric mean concentration of pneumococcal antibodies was found to be highest for
24 serotype 14, **followed** by serotype 18C and 6B (Black et al., 2000). The antibody level
25 concentrations for the remaining serotypes were similar (Black et al., 2000). Since the
26 vaccine uptake was low, it might take longer for the other serotypes to decrease. While
27 conjugate vaccines confer protection against carriage of types included in the vaccines and
28 occasionally other types in the same serogroups, they provide no protection against
29 serogroups that are not included in the vaccine. Therefore, conjugate vaccine use might be
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 expected to contribute to the observed increase in non-PCV7 types (19A, 6A, 6C, 23A and
2 15B). Nonetheless, we were unable to demonstrate any association between PCV7 use and
3 carriage of these serotypes.
4
5

6
7 In our locality, similar temporal changes have also been demonstrated among isolates
8 causing invasive disease in children (Ho et al., 2011). Among children aged <5 years, the
9 frequencies of serotype 14 and 18C had decreased from 36% to 15.7% and from 4.7% to 0%,
10 respectively, before (1995-2001) and shortly after (2005-2009) introduction of PCV7 to the
11 Hong Kong market (Ho et al., 2011). Over the same time periods, invasive disease caused by
12 serotype 19A had increased from 0% to 12.9% and serotype 6A increased from 2.3% to 7.1%
13 (Ho et al., 2011). Despite marked increases in prevalence of serotypes 6C, 15B and 23A
14 among isolates causing nasopharyngeal colonization, such serotypes were rarely found
15 among our invasive isolates. Although almost all children become colonized with
16 pneumococci during the first few years of life, only a small fraction of the colonization
17 proceed to invasive disease. In pneumococci, the capsular polysaccharide is assumed to be
18 the most important factor that determine the capacity of specific strains to cause invasive
19 diseases (Melin et al., 2010). According to one report, the invasive capacity of different
20 pneumococcal serotypes was found to vary by as many as 50 folds (highest for 3, 7F, 18C,
21 19A, 22F and 33F; while 6C, 15B and 23A were among the serotypes with the lowest
22 invasive capacity) (Yildirim et al., 2010). This could explain why increasing carriage of
23 serotypes 6C, 15B and 23A was not associated with corresponding changes in invasive
24 disease.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50
51 Besides pneumococcal conjugate vaccine use, excessive antibiotic use is another well-
52 established factor exerting pressure on pneumococcal ecology; with incorrect usage leading
53 to an increase in multidrug-resistant strains population. In Hong Kong, antimicrobials (such
54 as oral amoxicillin-clavulanate, cephalosporins and macrolides) are frequently prescribed to
55
56
57
58
59
60
61
62
63
64
65

1 children with febrile illnesses (Lam et al., 2009). Although the frequencies of recent
2 antibiotic uses in period 2 were slightly lower than those in period 1 (Chiu et al., 2001), the
3
4 30.6% among participating children and 27% among their household members still
5
6 constituted substantial consumption. Such high antibiotic uses partly reflect widespread
7
8 misconceptions about antibiotics among the laypersons in our community (Centre for Health
9
10 Protection, 2011b). In a territory-wide telephone survey conducted in November 2010, 66.9%
11
12 of 1569 respondents believed that antibiotics are effective in treating viral infections (Centre
13
14 for Health Protection, 2011b). Thus, the observed association between penicillin
15
16 nonsusceptibility and serotype prevalence is not surprising. Heavy antibiotic use could exert
17
18 pressure through selection of resistance in previously minor serotypes and subsequent clonal
19
20 expansion (Huang et al., 2009). Alternatively, pre-existing resistant clones could undergo
21
22 capsular switching and disseminate as non-vaccine serotypes (Ansaldi et al., 2011; Moore et
23
24 al., 2008). Our finding that a history of recent antibiotic use was associated with different
25
26 types of antibiotic-resistant *S. pneumoniae* in the two time periods corroborated previous
27
28 studies that the relationship between exposure to antibiotics and the spread of resistant
29
30 pneumococci is complex (Huang et al., 2009). Since details of the recent antibiotic use were
31
32 not collected, we were unable to conduct analysis according to antibiotic class, duration of
33
34 use and dosage. Unless antibiotic usage are curtailed, the overall antimicrobial resistance
35
36 among pneumococci would remain high (Dagan, 2009b; Sa-Leao et al., 2009).

37
38
39
40
41
42
43
44
45
46 The observed increases in prevalence of non-PCV7 serotypes such as 19A, 6C, 23A
47
48 and 15B were broadly in line with secular trends in the developed countries following the
49
50 introduction of PCV7 (Hanage et al., 2005; Huang et al., 2009). In Asian countries,
51
52 emergence of the non-PCV7 serotype 19A has been associated with expansion of multidrug-
53
54 resistant ST320 isolates and could be unrelated to PCV7 use (Choi et al., 2008; Ho et al.,
55
56 2011; Shin et al., 2011). Nonetheless, marked variations in the prevalence of serotype 19A
57
58
59
60
61
62
63
64
65

1 were observed. According to a multi-national surveillance (Shin et al., 2011), the prevalence
2 of serotype 19A among clinical isolates in 2008-2009 was high in India (13.0%), Japan
3 (11.1%), South Korea (9.9%), Malaysia (9.1%), and Saudi Arabia (9.0%), but low in Hong
4 Kong (3.1%), Taiwan (4.8%), Thailand (5.2%), the Philippines (1.7%), and Vietnam (0.4%).
5
6

7
8
9 It is recognized that antibody induced by the 6B component in PCV7 may provide cross
10 protection against the vaccine-related serotype 6A (Whitney et al., 2006). However, we found
11 that the prevalence of serotype 6A among nasopharyngeal isolates has increased and were
12 associated with emergence of dual penicillin/erythromycin nonsusceptibility in this serotype.
13
14 We have excluded potential confounding with other highly similarly subtypes by using
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1 PCV13 could provide substantial benefits over PCV7 (Ho et al., 2011). Hence, Hong Kong
2 has announced to switch to PCV13 in the childhood immunization program in the last quarter
3
4 of 2011. Such change in vaccine policy is expected to impact upon the future evolution of
5
6 pneumococcal serotypes in this locality. Future studies should monitor how the emerging
7
8 serotypes will evolve and attention should be paid to non-PCV13 serotypes with higher
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

This study is supported by research grants from the Research Fund for the Control of Infectious Diseases (RFCID) of the Health and Food Bureau of the Hong Kong SAR Government. Authors had no conflict of interest to declare.

References

1. Ansaldo F, Canepa P, de Florentiis D, Bandettini R, Durando P, Icardi G (2011) Increasing incidence of *Streptococcus pneumoniae* serotype 19A and emergence of two vaccine escape recombinant ST695 strains in Liguria, Italy, 7 years after implementation of the 7-valent conjugated vaccine. *Clin Vaccine Immunol* 18: 343-345.
2. Black S, Shinefield H, Fireman B, Lewis E, Ray P, Hansen JR, Elvin L, Ensor KM, Hackell J, Siber G, Malinoski F, Madore D, Chang I, Kohberger R, Watson W, Austrian R, Edwards K (2000) Efficacy, safety and immunogenicity of heptavalent pneumococcal conjugate vaccine in children. Northern California Kaiser Permanente Vaccine Study Center Group. *Pediatr Infect Dis J* 19: 187-195.
3. Centers for Disease Control and Prevention (2005) Direct and indirect effects of routine vaccination of children with 7-valent pneumococcal conjugate vaccine on incidence of invasive pneumococcal disease--United States, 1998-2003. *MMWR Morb Mortal Wkly Rep* 54: 893-897.
4. Centers for Disease Control and Prevention (2010a) Invasive pneumococcal disease in young children before licensure of 13-valent pneumococcal conjugate vaccine - United States, 2007. *MMWR Morb Mortal Wkly Rep* 59: 253-257.
5. Centers for Disease Control and Prevention (2010b) Licensure of a 13-valent pneumococcal conjugate vaccine (PCV13) and recommendations for use among children - Advisory Committee on Immunization Practices (ACIP), 2010. *MMWR Morb Mortal Wkly Rep* 59: 258-261.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
6. Centre for Health Protection HK (2011a) Inclusion of Pneumococcal Vaccine in Childhood Immunisation Programme . Last accessed 25 June 2011.
7. Centre for Health Protection HK (2011b) Safe use of antibiotics save lives. Last accessed 25 June 2011.
8. Chiu SS, Ho PL, Chow FK, Yuen KY, Lau YL (2001) Nasopharyngeal carriage of antimicrobial-resistant *Streptococcus pneumoniae* among young children attending kindergartens and day care centers in Hong Kong. *Antimicrob Agents Chemother* 45: 2765-2770.
9. Choi EH, Kim SH, Eun BW, Kim SJ, Kim NH, Lee J, Lee HJ (2008) *Streptococcus pneumoniae* serotype 19A in children, South Korea. *Emerg Infect Dis* 14: 275-281.
10. Clinical and Laboratory Standard Institute. Performance standards for antimicrobial susceptibility testing: Twentieth informational supplement M100-S20. Wayne, Pa, National Committee for Clinical Laboratory Standards, 2010.

Ref Type: Generic

11. Dagan R (2009a) New insights on pneumococcal disease: what we have learned over the past decade. *Vaccine* 27 Suppl 3: C3-C5.
12. Dagan R (2009b) Serotype replacement in perspective. *Vaccine* 27 Suppl 3: C22-C24.
13. Hanage WP, Kaijalainen TH, Syrjanen RK, Auranen K, Leinonen M, Makela PH, Spratt BG (2005) Invasiveness of serotypes and clones of *Streptococcus pneumoniae* among children in Finland. *Infect Immun* 73: 431-435.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
14. Hanquet G, Kissling E, Fenoll A, George R, Lepoutre A, Lernout T, Tarrago D, Varon E, Verhaegen J (2010) Pneumococcal serotypes in children in 4 European countries. *Emerg Infect Dis* 16: 1428-1439.
15. Ho PL, Ang I, Chow KH, Lai EL, Chiu SS (2010) The prevalence and characteristics of *Streptococcus pneumoniae* isolates expressing serotypes 6C and 6D in Hong Kong prior to the introduction of the 7-valent pneumococcal conjugate vaccine. *Diagn Microbiol Infect Dis* 68: 439-444.
16. Ho PL, Chiu SS, Ang I, Lau YL (2011) Serotypes and antimicrobial susceptibilities of invasive *Streptococcus pneumoniae* before and after introduction of 7-valent pneumococcal conjugate vaccine, Hong Kong, 1995-2009. *Vaccine* 29: 3270-3275.
17. Ho PL, Chiu SS, Cheung CH, Lee R, Tsai TF, Lau YL (2006a) Invasive pneumococcal disease burden in Hong Kong children. *Pediatr Infect Dis J* 25: 454-455.
18. Ho PL, Chiu SS, Chow FK, Mak GC, Lau YL (2007) Pediatric hospitalization for pneumococcal diseases preventable by 7-valent pneumococcal conjugate vaccine in Hong Kong. *Vaccine* 25: 6837-6841.
19. Ho PL, Lam KF, Chow FK, Lau YL, Wong SS, Cheng SL, Chiu SS (2004) Serotype distribution and antimicrobial resistance patterns of nasopharyngeal and invasive *Streptococcus pneumoniae* isolates in Hong Kong children. *Vaccine* 22: 3334-3339.
20. Ho PL, Que TL, Ng TK, Chiu SS, Yung RW, Tsang KW (2006b) Clinical outcomes of bacteremic pneumococcal infections in an area with high resistance. *Eur J Clin Microbiol Infect Dis* 25: 323-327.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
21. Huang SS, Hinrichsen VL, Stevenson AE, Rifas-Shiman SL, Kleinman K, Pelton SI, Lipsitch M, Hanage WP, Lee GM, Finkelstein JA (2009) Continued impact of pneumococcal conjugate vaccine on carriage in young children. *Pediatrics* 124: e1-11.
 22. Lam TP, Ho PL, Lam KF, Choi K, Yung R (2009) Use of antibiotics by primary care doctors in Hong Kong. *Asia Pac Fam Med* 8: 5.
 23. Melin M, Trzcinski K, Meri S, Kayhty H, Vakevainen M (2010) The capsular serotype of *Streptococcus pneumoniae* is more important than the genetic background for resistance to complement. *Infect Immun* 78: 5262-5270.
 24. Moore MR, Gertz RE, Jr., Woodbury RL, Barkocy-Gallagher GA, Schaffner W, Lexau C, Gershman K, Reingold A, Farley M, Harrison LH, Hadler JL, Bennett NM, Thomas AR, Mcgee L, Pilishvili T, Brueggemann AB, Whitney CG, Jorgensen JH, Beall B (2008) Population snapshot of emergent *Streptococcus pneumoniae* serotype 19A in the United States, 2005. *J Infect Dis* 197: 1016-1027.
 25. Pai R, Gertz RE, Beall B (2006) Sequential multiplex PCR approach for determining capsular serotypes of *Streptococcus pneumoniae* isolates. *J Clin Microbiol* 44: 124-131.
 26. Pilishvili T, Lexau C, Farley MM, Hadler J, Harrison LH, Bennett NM, Reingold A, Thomas A, Schaffner W, Craig AS, Smith PJ, Beall BW, Whitney CG, Moore MR (2010) Sustained reductions in invasive pneumococcal disease in the era of conjugate vaccine. *J Infect Dis* 201: 32-41.
 27. Porat N, Amit U, Givon-Lavi N, Leibovitz E, Dagan R (2010) Increasing importance of multidrug-resistant serotype 6A *Streptococcus pneumoniae* clones in acute otitis media in southern Israel. *Pediatr Infect Dis J* 29: 126-130.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
28. Sa-Leao R, Nunes S, Brito-Avo A, Frazao N, Simoes AS, Crisostomo MI, Paulo AC, Saldanha J, Santos-Sanches I, de Lencastre H (2009) Changes in pneumococcal serotypes and antibiotypes carried by vaccinated and unvaccinated day-care centre attendees in Portugal, a country with widespread use of the seven-valent pneumococcal conjugate vaccine. *Clin Microbiol Infect* 15: 1002-1007.
 29. Shin J, Baek JY, Kim SH, Song JH, Ko KS (2011) Predominance of ST320 among *Streptococcus pneumoniae* serotype 19A isolates from 10 Asian countries. *J Antimicrob Chemother* 66: 1001-1004.
 30. Whitney CG, Pilishvili T, Farley MM, Schaffner W, Craig AS, Lynfield R, Nyquist AC, Gershman KA, Vazquez M, Bennett NM, Reingold A, Thomas A, Glode MP, Zell ER, Jorgensen JH, Beall B, Schuchat A (2006) Effectiveness of seven-valent pneumococcal conjugate vaccine against invasive pneumococcal disease: a matched case-control study. *Lancet* 368: 1495-1502.
 31. Yildirim I, Hanage WP, Lipsitch M, Shea KM, Stevenson A, Finkelstein J, Huang SS, Lee GM, Kleinman K, Pelton SI (2010) Serotype specific invasive capacity and persistent reduction in invasive pneumococcal disease. *Vaccine* 29: 283-288.

Table 1. Characteristics of children according to time periods, Hong Kong

	% (No/total)		
	Period 1 (n=1978)	Period 2 (n=2211)	<i>P</i>
Age, mean \pm SD (yr)	4.3 \pm 0.8	3.9 \pm 0.9	<0.001
Household size, mean \pm SD	4.5 \pm 1.3	4.4 \pm 1.2	0.5
Recent antibiotic use by			
subject	50.1 (883/1747)	30.6 (677/2211)	<0.001
household members	34.7 (616/1777)	27.0 (596/2211)	<0.001
Overcrowding	14.0 (277/1972)	4.8 (105/2183)	<0.001
Prior hospitalization	3.2 (63/1978)	7.5 (165/2211)	<0.001
Respiratory symptoms at time of sampling	25.6 (507/1978)	9.7 (214/2211)	<0.001
Has young siblings (\leq 5y)	27.6 (546/1978)	27.8 (614/2211)	0.9
Recent physician visit			
At least once	77.6 (1535/1978)	62.7 (1386/2211)	<0.01
Multiple (\geq 3 times)	23.7 (468/1931)	13.1 (289/2211)	<0.01

The time periods were 1999-2000 (period 1) and 2009-2010 (period 2). The data for period 1 was extracted from a previous work (Chiu et al., 2001) and those for period 2 were collected in the present study. Unless specified, the figures were number (%) of children. The number of participating institutes for period 1 and period 2 were 79 and 191, respectively.

Table 2. Comparison of the serotype distribution of nasopharyngeal isolates of *Streptococcus pneumoniae* collected in two time periods, Hong Kong

Serotype ^a	No. (%)		
	Period 1	Period 2	<i>P</i>
Total	383 (100)	347 (100)	-
PCV7			
4	0 (0)	0 (0)	-
6B	88 (23.0)	75 (21.6)	0.7
19F	69 (18.0)	62 (17.9)	0.9
23F	52 (13.6)	41 (11.8)	0.5
14	34 (8.9)	9 (2.6)	<0.01
18C	9 (2.3)	2 (0.6)	0.049
9V	1 (0.3)	1 (0.3)	0.9
PCV13-nonPCV7			
3	12 (3.1)	6 (1.7)	0.2
19A	0 (0)	8 (2.3)	<0.001
6A	13 (3.4)	41 (11.8)	<0.001
Non-PCV13			
6C	7 (1.8)	27 (7.8)	<0.001
23A	6 (1.6)	19 (5.5)	0.003
15B	1 (0.3)	9 (2.6)	0.007
Others ^b	91 (23.78)	47 (13.5)	-
Vaccine coverage			
PCV7	253 (66.1)	190 (54.8)	0.001
PCV10	253 (66.1)	190 (54.8)	0.001
PCV13	278 (72.6)	245 (70.6)	0.6

^aNo PCV10-nonPCV7 serotypes (1, 5 and 7F) were found in both periods. If serotype cross protection for 6B and 6A was assumed, the vaccine coverage for PCV7 and PCV10 in period 1/period 2 was both 69.5%/66.6%, respectively.

^bFor period 1, these included 7C (n=2), 8 (n=1), 13 (n=3), 11A (n=12), 15C (n=7), 15F (n=6), 19B (n=3), 22F (n=1), 28A/F (n=1), 29 (n=1), 33B/C/D (n=2), 34 (n=13), 35AC/42 (n=5), 35B (n=4), 35F (n=2) and nontypeable (n=28). For period 2, these included 8 (n=1), 10A (n=2), 11B/F (n=1), 11D (n=2), 15C (n=23), 15F (n=6), 22F (n=1), 23B (n=1), 29 (n=1), 33B/C/D (n=1), 34 (n=11), 35AC/42 (n=2), 35B (n=2), 35F (n=1), 37 (n=1) and 7C (n=1).

Table 3. Proportions of children carrying pneumococci within each sampling period, according to age

	% children with carriage			
	Period 1	Period 2	Difference (95% CI)	<i>P</i> value
PCV7 types				
2y	27.1	7.0	-20.1 (-9.3, -32.9)	<0.001
3y	22.7	9.1	-13.6 (-8.4, -19.2)	<0.001
4y	13.9	9.7	-4.2 (-0.5, -8.0)	0.02
5y	9.1	7.4	-	0.19
All ages	12.8	8.6	-4.2 (-2.3, -6.1)	<0.01
PCV13-nonPCV10				
2y	1.7	7.0	-	0.19
3y	2.4	1.6	-	0.43
4y	1.2	2.5	-	0.12
5y	1.0	2.1	-	0.05
All ages	1.3	2.5	1.2 (0.4, 2.1)	<0.01
Non-PCV13				
2y	0.0	2.9	-	0.33
3y	7.6	5.5	-	0.23
4y	5.0	4.7	-	0.85
5y	5.1	4.2	-	0.36
All ages	5.3	4.6	-	0.5

y, year

Table 4. Serotype distribution of nasopharyngeal isolates and their antimicrobial susceptibility according to serotype and time periods, Hong Kong.


Serotype	% Pen-NS ^{a, b}			% Ery-NS			% Pen/Ery-NS		
	Period 1	Period 2	<i>P</i>	Period 1	Period 2	<i>P</i>	Period 1	Period 2	<i>P</i>
All	58.2	70.6	<0.001	77.0	86.7	<0.001	52.5	70.3	<0.001
PCV7									
4	-	-	-	-	-	-	-	-	-
6B	84.1	77.3	0.3	95.5	90.7	0.4	81.8	77.3	0.5
19F	75.4	100	<0.001	95.7	100	0.2	75.4	100	<0.001
23F	88.5	78.0	0.2	92.3	82.9	0.2	86.5	78.0	0.3
14	55.9	100	0.01	97.1	100	1.0	55.9	100	0.01
18C	0	0	-	11.1	50	0.3	0	0	-
9V	0	0	-	100	100	-	0	0	-
PCV13-nonPCV7									
3	8.3	0	1.0	25.0	0	0.5	0	0	-
19A	-	75.0	-	-	87.5	-	-	75.0	-
6A	7.7	87.8	<0.001	76.9	95.1	0.08	7.7	87.8	<0.001
Non-PCV13									
6C	42.9	59.3	0.7	85.7	96.3	0.4	42.9	59.3	0.7
23A	0	78.9	0.001	100	100	-	0	78.9	0.001
15B	0	22.2	1.0	0	55.6	1.0	0	22.2	1.0
others	29.7	19.1	-	40.7	63.8	-	9.9	17.0	-

^a The percentages referred to the proportion of penicillin-nonsusceptible (Pen-NS, MIC >0.06 µg/ml), erythromycin-nonsusceptible (Ery-NS) or dual penicillin/erythromycin-nonsusceptible (Pen/Ery-NS) isolates within each serotype group.

^b The penicillin MIC₅₀ and MIC₉₀ for all the isolates were 0.5 µg/ml and 2 µg/ml, respectively. Forty-nine isolates had penicillin MIC 4 µg/ml. They belonged to 19F (n=35), 23F (n=9), 6B (n=2), 14 (n=1), 15C (n=1) and 19A (n=1). Two isolates had penicillin MIC 8 µg/ml (both 19F).

Figure 1. Relationship between serotype-specific prevalence and penicillin-nonsusceptibility rates for (A) period 1 and (B) period 2. The line showed regression result according to the linear model.

(A)



(B)

