

NEAR INFRARED LASER SPECTROSCOPY OF SCANDIUM MONOBROMIDE

YE XIA, A. S.-C. CHEUNG, *Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong*; ZHENWU LIAO, MEI YANG, MAN-CHOR CHAN, *Department of Chemistry, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong*.

High resolution laser spectrum of scandium monobromide (ScBr) between 787 and 845 nm has been investigated using the technique of laser vaporization/reaction with free jet expansion and laser induced fluorescence spectroscopy. ScBr was produced by reacting laser vaporized Sc atoms with ethyl bromide (C_2H_5Br). Spectra of six vibrational bands of both $Sc^{79}Br$ and $Sc^{81}Br$ isotopomers of the $C^1\Sigma^+ - X^1\Sigma^+$ transition and seven vibrational bands of the $e^3\Delta - a^3\Delta$ transition were obtained and analyzed. Least-squares fit of the measured line positions for the singlet transitions yielded accurate molecular constants for the $v = 0 - 3$ levels of the $C^1\Sigma^+$ state and the $v = 0 - 2$ levels of the $X^1\Sigma^+$ state. Similar least-squares fit for the triplet transitions yielded molecular constants for the $v = 0 - 2$ levels of both $e^3\Delta$ and $a^3\Delta$ states. The equilibrium bond length, r_0 , of the $a^3\Delta$ state has been determined to be 2.4789 Å. Financial support from the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No. HKU 701008P) is gratefully acknowledged