

ON CYCLE INTEGRALS OF WEAKLY HOLOMORPHIC MODULAR FORMS

KATHRIN BRINGMANN, PAVEL GUERZHOY, AND BEN KANE

ABSTRACT. In this paper, we investigate cycle integrals of weakly holomorphic modular forms. We show that these integrals coincide with the cycle integrals of classical cusp forms. We use these results to define a Shintani lift from integral weight weakly holomorphic modular forms to half-integral weight holomorphic modular forms.

1. INTRODUCTION AND STATEMENT OF RESULTS

The relationship between integral and half-integral weight modular forms has been pivotal in a number of applications. The link between these spaces was first established in a groundbreaking paper of Shimura [17] in which he constructed lifts from half-integral to integral weight modular forms. The level of the image of Shimura's lifts was then studied by a number of authors. Niwa [16] used theta lifts to find the optimal level of the lifts in general, while Kohnen [12] found a distinguished subspace of half-integral weight modular forms (known as Kohnen's plus space) which map to a lower level. A number of general results have a much more explicit form when restricted to Kohnen's plus space. For example, Waldspurger [21] used Shimura's lifts to establish connections between central values of L -functions of integral weight modular forms and coefficients of corresponding half-integral weight modular forms. By restricting to Kohnen's plus space, Kohnen and Zagier [13] refined Waldspurger's

Date: April 29, 2014.

2010 Mathematics Subject Classification. 11F37, 11F11.

Key words and phrases. Cycle integrals, weakly holomorphic modular forms, harmonic weak Maass forms, Shintani lift, Shimura lift.

The research of the first author was supported by the Alfred Krupp Prize for Young University Teachers of the Krupp Foundation and by the Deutsche Forschungsgemeinschaft (DFG) Grant No. BR 4082/3-1. The research of the second author was partially supported by a Simons Foundation Collaboration grant (209914 to P. Guerzhoy). Part of this research was conducted while the third author was a postdoc at the University of Cologne.

formula [21] to determine an explicit constant which proved non-negativity of the central values of these L -functions. Waldspurger's formula [21] (and Kohnen and Zagier's variant [13]) also yields a link between the vanishing of central L -values of integral weight modular forms and coefficients of half-integral weight modular forms. Specifically, for each fundamental discriminant D and integral weight Hecke eigenform f , the central value of the D th twist of the L -function of f vanishes if and only if the $|D|$ th coefficient of a half-integral weight modular form related to f by the Shimura correspondence is zero.

More recently, Bruinier and Ono [5] proved that the central values of derivatives of L -functions vanish precisely when certain coefficients of a newer modular object, known as harmonic weak Maass forms, are algebraic.

Central to the theory built around integral and half-integral weight modular forms is a two-sided interplay between the Shimura lifts and lifts of Shintani [18] which map integral weight to half-integral weight modular forms. The Shintani lifts are defined in terms of certain cycle integrals, which are, roughly speaking, integrals of modular objects along geodesics defined by certain integral binary quadratic forms of positive discriminant. In particular, for a continuous function F satisfying weight $2\kappa \in 2\mathbb{Z}$ modularity, the coefficients of the Shintani lifts are given as traces of cycle integrals of the type

$$\mathcal{C}(F; Q) := \int_{C_Q} F(z) Q(z, 1)^{\kappa-1} dz, \quad (1.1)$$

where Q is an integral binary quadratic form of positive discriminant and C_Q is explicitly defined in Section 2. In [2], the authors related two families of cycle integrals coming from weight $2 - 2k$ ($k \in \mathbb{N}$ fixed throughout) harmonic weak Maass forms \mathcal{M} . One family is formed with the traces of cycle integrals defining the δ th Shintani lift of the weight $2k$ cusp form $\xi_{2-2k}(\mathcal{M})$ (where $\xi_{2-2k} := 2iy^{2-2k} \frac{\partial}{\partial \bar{z}}$ and δ is a fundamental discriminant for which $(-1)^k \delta > 0$). These cycle integrals were shown to be fundamentally equal to the traces of cycle integrals of a (non-holomorphic) weight 0 Maass form which is essentially built out of the $(k - 1)$ th (holomorphic) derivative of \mathcal{M} . However, if one instead takes $2k - 1$ derivatives of \mathcal{M} , then the resulting function is a weight $2k$ *weakly holomorphic modular form* (i.e., a meromorphic modular form whose only possible poles occur at the cusps). Cycle integrals of weakly holomorphic modular forms are well-defined and were first considered in [9], where an exciting connection between such cycle integrals and rational period functions was established. Since $\xi_{2-2k}(\mathcal{M})$ and $\mathcal{D}^{2k-1}(\mathcal{M})$ (where $\mathcal{D} := \frac{1}{2\pi i} \frac{\partial}{\partial z}$) are both weight $2k$ weakly holomorphic modular forms, it is natural to compare their cycle integrals.

Defining a regularization, which is necessary to guarantee convergence of (1.1) in the case that the discriminant of Q is a square, our first result is that these integrals in essence coincide for every element \mathcal{M} in the space H_{2-2k}^{cusp} of harmonic weak Maass forms for which $\xi_{2-2k}(\mathcal{M})$ is a cusp form.

Theorem 1.1. *For every harmonic weak Maass form $\mathcal{M} \in H_{2-2k}^{\text{cusp}}$, every discriminant $D > 0$, and every $Q \in \mathcal{Q}_D$, the cycle integrals satisfy*

$$\mathcal{C}(\xi_{2-2k}(\mathcal{M}); Q) = -\frac{(4\pi)^{2k-1}}{(2k-2)!} \overline{\mathcal{C}(\mathcal{D}^{2k-1}(\mathcal{M}); Q)}. \quad (1.2)$$

Note that the kernel of ξ_{2-2k} (namely, the space $M_{2-2k}^!$ of weakly holomorphic modular forms) is non-trivial. In the special case that $\mathcal{M} \in M_{2-2k}^!$, Theorem 1.1 therefore immediately yields the following corollary.

Corollary 1.2. *If $G \in M_{2-2k}^!$, then*

$$\mathcal{C}(\mathcal{D}^{2k-1}(G); Q) = 0.$$

By taking a trace of the cycle integrals, we next define a Shintani lift for $f \in S_{2k}^!$, the space of *weakly holomorphic cusp forms* (i.e., those $f \in M_{2k}^!$ whose constant terms vanish). For δ given before Theorem 1.1 and $f \in S_{2k}^!$, this Shintani lift is given by

$$\mathcal{S}_\delta^*(f)(z) := \sum_{\substack{m>0 \\ (-1)^k m \equiv 0,1 \pmod{4}}} \sum_{Q \in \mathcal{Q}_{|\delta|m}/\text{SL}_2(\mathbb{Z})} \chi_\delta(Q) \mathcal{C}(f; Q) q^m, \quad (1.3)$$

where the genus character χ_δ is defined in (2.3) and $q := e^{2\pi iz}$. We obtain a relation between (1.3) and the classical δ th Shintani lift. If $f \in S_{2k}$, then a theorem of Shintani [18] implies that $\mathcal{S}_\delta^*(f)$ is an element of Kohnen's plus space [12] $\mathbb{S}_{k+\frac{1}{2}}$ of weight $k + \frac{1}{2}$ cusp forms.

We prove that this extends to $S_{2k}^!$ and include the relation between the weight $2k$ Eisenstein series G_{2k} (see (4.1)) and Cohen's weight $k + \frac{1}{2}$ Eisenstein series \mathcal{H}_k (defined in (4.3)) for completeness.

Theorem 1.3. *Suppose that δ is a fundamental discriminant satisfying $(-1)^k \delta > 0$.*

- (1) *If $f \in S_{2k}^!$, then $\mathcal{S}_\delta^*(f) \in \mathbb{S}_{k+\frac{1}{2}}$.*
- (2) *If $\mathcal{M} \in H_{2-2k}^{\text{cusp}}$, then*

$$\mathcal{S}_\delta^*(\xi_{2-2k}(\mathcal{M})) = -\frac{(4\pi)^{2k-1}}{(2k-2)!} (\mathcal{S}_\delta^*(\mathcal{D}^{2k-1}(\mathcal{M})))^c, \quad (1.4)$$

where for $g \in \mathbb{S}_{k+\frac{1}{2}}$ we use $g^c(z) := \overline{g(-\bar{z})}$ for any function g on the upper half-plane.

(3) If $\delta = 1$, then we have

$$\mathcal{S}_1^*(G_{2k}(z)) = \frac{1}{2}\zeta(1-k)\mathcal{H}_k(z).$$

Remarks.

- (1) If $\delta = 1$, then (1.4) holds for all harmonic weak Maass forms.
- (2) We expect that a twisted version of Theorem 1.3 (3) holds for general fundamental discriminants δ satisfying $(-1)^k\delta > 0$.
- (3) One can prove that for $\delta \neq 1$ and $f \in S_{2k}^!$, the Shintani lift $\mathcal{S}_\delta^*(f)$ is not influenced by the even periods of f . That is to say, if $f_1, f_2 \in S_{2k}^!$ have the same odd periods, then $\mathcal{S}_\delta^*(f_1) = \mathcal{S}_\delta^*(f_2)$. Computational evidence further indicates that in this case $\mathcal{C}(f_1; Q) = \mathcal{C}(f_2; Q)$ for every $Q \in \mathcal{Q}_D$ and every $D > 0$. Since we do not need this for our purposes, we omit the proof here.
- (4) In [10] another geometric extension of the Shintani lift was given, which for Eisenstein series coincides with our definition (1.3). Theorem 1.3 (3) can also directly be concluded from Section 9 of [10].
- (5) The restriction that δ is fundamental may be omitted by using the action of the Hecke operators to define the Shintani lift for non-fundamental discriminants, but we do not work out the details here.

The paper is organized as follows. In Section 2, we introduce the regularization necessary to define the cycle integrals for square discriminants and relate the cycle integrals for non-square discriminants to the periods of weakly holomorphic modular forms. In Section 3 we prove Theorem 1.1. Finally, we prove Theorem 1.3 in Section 4.

ACKNOWLEDGEMENTS

The authors thank Jens Funke for helpful discussion.

2. CYCLE INTEGRALS AND PERIODS

In this section, we recall known facts on cycle integrals and periods of weakly holomorphic modular forms and additionally prove a relation between them. The generalization of these definitions to weakly holomorphic modular forms requires a regularized integral defined by the first author, Fricke, and Kent [1]. Assume that

$f : \mathbb{H} \rightarrow \mathbb{C}$ is a continuous function and that there exists $c \in \mathbb{R}^+$ such that f satisfies the growth condition

$$f(z) = O\left(e^{c\operatorname{Im}(z)}\right) \quad (2.1)$$

uniformly in $\operatorname{Re}(z)$ as $\operatorname{Im}(z) \rightarrow \infty$. Then, for each $z_0 \in \mathbb{H}$, the integral

$$\int_{z_0}^{i\infty} e^{uiv} f(w) dw$$

(where the path of integration lies within a vertical strip) is convergent for $u \in \mathbb{C}$ with $\operatorname{Re}(u) \gg 0$. If this integral has an analytic continuation to $u = 0$, then the *regularized integral* of f is given by

$$R. \int_{z_0}^{i\infty} f(w) dw := \left[\int_{z_0}^{i\infty} e^{uiv} f(w) dw \right]_{u=0},$$

where the right-hand side means that we take the value at $u = 0$ of the analytic continuation of the integral. We similarly define integrals at other cusps \mathfrak{a} . Specifically, suppose that $\mathfrak{a} = \sigma_{\mathfrak{a}}(i\infty)$ for a scaling matrix $\sigma_{\mathfrak{a}} \in \operatorname{SL}_2(\mathbb{Z})$. If $f(\sigma_{\mathfrak{a}}z)$ satisfies (2.1), then we use the regularization

$$R. \int_{z_0}^{\mathfrak{a}} f(w) dw := R. \int_{\sigma_{\mathfrak{a}}^{-1}z_0}^{i\infty} f|_2\sigma_{\mathfrak{a}}(w) dw.$$

For cusps $\mathfrak{a}, \mathfrak{b}$, we set

$$R. \int_{\mathfrak{a}}^{\mathfrak{b}} f(w) dw := R. \int_{z_0}^{\mathfrak{b}} f(w) dw + R. \int_{\mathfrak{a}}^{z_0} f(w) dw \quad (2.2)$$

for any $z_0 \in \mathbb{H}$. It is not hard to see that this integral is independent of $z_0 \in \mathbb{H}$.

This regularization can be used to generalize the classical definition of periods of cusp forms. To be more precise, the n th periods ($n \in \mathbb{N}_0$) for *weakly holomorphic cusp forms* $f \in S_{2k}^!$ (those weakly holomorphic modular forms whose constant terms vanish) are given by

$$r_n(f) := R. \int_0^{\infty} f(it)t^n dt,$$

which may be packaged into *period polynomials*

$$r(f; z) := \sum_{n=0}^{2k-2} i^{-n+1} \binom{2k-2}{n} r_n(f) z^{k-2-n}.$$

The first two authors, Kent, and Ono (see (1.7) and (1.9) of [3]) proved that the period polynomials of $f(z) = \sum_{n \in \mathbb{Z}} a_n q^n \in S_{2k}^!$ are constant multiples of the error to modularity under $S := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ of the (holomorphic) Eichler integrals

$$\mathcal{E}_f(z) := \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{a_n}{n^{2k-1}} q^n.$$

For $t_0 > 0$, $c \in \mathbb{Z}$, and $d \in \mathbb{N}$, we also require the twisted L -series

$$L_f^*(\zeta_c^d, s) := \sum_{m \geq m_0} \frac{a_f(m) \zeta_c^{dm} \Gamma(s, 2\pi m t_0)}{(2\pi m)^s} + i^k c^{k-2s} \sum_{m \geq m_0} \frac{a_f(m) \zeta_c^{-am} \Gamma\left(k - s, \frac{2\pi m}{c^2 t_0}\right)}{(2\pi m)^{k-s}}.$$

Here for $y > 0$, $\Gamma(s, y) := \int_y^\infty t^{s-1} e^{-t} dt$ is the *incomplete* Γ -function, and the definition can be shown to be independent of the choice of t_0 .

Before clarifying the role played by the above regularization if the discriminant is a square, we formally define the cycle integrals given in (1.1). For each non-square discriminant $D > 0$, denote the set of integral binary quadratic forms of discriminant D by \mathcal{Q}_D . If D is not a square, then the (infinite cyclic) group of automorphs of an indefinite binary quadratic form $Q = [a, b, c] \in \mathcal{Q}_D$ is defined by

$$g_Q := \begin{pmatrix} \frac{t+bu}{2} & cu \\ -au & \frac{t-bu}{2} \end{pmatrix},$$

where (t, u) is the smallest positive solution to the Pell equation $t^2 - Du^2 = 4$ (the automorphs are trivial if D is a square, and we let g_Q be the identity in this case). For $a \neq 0$ (resp. $a = 0$), let S_Q be the oriented semicircle (resp. vertical line) given by

$$a|z|^2 + bx + c = 0$$

directed counterclockwise if $a > 0$, clockwise if $a < 0$, and up from the real axis if $a = 0$. Note that $\frac{dz}{Q(z, 1)}$ is an invariant measure on S_Q . Since every $Q \in \mathcal{Q}_D$ is modular of weight -2 for $\Gamma_Q := \langle g_Q \rangle$, for $\kappa \in \mathbb{Z}$ it is natural to integrate the product of $Q(z, 1)^{\kappa-1}$ times functions F satisfying weight $2\kappa \in 2\mathbb{Z}$ modularity on Γ_Q along $\Gamma_Q \setminus S_Q$. Hence, for $z \in S_Q$, we define C_Q to be the directed arc from z to $g_Q z$ along S_Q if D is non-square and $C_Q := S_Q$ if D is a square. As in Lemma 6 of [8], one concludes that the cycle integral in (1.1) is well-defined for any continuous function F satisfying weight $2k \in \mathbb{Z}$ modularity on Γ_Q for which the integral converges.

More generally, in the case of convergence, we extend the definition of the cycle integrals in (1.1) by

$$\mathcal{C}(F; Q) := R. \int_{C_Q} F(z) Q(z, 1)^{k-1} dz.$$

In particular, this is necessary for square D whenever F does not decay towards the cusps. Hence this regularization allows one to take cycle integrals of *harmonic weak Maass forms*. Denoting the weight $2 - 2k$ *hyperbolic Laplacian* by

$$\Delta_{2-2k} := -y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + i(2 - 2k)y \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right),$$

these are functions $\mathcal{M} : \mathbb{H} \rightarrow \mathbb{C}$ satisfying:

- (i) $\mathcal{M}|_{2-2k}\gamma = \mathcal{M}$ for all $\gamma \in \mathrm{SL}_2(\mathbb{Z})$,
- (ii) $\Delta_{2-2k}(\mathcal{M}) = 0$,
- (iii) \mathcal{M} has at most linear exponential growth at $i\infty$.

The subspace of harmonic weak Maass forms \mathcal{M} for which $\xi_{2-2k}(\mathcal{M})$ is a cusp form is denoted by H_{2-2k}^{cusp} . We aim to compare the cycle integrals of the weakly holomorphic modular forms $\mathcal{C}(\xi_{2-2k}(\mathcal{M}); Q)$ and $\overline{\mathcal{C}(\mathcal{D}^{2k-1}(\mathcal{M}); Q)}$ for $\mathcal{M} \in H_{2-2k}^{\mathrm{cusp}}$.

Having established the definition of cycle integrals for weakly holomorphic modular forms, we have almost all of the notation required for the definition of the Shintani lift (1.3). To complete the definition, recall that for every pair of discriminants D_1 and D_2 , the corresponding *genus character* (for example, see pp. 59–62 of [19]) of a binary quadratic form $Q(X, Y) = [a, b, c](X, Y) := aX^2 + bXY + cY^2 \in \mathcal{Q}_{D_1 D_2}$ is given by

$$\chi_{D_1}(Q) := \begin{cases} \left(\frac{D_1}{r}\right) & \text{if } (a, b, c, D_1) = 1 \text{ and } Q \text{ represents } r \text{ with } (r, D_1) = 1, \\ 0 & \text{if } (a, b, c, D_1) > 1. \end{cases} \quad (2.3)$$

Here $\left(\frac{D_1}{\cdot}\right)$ denotes the Kronecker character.

We make frequent use of a generalization of Theorem 7 of [14] to include weakly holomorphic modular forms. To state the theorem, we require some notation. A binary quadratic form $Q = [a, b, c] \in \mathcal{Q}_D$ ($D > 0$) is called *reduced* if $a > 0$, $c > 0$, and $b > a + c$. Let \mathcal{A} be an $\mathrm{SL}_2(\mathbb{Z})$ -equivalence class of integral binary quadratic forms of discriminant $D > 0$. We enumerate the reduced forms in \mathcal{A} by $Q_0, \dots, Q_r = Q_0$, where

$$Q_j = Q_{j-1} \circ M_j := Q_{j-1}|_{-2} M_j, \quad (2.4)$$

with

$$M_j := \begin{pmatrix} m_j & 1 \\ -1 & 0 \end{pmatrix} \quad (2.5)$$

for some integers $m_j \geq 2$. Here $|-_2$ is the usual weight -2 slash operator and we use $Q_{j-1}|_{-2}$ to abbreviate the action on the function $Q_{j-1}(z, 1)$. We use the reduced forms to define the polynomial

$$Q_{k,D,\mathcal{A}}(X) := \sum_{\substack{Q \in \mathcal{A} \\ Q \text{ reduced}}} Q(X, -1)^{k-1} = \sum_{j=1}^r Q_j(X, -1)^{k-1}$$

and denote the coefficient of X^n by $q_{k,D,\mathcal{A}}^{(n)}$. We are now ready to give the relation between cycle integrals and periods, which was shown for cusp forms in [14].

Theorem 2.1. *For each equivalence class \mathcal{A} of binary quadratic forms of non-square discriminant containing $Q \in \mathcal{A}$ and $f \in S_{2k}^!$, one has*

$$\mathcal{C}(f; Q) = \sum_{n=0}^{2k-2} i^{-n+1} q_{k,D,\mathcal{A}}^{(n)} r_n(f). \quad (2.6)$$

Proof. We rewrite the right-hand side of (2.6), using the definition of r_n

$$\begin{aligned} \sum_{n=0}^{2k-2} i^{-n+1} q_{k,D,\mathcal{A}}^{(n)} r_n(f) &= \sum_{n=0}^{2k-2} (-1)^n q_{k,D,\mathcal{A}}^{(n)} R \cdot \int_0^{i\infty} f(z) z^n dz \\ &= R \cdot \int_0^{i\infty} f(z) Q_{k,D,\mathcal{A}}(-z) dz = R \cdot \sum_{j=1}^r \int_0^{i\infty} f(z) Q_{j-1}(z, 1)^{k-1} dz. \end{aligned} \quad (2.7)$$

On the left-hand side of (2.6), we follow the formal argument at the beginning of the proof in [14] mutatis mutandis to obtain

$$\mathcal{C}(f; Q) = \sum_{j=1}^r \int_{z_0}^{M_j z_0} f(z) Q_{j-1}(z, 1)^{k-1} dz.$$

However, for every $z_0 \in \mathbb{H}$, continuity in u implies that

$$\left[\sum_{j=1}^r \int_{z_0}^{M_j z_0} f(z) e^{uiw} Q_{j-1}(z, 1)^{k-1} dz \right]_{u=0} = \sum_{j=1}^r \int_{z_0}^{M_j z_0} f(z) Q_{j-1}(z, 1)^{k-1} dz. \quad (2.8)$$

Since $\mathcal{C}(f; Q)$ is independent of z_0 , the left-hand side of (2.8) is hence also independent. We then take $z_0 \rightarrow 0$, precisely yielding the right-hand side of (2.7). \square

3. PROOF OF THEOREM 1.1

The argument to prove Theorem 1.1 naturally separates for D square and non-square. We begin with the case that D is not a square and first prove the existence of $\mathcal{M} \in H_{2-2k}^{\text{cusp}}$ for which (1.2) holds.

Proposition 3.1. *For every $g \in S_{2k}$, there exists $\mathcal{M} \in H_{2-2k}^{\text{cusp}}$ for which $\xi_{2-2k}(\mathcal{M}) = g$ such that for every non-square discriminant D and every $Q \in \mathcal{Q}_D$, (1.2) holds.*

Proof. We may choose \mathcal{M} as in the second statement of Theorem 3.1 of [1], namely satisfying

$$r(\xi_{2-2k}(\mathcal{M}); z) = -\frac{(4\pi)^{2k-1}}{(2k-2)!} \overline{r(\mathcal{D}^{2k-1}(\mathcal{M}); z)}.$$

By comparing coefficients of z^n , one obtains

$$r_n(\xi_{2-2k}(\mathcal{M})) = (-1)^n \frac{(4\pi)^{2k-1}}{(2k-2)!} \overline{r_n(\mathcal{D}^{2k-1}(\mathcal{M}))}.$$

For every $Q \in \mathcal{A}$, we then use Theorem 2.1 to yield

$$\begin{aligned} \mathcal{C}(\xi_{2-2k}(\mathcal{M}); Q) &= \sum_{n=0}^{2k-2} i^{-n+1} q_{k,D,\mathcal{A}}^{(n)} r_n(\xi_{2-2k}(\mathcal{M})) \\ &= \frac{(4\pi)^{2k-1}}{(2k-2)!} \sum_{n=0}^{2k-2} i^{n+1} q_{k,D,\mathcal{A}}^{(n)} \overline{r_n(\mathcal{D}^{2k-1}(\mathcal{M}))} \\ &= -\frac{(4\pi)^{2k-1}}{(2k-2)!} \sum_{n=0}^{2k-2} i^{-n+1} q_{k,D,\mathcal{A}}^{(n)} r_n(\mathcal{D}^{2k-1}(\mathcal{M})) = -\frac{(4\pi)^{2k-1}}{(2k-2)!} \overline{\mathcal{C}(\mathcal{D}^{2k-1}(\mathcal{M}); Q)}. \end{aligned}$$

This gives the claim. \square

Proposition 3.1 shows that (1.2) holds for at least one $\mathcal{M} \in H_{2-2k}^{\text{cusp}}$ which maps to $g \in S_{2k}$ under ξ_{2-2k} and the difference of any two such elements of H_{2-2k}^{cusp} is weakly holomorphic. To show the non-square case it hence remains to prove Corollary 1.2 for D non-square, which we now state as a separate proposition.

Proposition 3.2. *If $G \in M_{2-2k}^1$ and $Q \in \mathcal{Q}_D$ with D non-square, then*

$$\mathcal{C}(\mathcal{D}^{2k-1}(G); Q) = 0.$$

Before proving Proposition 3.2, we show a useful lemma.

Lemma 3.3. *For every $k \geq 2$ and discriminant $D > 0$ we have*

$$q_{k,D,\mathcal{A}}^{(0)} = q_{k,D,\mathcal{A}}^{(2k-2)}. \quad (3.1)$$

Proof. Denote the reduced forms $Q_j = [a_j, b_j, c_j]$ as in (2.4). The leading term of

$$Q_{k,D} \Big|_{2-2k} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (X) = \sum_{j=1}^r Q_j(1, -X)^{k-1}$$

is then

$$\sum_{j=1}^r c_j^{k-1} = q_{k,D,\mathcal{A}}^{(2k-2)}. \quad (3.2)$$

However, since $Q_j = Q_{j-1} \Big|_{-2} M_j = Q_{j-1} \circ M_j$ by (2.4) (where M_j is defined in (2.5)), we have

$$\begin{aligned} Q_{k,D} \Big|_{2-2k} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (X) &= \sum_{j=1}^r Q_{j-1} \circ \begin{pmatrix} m_j & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (X, 1)^{k-1} \\ &= \sum_{j=1}^r Q_{j-1} \circ \begin{pmatrix} -1 & m_j \\ 0 & -1 \end{pmatrix} (X, 1)^{k-1} = \sum_{j=1}^r Q_{j-1}(m_j - X, -1)^{k-1} \\ &= \sum_{j=1}^r Q_j(X - m_{j+1}, 1)^{k-1}. \end{aligned}$$

Since translation does not change the leading coefficient, we conclude that the leading coefficient is

$$\sum_{j=1}^r a_j^{k-1} = q_{k,D,\mathcal{A}}^{(0)}.$$

Comparing with (3.2) completes the proof of the lemma. \square

Proof of Proposition 3.2. For weakly holomorphic forms $f \in \mathcal{D}^{2k-1}(M_{2-2k}^!)$, one may greatly simplify (2.6). In this case, $r_n(f) = 0$ for $0 < n < 2k - 2$. Moreover, it is known [3, 14] that for every $f \in S_{2k}^!$ one has the relations

$$i^{-n+1}r_n(f) = (-1)^{n+1}i^{-(2k-2-n)+1}r_{2k-2-n}(f). \quad (3.3)$$

Hence in particular

$$i^1r_0(f) = -i^{-(2k-2)+1}r_{2k-2}(f).$$

Plugging this in, (2.6) becomes

$$\mathcal{C}(f; Q) = ir_0(f) \left(q_{k,D,\mathcal{A}}^{(0)} - q_{k,D,\mathcal{A}}^{(2k-2)} \right).$$

We finally use Lemma 3.3 to obtain $\mathcal{C}(f; Q) = 0$. □

We next consider the square case.

Proposition 3.4. *If D is a square, then (1.2) holds for every $\mathcal{M} \in H_{2-2k}^{cusp}$ and $Q \in \mathcal{Q}_D$.*

Proof. For the equivalence classes of quadratic forms of discriminant D , we may take as a set of representatives (cf. Lemma 3.31 and Lemma 3.32 of [15] for a modern version of Gauss's theory [11])

$$\left\{ [0, \sqrt{D}, c] \mid 0 \leq c < \sqrt{D} \right\}.$$

We hence have

$$\mathcal{C}(\xi_{2-2k}(\mathcal{M}); Q) = R \int_{-\frac{c}{\sqrt{D}}}^{i\infty} \xi_{2-2k}(\mathcal{M})(z) \left(\sqrt{D}z + c \right)^{k-1} dz. \quad (3.4)$$

The right-hand side of (3.4) was evaluated in Theorems 4.2 and 4.3 of [1] as

$$\begin{aligned} & i^{-k} D^{\frac{k-1}{2}} L_{\xi_{2-2k}(\mathcal{M})}^* \left(\zeta_{-c}^{\sqrt{D}}, k \right) \\ &= -\frac{(4\pi)^{2k-1}}{(2k-2)!} i^k D^{\frac{k-1}{2}} \overline{L_{\mathcal{D}^{2k-1}(\mathcal{M})}^* \left(\zeta_{-c}^{\sqrt{D}}, k \right)} = -\frac{(4\pi)^{2k-1}}{(2k-2)!} \overline{\mathcal{C}(\mathcal{D}^{2k-1}(\mathcal{M}); Q)}, \end{aligned}$$

where in the last equality we again use Theorem 4.2 of [1] and then reverse the argument in (3.4). □

4. PROOF OF THEOREM 1.3

In this section, we concentrate on the Shintani lifts defined in (1.3). Using Theorem 1.1 to relate the Shintani lift on $S_{2k}^!$ to the classical Shintani lift on S_{2k} , in order to compute the Shintani lifts of all weakly holomorphic modular forms, it remains to determine the cycle integrals of the weight $2k$ Eisenstein series

$$G_{2k}(z) := -\frac{B_{2k}}{4k} + \sum_{n=1}^{\infty} \sigma_{2k-1}(n)q^n, \quad (4.1)$$

where B_ℓ is the ℓ th Bernoulli number and $\sigma_s(n) := \sum_{d|n} d^s$. Kohnen and Zagier (see pages 240–241 of [14]) related the cycle integrals of Eisenstein series to the zeta functions

$$\zeta_{[Q]}(s) := \sum_{\substack{(m,n) \in \Gamma_Q \backslash \mathbb{Z}^2 \\ Q(m,n) > 0}} Q(m,n)^{-s}.$$

Note that these series converge for $\operatorname{Re}(s)$ sufficiently large and are independent of the representatives Q of $\mathcal{Q}_D/\operatorname{SL}_2(\mathbb{Z})$. The relation between the cycle integrals of Eisenstein series and $\zeta_{[Q]}(s)$ is given in the following lemma.

Lemma 4.1. *For every $Q \in \mathcal{Q}_D$, one has*

$$\mathcal{C}(G_{2k}; Q) = \frac{1}{2}(-1)^k \zeta_{[Q]}(1-k). \quad (4.2)$$

The image of G_{2k} under \mathcal{S}_1^* is well-known to be a constant multiple of Cohen's [7] weight $k + \frac{1}{2}$ Eisenstein series (this statement is essentially contained in [14], but we add the details below for the convenience of the reader)

$$\mathcal{H}_k(z) := \zeta(1-2k) + \sum_{\substack{m \in \mathbb{N} \\ (-1)^k m \equiv 0,1 \pmod{4}}} H(k, (-1)^k m) q^m, \quad (4.3)$$

where for $D = D_0 f^2$ with a fundamental discriminant D_0 and $f \in \mathbb{N}$ we define (μ is the usual Möbius function)

$$H(k, D) := L\left(1-k, \left(\frac{D_0}{\cdot}\right)\right) \sum_{d|f} \mu(d) \left(\frac{D_0}{d}\right) d^{k-1} \sigma_{2k-1}\left(\frac{f}{d}\right).$$

Combining the fact that G_{2k} maps to a modular form under \mathcal{S}_1^* with Theorem 2.1, we are able to prove that the Shintani lift (1.3) maps weakly holomorphic modular forms to modular forms.

Proof of Theorem 1.3. Using Theorem 1.1 termwise immediately implies that (1.4) holds for every $\mathcal{M} \in H_{2-2k}^{\text{cusp}}$, yielding (2).

Since ξ_{2-2k} is surjective [4] and g^c is a cusp form if g is a cusp form, (1) then follows from (2) and the classical results of Shintani [18] for cusp forms.

In order to prove (3), we use the following identity of Siegel [20]:

$$\sum_{Q \in \mathcal{Q}_D / \text{SL}_2(\mathbb{Z})} \zeta_{[Q]}(1-k) = \zeta(1-k)H(k, D).$$

Combining this with Lemma 4.1 and noting that $\chi_1(Q) = 1$ for every Q yields the third statement of the theorem. \square

REFERENCES

- [1] K. Bringmann, K. Fricke, and Z. Kent, *Special L-values and periods of weakly holomorphic modular forms*, Proc. Amer. Math. Soc., accepted for publication.
- [2] K. Bringmann, P. Guerzhoy, and B. Kane, *Shintani lifts and fractional derivatives for harmonic weak Maass forms*, Adv. Math., accepted for publication.
- [3] K. Bringmann, P. Guerzhoy, Z. Kent, and K. Ono, *Eichler-Shimura theory for mock modular forms*, Math. Ann. **355** (2013), 1085–1121.
- [4] J. Bruinier and J. Funke, *On two geometric theta lifts*, Duke Math. J. **125** (2004), no. 1, 45–90.
- [5] J. Bruinier and K. Ono, *Algebraic formulas for the coefficients of half-integral weight harmonic weak Maass forms*, Adv. Math. **246** (2013), 198–219.
- [6] Y. Choie and D. Zagier, *Rational period functions for $\text{PSL}_2(\mathbb{Z})$* , in “A Tribute to Emil Grosswald: Number Theory and Related Analysis”, Contemporary Math. **143** (1993), 89–107.
- [7] H. Cohen, *Sums involving the values at negative integers of L-functions of quadratic characters*, Math. Ann. **217** (1975), 271–285.
- [8] W. Duke, Ö. Imamoglu, and Á. Tóth, *Cycle integrals of the j-function and mock modular forms*, Ann. of Math. **173** (2011), 947–981.
- [9] W. Duke, Ö. Imamoglu, and Á. Tóth, *Rational period functions and cycle integrals*, Abh. Math. Semin. Univ. Hambg. **80** (2010), 255–264.
- [10] J. Funke and J. Millson, *Spectacle cycles with coefficients and modular forms of half-integral weight* in Arithmetic geometry and automorphic forms, Adv. Lect. Math. **19** (2011), 91–154.
- [11] C. Gauss, *Disquisitiones Arithmeticae*, Leipzig, 1801.
- [12] W. Kohnen, *Modular forms of half-integral weight on $\Gamma_0(4)$* , Math. Ann. **248** (1980), 249–266.
- [13] W. Kohnen and D. Zagier, *Values of L-series of modular forms at the center of the critical strip*, Invent. Math. **64** (1981), 175–198.
- [14] W. Kohnen and D. Zagier, *Modular forms with rational periods* in “Modular Forms,” ed. by R. A. Rankin, Ellis Horwood (1984), 197–249.
- [15] F. Lemmermeyer, *Binary quadratic forms: An elementary approach to the arithmetic of elliptic and hyperelliptic curves*, book to appear in 2014.

- [16] S. Niwa, *Modular forms of half integral weight and the integral of certain theta-functions*, Nagoya Math. J. **56** (1974), 174–161.
- [17] G. Shimura, *On modular forms of half integral weight*, Ann. of Math. **97** (1973), 440–481.
- [18] T. Shintani, *On construction of holomorphic cusp forms of half integral weight*, Nagoya Math. J. **58** (1975), 83–126.
- [19] C. Siegel, *Advanced analytic number theory*, 2nd ed., Tata Institute of Fundamental Research Studies in Mathematics, 9, Tata Institute of Fundamental Research, Bombay, 1980.
- [20] C. Siegel, *Berechnung von Zetafunktionen an ganzzahligen Stellen*, Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II **10** (1969), 87–102.
- [21] J. Waldspurger, *Sur les coefficients de Fourier des formes modulaires de poids demi-entier*, J. Math. Pures et Appl. **60** (1981), 375–484.
- [22] D. Zagier, *Modular forms whose Fourier coefficients involve zeta-functions of quadratic fields*, in Modular forms of One Variable VI, Lecture notes in Math. **627**, Springer, Berlin, 1977.
- [23] D. Zagier, *From quadratic functions to modular functions*, in Number Theory in Progress. **2** (1999), 1147–1178.

MATHEMATICAL INSTITUTE, UNIVERSITY OF COLOGNE, WEYERTAL 86-90, 50931 COLOGNE, GERMANY

E-mail address: `kbringma@math.uni-koeln.de`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAWAII, HONOLULU, HI 96822-2273

E-mail address: `pavel@math.hawaii.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HONG KONG, POKFULAM, HONG KONG

E-mail address: `bkane@maths.hku.hk`