



ORIGINAL ARTICLE

# Outcome and risk factor analysis of patients who underwent open infrarenal aortic aneurysm repair



Yuk Law, Yiu Che Chan\*, Grace C.Y. Cheung, Albert Chi Wai Ting, Stephen Wing Keung Cheng

Division of Vascular and Endovascular Surgery, Department of Surgery, University of Hong Kong Medical Center, Pokfulam, Hong Kong

Received 24 January 2015; received in revised form 26 February 2015; accepted 2 March 2015  
Available online 14 May 2015

## KEYWORDS

abdominal aortic aneurysm;  
abdominal aortic aneurysm repair;  
long term survival;  
reintervention

**Summary** *Introduction:* The aim of this study was to evaluate the short- and long-term outcomes in patients who underwent open infrarenal aortic aneurysm repair.

*Methods:* Consecutive patients who underwent open repair of infrarenal aortic aneurysms at our institution from July 1<sup>st</sup> 1990 to June 30<sup>th</sup> 2012 were reviewed from a prospective collected departmental database. Short-term outcomes included 30-day mortality and peri-operative complications. Independent risk factors to predict 30-day mortality were identified. Long-term survival and secondary interventions were also reported.

*Results:* Three hundred and eighty-three patients (317 males, median age 72 years with a range of 15–90 years) underwent open infrarenal aortic aneurysm repair during the period, of whom 266 (69.5%) were elective, 18 (4.7%) were urgent for symptomatic but nonruptured cases, and 99 (25.8%) were emergency procedures for ruptured aneurysms. Mean aneurysm size was 6.5 cm (ranging from 2.5 cm to 15 cm). All patients were followed up for at least 24 months with a mean follow up period 163 months. Overall 30-day mortality was 11.0% (36.4% for ruptured cases, 11.1% for symptomatic cases, and 1.5% for elective cases;  $p < 0.001$ ). Preexisting renal disease and ruptured aneurysms were independent risk factors for 30-day mortality ( $p = 0.001$  and  $p = 0.006$  respectively). Systemic complications included 50 cardiac events, 52 respiratory events, six renal events, three cerebral vascular accidents, and one deep vein thrombosis/pulmonary embolism. Local complications included two anastomotic/graft hemorrhage, 10 distal thrombosis/embolisms, five bowel ischemias, one spinal cord ischemia, and 17 wound complications. The ruptured group presented survival rates of 53.5%, 50.5%, 47.5%, 42.3%, 38.0%, 21.9%, and 12.5% at 1 year, 2 years, 3 years, 4 years, 5 years, 10 years, and 15 years, respectively; while nonruptured survival rates were 91.5%,

Conflicts of interest: No conflicts of interest declared.

\* Corresponding author. Division of Vascular and Endovascular Surgery, Department of Surgery, University of Hong Kong Medical Centre, South Wing, 14th Floor K Block, Queen Mary Hospital, Pokfulam Road, Hong Kong.

E-mail address: [ycchan88@hkucc.hku.hk](mailto:ycchan88@hkucc.hku.hk) (Y.C. Chan).

<http://dx.doi.org/10.1016/j.asjsur.2015.03.009>

1015-9584/Copyright © 2015, Asian Surgical Association. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

88.0%, 83.7%, 78.3%, 73.0%, 43.0%, and 25.3%, respectively (log rank  $p < 0.001$ ). For those who died 30 days after the operation, only six patients (1.8%) died from aneurysm related mortality. A total of three (0.9%) patients underwent late re-interventions, one for late aorto-enteric fistulae and two for anastomotic pseudoaneurysms.

**Conclusion:** In the current era of endovascular repair, open infrarenal aneurysm repair is effective and durable, and has very low secondary interventions rates.

Copyright © 2015, Asian Surgical Association. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Ever since Nicolai Volodos in 1986<sup>1</sup> and Juan Carlos Parodi in 1990<sup>2</sup> demonstrated the feasibility of aortic aneurysm exclusion with endografts, endovascular stent graft repair (EVAR) became the standard of care for infrarenal abdominal aortic aneurysms (AAA) at the end of past century. There was a resurgence of interest in open repair when long-term results from the EVAR1 trial,<sup>3</sup> DREAM trial,<sup>4</sup> OVER trial,<sup>5</sup> and ACE<sup>6</sup> trial were published recently. They all showed that the early mortality advantage of EVAR was lost at around 2 years of follow up, and re-intervention, long term complications, and cost were significantly higher in the EVAR group compared to the open repair group. This observation was reconfirmed using meta-analysis.<sup>7</sup>

We present the long-term outcomes of patients who underwent open infrarenal AAA repair in our department. This is the first paper reporting the long-term durability of open repairs in the Han Chinese population. Patients and clinicians can now be informed about the relative merits of open treatment versus endovascular treatment.

## 2. Methods

We reviewed all patients who received open repair of infrarenal aortic aneurysm at our institution, a tertiary referral center in Hong Kong for the period of July 1<sup>st</sup> 1990 to June 30<sup>th</sup> 2012. Data were subtracted from a prospective collected departmental database, supplemented by clinical notes and computer records. Only infrarenal aneurysms were included; while pararenal, suprarenal, and thoracoabdominal aneurysms were excluded.

Patients presented with ruptures, either suspected clinically or diagnosed using imaging. These patients were treated as surgical emergencies. Midline laparotomy and infrarenal clamps were the preferred approaches. Often a supraceliac clamp may be needed temporarily. Asymptomatic patients were offered repair when the aneurysm size reached 5.0 cm. Symptomatic patients included those with pain, infection, embolic phenomenon, and an aneurysm expansion rate  $>0.5$  cm/6 mo. They were treated in an early elective basis. Incision was rooftop or midline with an infrarenal clamp. Tube or bifurcated grafts of woven Dacron or knitted Dacron were used.

Patients were nursed in intensive care units after their operations. After discharge, patients were regularly followed up in the outpatient clinic. Surveillance duplex or computer tomography (CT) scans were not routine.

During the same period of time, our center also performed endovascular stenting but these patients were not reported in the current study.


Patients' baseline characteristics and follow-up periods were reported. Short-term outcomes included perioperative morbidities and mortalities. Variables including age, sex, comorbidities, preoperative hemoglobin level, preoperative base excess, whether the aneurysm had ruptured or not, and American Society of Anesthesiologists (ASA) grades were used to predict the 30-day mortality using a binary logistic regression model. Long-term survival and secondary interventions were reported. Chi-square test was used to differentiate significant differences between categorical variables, while Student  $t$  test was used for continuous variables. Statistical analysis was calculated by SPSS version 22 (SPSS Inc., Chicago, IL, USA). A  $p$  value  $<0.05$  was defined as significant.

Definitions of comorbidities were as follows. Cardiac history included stable angina, unstable angina, myocardial infarction, congestive heart failure, and arrhythmia. Pulmonary history included all patients with dyspnea or chest roentgenographic changes. Renal impairment was defined as serum creatinine level  $>120$   $\mu\text{mol/L}$ . Positive smoking history included all patients who had quit  $<10$  years.

Thirty-day mortality was defined as death  $\leq 30$  days after index open repair. Late aneurysm-related mortality was defined as death  $>30$  days after index operation and as a direct result of aneurysm rupture. Late re-intervention was defined as further intervention  $>30$  days after the index operation. Graft related events were defined as those occurring as a direct consequence of prosthetic aortic replacement and consisted of graft thrombosis, pseudoaneurysm formation, graft infection, and graft-enteric fistula. Other aneurysms remote from the index operation, e.g., thoracic or iliac aneurysm, requiring further intervention were not counted and were reported separately.

## 3. Results

Three hundred and eighty-three patients underwent open infrarenal aortic aneurysm repair during the study period (Fig. 1). Three hundred and seventeen (82.8%) were males while 66 (17.2%) were female. The median age was 72 years ranging from 15 years to 90 years, of whom 266 (69.5%) were elective, 18 (4.7%) were urgent for symptomatic but nonruptured cases, and 99 (25.8%) were emergency procedures for ruptured aneurysms. Mean aneurysm size was 6.5 cm ranging from 2.5 cm to 15.0 cm. Other baseline



**Fig. 1** Number of open infrarenal abdominal aortic aneurysm repairs over the years.

characteristics are shown in [Table 1](#). All patients were followed up for at least 24 months, with a mean follow up period 163 months.

### 3.1. Short-term results

Overall, 30-day mortality was 11.0% (36.4% for ruptured cases, 11.1% for symptomatic cases, and 1.5% for elective

**Table 1** Baseline characteristics of patients who underwent open surgical repair ( $n = 383$ ).

Variable	No.	%
Age	Median age 72 y (range, 15–90 y)	
≥70 y	230	60.1
<70 y	153	39.9
Gender		
Male	317	82.8
Female	66	17.2
Presentation		
Ruptured	99	25.8
Symptomatic	18	4.7
Elective	266	69.5
Coexisting condition		
Smoking	226	59.0
Hypertension	213	55.6
Diabetes mellitus	39	10.2
Cardiac disease	129	33.7
Cerebral vascular accident	44	11.5
Pulmonary disease	49	12.8
Renal impairment	89	23.2
Family history of aneurysm	3	0.8
Aneurysm size	Median size 6.5 cm (range, 2.5–15.0 cm)	
<5 cm	21	5.5
5–7 cm	251	65.5
>7 cm	111	29.0

cases;  $p < 0.001$ ). Univariate logistic regression analysis showed age, smoking history, hypertension, pre-existing renal disease, preoperative hemoglobin level, preoperative base excess, ruptured aneurysm, ASA grades 3–5, and aneurysm size were predictors of 30-day mortality ([Table 2](#)). With subsequent multivariate logistic regression, only pre-existing renal disease and ruptured aneurysms were found to be independent significant risk factors for 30-day mortality ( $p = 0.001$  and  $p = 0.006$ , respectively; [Table 3](#)).

Compared with nonruptured aneurysms, ruptured aneurysms led to significant longer aortic clamp time, prolonged operative duration, larger amount of blood loss, and increased postoperative ventilator assisted time ([Table 4](#)). Systemic complications included 50 cardiac events, 52 respiratory events, six renal events, three cerebral vascular accidents, and one deep vein thrombosis/pulmonary embolism. Local complications included two anastomotic/graft hemorrhage, 10 distal thrombosis/embolism, five bowel ischemias, one spinal cord ischemia, and 17 wound complications. Compared with nonruptured aneurysms, ruptured aneurysms led to significantly more adverse events in cardiac, respiratory, renal, and bowel ischemia complication subgroups ([Table 4](#)).

### 3.2. Long-term results

The ruptured group presented survival rates of 53.5%, 50.5%, 47.5%, 42.3%, 38.0%, 21.9%, and 12.5% at 1 year, 2 years, 3 years, 4 years, 5 years, 10 years, and 15 years, respectively; while the nonruptured group had survival rates of 91.5%, 88.0%, 83.7%, 78.3%, 73.0%, 43.0%, and 25.3% (log rank  $p < 0.001$ ; [Fig. 2](#)). For those who died 30 days after their operation, only six patients died from

**Table 2** Univariate analysis of preoperative variables to predict 30-day mortality.<sup>a</sup>

Variable	Odds ratio	95% Confidence interval	$p$
Male	1.046	0.443–2.469	0.918
Age	1.055	1.012–1.100	0.013
Smoking	0.269	0.135–0.537	<0.001
Hypertension	0.401	0.206–0.782	0.007
DM	0.411	0.095–1.770	0.233
Cardiac disease	1.390	0.721–2.680	0.325
CVA	0.792	0.269–2.337	0.673
Pulmonary disease	2.052	0.915–4.602	0.081
Renal disease	2.541	1.302–4.959	0.006
Preoperative hemoglobin level	0.669	0.578–0.773	<0.001
Preoperative base excess	0.883	0.832–0.936	<0.001
AAA size	1.314	1.097–1.573	0.003
Ruptured	26.476	10.695–65.546	<0.001
ASA 3–5	5.807	1.747–19.303	0.004

AAA = abdominal aortic aneurysm; ASA = American Society of Anaesthesiologist; CVA = cerebrovascular accident; DM = diabetes mellitus.

<sup>a</sup> By binary logistic regression. Age, preoperative hemoglobin level, preoperative base excess, and AAA size are continuous variables, while others are categorical variables.


**Table 3** Multivariate analysis of preoperative variables to predict 30-day mortality.<sup>a</sup>

Variable	Odd ratio	95% Confidence interval	<i>p</i>
Age	1.037	0.960 1.119	0.358
Smoking	1.005	0.322 3.138	0.994
Hypertension	0.385	0.106 1.389	0.145
Renal disease	8.434	2.349 30.284	0.001
Preoperative hemoglobin level	0.867	0.668 1.124	0.280
Preoperative base excess	0.952	0.889 1.019	0.156
AAA size	1.193	0.909 1.566	0.203
Ruptured	7.803	1.778 34.233	0.006
ASA 3–5	2.266	0.223 23.032	0.489

AAA = abdominal aortic aneurysm; ASA = American Society of Anesthesiologists score.

<sup>a</sup> By binary logistic regression. Age, preoperative hemoglobin level, preoperative base excess, and AAA size are continuous variables, while others are categorical variables.

aneurysm related causes, giving a late aneurysm-related mortality of 1.8% (Table 5). There were a total of three (0.9%) patients who underwent late re-interventions due to graft related events. The first patient underwent open repair in 1994 at the age of 66 years. The 4.4 cm infrarenal aortic aneurysm and 5.7 cm left iliac aneurysm were replaced with aorto-bi-iliac knitted Dacron graft. He presented with recurrent gastrointestinal bleeding of unknown origin 6 years later. Upper endoscopy and colonoscopy failed to localize the bleeding source. Laparotomy and enteroscopy showed suspicious bleeding from the proximal jejunum and possible graft-to-jejunum fistula. Bleeding

**Fig. 2** Survival curve of ruptured versus nonruptured (log rank *p* < 0.001).

was settled with aorto-uni-iliac endovascular stenting and cross femoral bypass. The patient is still alive to date (Fig. 3). The second patient underwent open repair with aorto-bi-femoral woven Dacron graft in 1993 at the age of 72 years. Seven years later, he was incidentally found to have a pulsatile mass at the epigastrium with a CT scan showing an infrarenal 8 cm proximal anastomosis pseudoaneurysm. Treatment was standard endovascular repair (Fig. 4). The last patient had open repair surgery with aorto-bi-iliac woven Dacron graft in 1993 at the age of 76 years. He complained of epigastric pain 20 years later with a CT scan showing a 9 cm proximal perivisceral pseudoaneurysm, which was settled with endovascular stenting

**Table 4** Short-term outcomes.

	Overall ( <i>n</i> = 383)	Ruptured ( <i>n</i> = 99)	Nonruptured ( <i>n</i> = 284)	<i>p</i>
Mean aortic clamp time (min)	62.4	72.8	59.5	<0.001 <sup>a</sup>
Mean operative time (min)	202.5	180.3	209.7	0.001 <sup>a</sup>
Blood loss (mL)	1748	3858	1055	<0.001 <sup>a</sup>
Mean days of assisted ventilation (d)	2.2	7.6	0.5	<0.001 <sup>a</sup>
30 d mortality	42 (11.0)	36 (36.4)	6 (2.1)	<0.001 <sup>b</sup>
Systemic complications				
Cardiac	50 (13.0)	21 (21.2)	29 (10.2)	0.009 <sup>b</sup>
Respiratory	52 (13.6)	32 (32.3)	20 (7.0)	<0.001 <sup>b</sup>
Renal	6 (1.6)	5 (5.1)	1 (0.4)	0.005 <sup>b</sup>
CVA	3 (0.8)	1 (1.0)	2 (0.7)	1.000 <sup>b</sup>
DVT/PE	1 (0.3)	1 (1.0)	0 (0)	0.258 <sup>b</sup>
Local complications				
Graft complication	2 (0.5)	0 (0)	2 (0.7)	1.000 <sup>b</sup>
Distal thrombosis/embolism	10 (2.6)	5 (5.1)	5 (1.8)	0.134 <sup>b</sup>
Bowel ischemia	5 (1.3)	4 (4.0)	1 (0.4)	0.017 <sup>b</sup>
Spinal ischemia	1 (0.3)	1 (1.0)	0 (0)	0.258 <sup>b</sup>
Wound complication	17 (4.4)	7 (7.1)	10 (3.5)	0.158 <sup>b</sup>

Data are presented as *n* or *n* (%).

CVA = cerebrovascular accident; DVT = deep vein thrombosis; PE = pulmonary embolism.

<sup>a</sup> Student *t* test.

<sup>b</sup> Chi-square test.

**Table 5** Causes of death after 30 days.

Causes	No. ( <i>n</i> = 341)	
Cardiovascular related	41	
Noncardiovascular related	108	
Ruptured aneurysm/complications	6	1 type A dissection with hemopericardium 1 ruptured arch aneurysm 3 ruptured thoracic aneurysm 1 suspected perigraft infection
Unknown	71	
Still alive	115	

with fenestration to all four visceral vessels (Fig. 5; Table 6). In addition, 17 (5.0%) patients developed thoracic or iliac aneurysms in later life. They were treated with open or endovascular repair (Table 7).

#### 4. Discussion

Our indications for surgical management of infrarenal abdominal aortic aneurysm were similar to well established standards.<sup>8–13</sup> One-fourth of open repairs were for ruptured cases with 36.4% 30-day mortality. The mortality rate for open elective infrarenal aortic aneurysm repair was 1.5%. Symptomatic aneurysms only account for a small proportion.

World reported perioperative mortality and morbidity varied greatly. A meta-analysis of the reported literature showed 30-day or in-hospital mortality from ruptured aneurysm ranging from 27% to 69%.<sup>14</sup> As for elective cases, 30-day or in-hospital mortality ranged from 1.4% to 6.5%. Overall 30-day mortality of the open arm of EVAR1,<sup>3</sup>

DREAM,<sup>4</sup> OVER,<sup>5</sup> and ACE<sup>6</sup> trials was 3.69% (range, 0.6–4.6%). While certain high volume centers reported excellent mortality rates of 0% to 1.2%,<sup>15–17</sup> multi-institutional nationwide data continued to show higher incidences of mortality ranging from 4.2% to 8.4%.<sup>16,17</sup> There were strong relationships between operative outcomes and the case volume of the surgeon or hospital. Birkmeyer et al<sup>18,19</sup> reported an elective AAA repair mortality rate of 7.8% in low volume centers (<17 cases/y), as opposed to 4.9% in high volume centers (>79 cases/y).

Despite advancements in operative, anesthetic, and intensive care, perioperative mortality of ruptured aneurysms was still extremely high. Several meta-analyses are available in literature. Hoornweg et al<sup>20</sup> reported an overall mortality of 48.5%. Bown et al<sup>21</sup> reported the rate as 48%. Kantonen et al<sup>22</sup> reported the rate as 68%. This high mortality rate is most likely related to factors such as delays in recognition and intervention, hemodynamic instability, profound blood loss, and suboptimal perioperative care in emergency setting.<sup>21,22</sup> The relatively low rates in our institution were probably related to short travel times to




Fig. 3 Late intervention, Patient 1.






Fig. 4 Late intervention Patient 2.

the hospital, availability of dedicated vascular surgeons, and outstanding anesthetic and intensive care.

Several factors have been identified as predictive factors for operative mortality. These included advanced age, women, elevated body mass index (BMI), ruptured aneurysm, large aneurysm size, smoking, renal impairment, chronic obstructive pulmonary disease, coronary artery disease, prolonged aortic clamp time, greater blood loss, and positive fluid balance.<sup>23–34</sup> Furthermore, a few risk scoring systems have been developed to identify high risk patients undergoing open AAA repair. The Glasgow Aneurysm score (GAS),<sup>35</sup> Vascular Physiology only-Physiological and Operative Severity Score for enUmeration of Mortality

Table 6 Late re-interventions.

Causes	No. (n = 341)	Treatments
Aorto-enteric fistula	1	EVAR
Proximal anastomotic pseudoaneurysm	2	EVAR

Table 7 Treatment of aneurysm elsewhere.

Types of aneurysm	No. (n = 341)	Treatments
Thoracoabdominal aneurysm	10	TEVAR
	1	Open
Iliac aneurysm	5	EVAR
	1	Open

(T)EVAR = (Thoracic) endovascular aortic repair.

[V(p)-POSSUM],<sup>36</sup> Vascular Biochemical and Hematological Outcome Model (VBHOM),<sup>37</sup> Lee’s Revised Cardiac Risk Index (RCRI),<sup>38</sup> and Preoperative Risk Score of the Estimation of Physiological Ability and Surgical Stress Score (PRS of E-PASS)<sup>39</sup> were the best known predictors. Bryce et al<sup>40</sup> and Tang et al<sup>41</sup> showed that V(p)-POSSUM and E-PASS better predict outcome than the other two. However, due to the size of our patient cohort, our study only showed that pre-existing renal disease and ruptured aneurysms were associated with increased mortality in elective surgery and in emergency open repair. The most commonly reported risk factors of early mortality, such as advanced age and female gender, were insignificant in this study.

The merits of open repair compared with endovascular repair were lower late aneurysm-related mortality and lower late re-intervention rates. Both EVAR<sup>3</sup> and DREAM<sup>4</sup> trials reported their long-term survival after a mean




Fig. 5 Late intervention, Patient 3.

follow up of 6 years and 6.4 years respectively. Late aneurysm related mortality after open repair was 2.2% and 0.6% in EVAR1 and DREAM respectively. Late re-intervention rate was 1.7% per year in the EVAR1 trial, and 18.1% in the DREAM trial. Both studies demonstrated significant lower re-intervention rates after open repair compared with those who underwent endovascular repair (5.1%/y and 29.6%/y in the EVAR1 trial and DREAM trial respectively). The Mayo clinic reported re-intervention rates for late aortic graft-related events of 5.1–5.3% at a median follow up of 5.8–7 years, with an associated operative mortality rate of 17–19%.<sup>42,43</sup> Biancari et al<sup>44</sup> from Finland reported a re-intervention rate of 10.6% at the median follow up of 8 years, with an associated operative mortality rate of 9%. Adam et al<sup>45</sup> from Australia reported a re-intervention rate of 3.4% at the median follow up of 41 months in non-ruptured aneurysms and a rate of 5% at the median follow up of 30 months in ruptured cases. Our data of 1.8% late aneurysm-related mortality and 0.9% late re-intervention rate demonstrated that open AAA repair has excellent long-term durability in our population, and furthermore, the results compare favorably with previous reports from North America, Europe, and Australia (Table 8).

Graft occlusion and anastomotic pseudoaneurysm were the more frequent graft-related indications for re-intervention, while graft infection and graft-enteric fistula occurred infrequently.<sup>42–47</sup> All re-interventions in our study were for symptomatic complications, while it may be difficult to identify asymptomatic complications as imaging follow ups after open repair were not routine in our center. CT scans were included in the follow up protocol for patient randomized to open repair in EVAR1 trial,<sup>48</sup> and it is anticipated that further reports from the trial will determine the true incidence of late graft-related events.

There are several limitations in this study, as this study had a relatively small case number and incomplete long-term data collection. Completeness of patient follow up was an inherited limitation in our retrospective study. Being a tertiary referral center, every patient with symptomatic late graft related events would be referred back to us for management. The late re-intervention rate should therefore be accurate. However, the causes of death of a vast number of patients were unknown despite every effort to determine the cause of death, adding uncertainty to the

late aneurysm-related mortality. Another important limitation of this study was the lack of comparison with the endovascular group. Our center began endovascular treatment for AAA in 1999. We were eagerly waiting the long-term outcomes of endovascular AAA repair. In the current era of endovascular repair, open infrarenal aneurysm repair is safe and effective, and still has its place in cases with difficult necks and access. As our study spanned from 1990 to 2012, there is always an issue with different emerging techniques over time, but nonetheless it represents one of the largest databases of Chinese patients receiving open AAA management.

## Acknowledgments

This study received no financial support.

## References

- Volodos NL, Karpovich IP, Troyan VI, et al. Clinical experience of the use of self-fixing synthetic prostheses for remote endoprosthetics of the thoracic and the abdominal aorta and iliac arteries through the femoral artery and as intraoperative endoprosthesis for aorta reconstruction. *Vasa Suppl.* 1991;33: 93–95.
- Parodi JC, Palmaz JC, Barone HD. Transfemoral intraluminal graft implantation for abdominal aortic aneurysms. *Ann Vasc Surg.* 1991;5:491–499.
- The United Kingdom EVAR Trial Investigators. Endovascular versus open repair of abdominal aortic aneurysm. *N Engl J Med.* 2010;362:1863–1871.
- De Bruin JL, Baas AF, Buth J, et al. Long-term outcome of open or endovascular repair of abdominal aortic aneurysm. *N Engl J Med.* 2010;362:1881–1889.
- Lederle FA, Frieschlag JA, Kyriakides TC, et al. Outcomes following endovascular vs open repair of abdominal aortic aneurysm: a randomized trial. *JAMA.* 2009;302:1535–1542.
- Becquemain JP. The ACE trial: a randomized comparison of open versus endovascular repair in good risk patients with abdominal aortic aneurysm. *J Vasc Surg.* 2009;50:222–2224. discussion 224.
- Dangas G, O'Connor D, Firwana B, et al. Open versus endovascular stent graft repair of abdominal aortic aneurysms: a meta-analysis of randomized trials. *JACC Cardiovasc Interv.* 2012;5:1071–1080.
- Mortality results for randomised controlled trial of early elective surgery or ultrasonographic surveillance for small abdominal aortic aneurysms. The UK Small Aneurysm Trial Participants. *Lancet.* 1998;352:1649–1655.
- Lederle FA, Wilson SE, Johnson JR, et al. Immediate repair compared with surveillance of small abdominal aortic aneurysms. *N Engl J Med.* 2002;346:1437–1444.
- Cao P, De Rango P, Verzini F, et al. Comparison of surveillance versus aortic endografting for small aneurysm repair (CAESAR): results from a randomised trial. *Eur J Vasc Endovasc Surg.* 2011;41:13–25.
- Ouriel K. The PIVOTAL study: a randomized comparison of endovascular repair versus surveillance in patients with smaller abdominal aortic aneurysms. *J Vasc Surg.* 2009; 49:266–2669.
- Eliason JL, Clouse WD. Current management of infrarenal abdominal aortic aneurysms. *Surg Clin North Am.* 2007;87: 1017–1033.
- Chaikof EL, Brewszter DC, Dalman RL, et al. The care of patients with an abdominal aortic aneurysm: the Society for

**Table 8** Late aneurysm related mortality and late re-intervention rate in other studies.

	Late aneurysm related mortality (%)	Late re-intervention rate (%)	Mean follow up period
EVAR 1	2.2	1.7%/y	6 y
DREAM	0.6	18.1	6.4 y
Mayo clinic		5.1–5.3	5.8–7 y
Biancari et al <sup>44</sup> (Finland)		10.6	8 y
Adam et al <sup>45</sup> (Australia)		3.4 (nonruptured) 5 (ruptured)	41 mo 30 mo
Our data	1.8	0.9	13.6 y

- Vascular Surgery practice guidelines. *J Vasc Surg.* 2009;50: S2–49.
14. Hallin A, Bergqvist D, Holmberg L. Literature review of surgical management of abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg.* 2001;22:197–204.
  15. Hertzner NR, Mascha EJ, Karafa MT, O'Hara PJ, Krajewski LP, Beven EG. Open infrarenal abdominal aortic aneurysm repair: the Cleveland Clinic experience from 1989 to 1998. *J Vasc Surg.* 2002;35:1145–1154.
  16. Lawrence PF, Gazak C, Bhirangi L, et al. The epidemiology of surgically repaired aneurysms in the United States. *J Vasc Surg.* 1999;30:632–640.
  17. Dimick JB, Cowan Jr JA, Stanley JC, Henke PK, Pronovost PJ, Upchurch Jr GR. Surgeon specialty and provider volumes are related to outcome of intact abdominal aortic aneurysm repair in the United States. *J Vasc Surg.* 2003;38:739–744.
  18. Birkmeyer JD, Siewers AE, Finlayson EVA, et al. Hospital volume and surgical mortality in the United States. *N Engl J Med.* 2002;346:1128–1137.
  19. Birkmeyer JD, Stukel TA, Siewers AE, et al. Surgeon volume and operative mortality in the United States. *N Engl J Med.* 2003;349:2117–2127.
  20. Hoornweg LL, Storm-versloot MN, Ubbink DT, Koelemay MJ, Legemate DA, Balm R. Meta analysis on mortality of ruptured abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg.* 2008;35:558–570.
  21. Bown MJ, Sutton AJ, Bell PR, Sayers RD. A meta-analysis of 50 years of ruptured abdominal aortic aneurysm repair. *Br J Surg.* 2002;89:714–730.
  22. Kantonen I, Lepäntalo M, Brommels M, Luther M, Salenius JP, Ylönen K. Mortality in ruptured abdominal aortic aneurysms. The Finnvasc Study Group. *Eur J Vasc Endovasc Surg.* 1999;17: 208–212.
  23. Bauer EP, Redaelli C, von Segesser LK, Turina MI. Ruptured abdominal aortic aneurysms: predictors for early complications and death. *Surgery.* 1993;114:31–315.
  24. Sasaki S, Takigami K, Kuniyama T, et al. Abdominal aortic aneurysms in aged patients: analysis of risk factors in non-ruptured cases. *J Cardiovasc Surg (Torino).* 1999;40:1–5.
  25. Berry AJ, Smith RB, Weintraub WS, et al. Age versus comorbidities as risk factors for complications after elective abdominal aortic reconstructive surgery. *J Vasc Surg.* 2001;33: 345–352.
  26. Roger VL, Ballard DJ, Hallett Jr JW, Osmundson PJ, Puetz PA, Gersh BJ. Influence of coronary artery disease on morbidity and mortality after abdominal aortic aneurysmectomy: a population-based study, 1971–1987. *J Am Coll Cardiol.* 1989; 14:1245–1252.
  27. Stone DH, Goodney PP, Kalish J, et al. Severity of chronic obstructive pulmonary disease is associated with adverse outcomes in patients undergoing elective abdominal aortic aneurysm repair. *J Vasc Surg.* 2013;57:1531–1536.
  28. Lo RC, Bensley RP, Hamdan AD, et al. Gender differences in abdominal aortic aneurysm presentation, repair, and mortality in the Vascular Study Group of New England. *J Vasc Surg.* 2013; 57:1261–1268. e1–5.
  29. Kennedy NA, Flynn LM, Berg RM, Lorelli DR, Rama K, Rizk Y. The evaluation of morbidity and mortality in abdominal aortic aneurysm repair patients as related to body mass index. *Am J Surg.* 2010;199:369–371. discussion 371.
  30. Grant SW, Hickey GL, Grayson AD, Mitchell DC, McCollum CN. National risk prediction model for elective abdominal aortic aneurysm repair. *Br J Surg.* 2013;100:645–653.
  31. Hertzner NR, Mascha EJ. A personal experience with factors influencing survival after elective open repair of infrarenal aortic aneurysms. *J Vasc Surg.* 2005;42:898–905.
  32. McArdle GT, Price G, Lewis A, et al. Positive fluid balance is associated with complications after elective open infrarenal abdominal aortic aneurysm repair. *Eur J Vasc Endovasc Surg.* 2007;34:522–527.
  33. Beck AW, Goodney PP, Nolan BW, et al. Predicting 1-year mortality after elective abdominal aortic aneurysm repair. *J Vasc Surg.* 2009;49:838–843. discussion 843–844.
  34. Nathan DP, Brinster CJ, Jackson BM, et al. Predictors of decreased short- and long-term survival following open abdominal aortic aneurysm repair. *J Vasc Surg.* 2011;54: 1237–1243.
  35. Samy AK, Murray G, MacBain G. Glasgow aneurysm score. *Cardiovasc Surg.* 1994;2:41–44.
  36. Prytherch DR, Sutton GL, Boyle JR. Portsmouth POSSUM models for abdominal aortic aneurysm surgery. *Br J Surg.* 2001;88: 958–963.
  37. Tang T, Walsh SR, Prytherch DR, et al. VBHOM, a data economic model for predicting the outcome after open abdominal aortic aneurysm surgery. *Br J Surg.* 2007;94:717–721.
  38. Lee TH, Marcantonio ER, Mangione CM, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation.* 1999;100: 1043–1049.
  39. Tang T, Walsh SR, Fanshawe TR, et al. Estimation of physiologic ability and surgical stress (E-PASS) as a predictor of immediate outcome after elective abdominal aortic aneurysm surgery. *Am J Surg.* 2007;194:176–182.
  40. Bryce GJ, Payne CJ, Gibson SC, Kingsmore DB, Byrne DS, Delles C. Risk stratification scores in elective open abdominal aortic aneurysm repair: are they suitable for preoperative decision making? *Eur J Vasc Endovasc Surg.* 2012;44:55–61.
  41. Tang TY, Walsh SR, Fanshawe TR, et al. Comparison of risk-scoring methods in predicting the immediate outcome after elective open abdominal aortic aneurysm surgery. *Eur J Vasc Endovasc Surg.* 2007;34:505–513.
  42. Hallett Jr JW, Marshall DM, Petterson TM, et al. Graft-related complications after abdominal aortic aneurysm repair: reassurance from a 36-year population-based experience. *J Vasc Surg.* 1997;25:277–284. discussion 285–286.
  43. Cho JS, Głowiczki P, Martelli E, et al. Long-term survival and late complications after repair of ruptured abdominal aortic aneurysms. *J Vasc Surg.* 1998;27:813–819. discussion 819–820.
  44. Biancari F, Ylönen K, Anttila V, et al. Durability of open repair of infrarenal abdominal aortic aneurysm: a 15-year follow-up study. *J Vasc Surg.* 2002;35:87–93.
  45. Adam DJ, Fitridge RA, Raptis S. Late reintervention for aortic graft-related events and new aortoiliac disease after open abdominal aortic aneurysm repair in an Australian population. *J Vasc Surg.* 2006;43:701–705. discussion 705–706.
  46. Crawford ES, Saleh SA, Babb 3rd JW, Glaeser DH, Vaccaro PS, Silvers A. Infrarenal abdominal aortic aneurysm: factors influencing survival after operation performed over a 25-year period. *Ann Surg.* 1981;193:699–709.
  47. Plate G, Hollier LA, O'Brien P, Pairolero PC, Cherry KJ, Kazmier FJ. Recurrent aneurysms and late vascular complications following repair of abdominal aortic aneurysms. *Arch Surg.* 1985;120:590–594.
  48. Endovascular aneurysm repair versus open repair in patients with abdominal aortic aneurysm (EVAR trial 1): randomised controlled trial. *Lancet.* 2005;365:2179–2186.