

Research Article

Calycosin and Formononetin Induce Endothelium-Dependent Vasodilation by the Activation of Large-Conductance Ca^{2+} -Activated K^+ Channels (BK_{Ca})

Hisa Hui Ling Tseng,¹ Chi Teng Vong,¹ George Pak-Heng Leung,² Sai Wang Seto,³ Yiu Wa Kwan,⁴ Simon Ming-Yuen Lee,¹ and Maggie Pui Man Hoi¹

¹State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Avenida da Universidade, Taipa, Macau

²Department of Pharmacology and Pharmacy, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong

³National Institute of Complementary Medicine, Western Sydney University, Penrith, NSW, Australia

⁴School of Biomedical Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong

Correspondence should be addressed to Maggie Pui Man Hoi; maghoi@umac.mo

Received 5 July 2016; Revised 26 September 2016; Accepted 19 October 2016

Academic Editor: Kuzhuvelil B. Harikumar

Copyright © 2016 Hisa Hui Ling Tseng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Calycosin and formononetin are two structurally similar isoflavonoids that have been shown to induce vasodilation in aorta and conduit arteries, but study of their actions on endothelial functions is lacking. Here, we demonstrated that both isoflavonoids relaxed rat mesenteric resistance arteries in a concentration-dependent manner, which was reduced by endothelial disruption and nitric oxide synthase (NOS) inhibition, indicating the involvement of both endothelium and vascular smooth muscle. In addition, the endothelium-dependent vasodilation, but not the endothelium-independent vasodilation, was blocked by BK_{Ca} inhibitor iberiotoxin (IbTX). Using human umbilical vein endothelial cells (HUVECs) as a model, we showed calycosin and formononetin induced dose-dependent outwardly rectifying K^+ currents using whole cell patch clamp. These currents were blocked by tetraethylammonium chloride (TEACl), charybdotoxin (ChTX), or IbTX, but not apamin. We further demonstrated that both isoflavonoids significantly increased nitric oxide (NO) production and upregulated the activities and expressions of endothelial NOS (eNOS) and neuronal NOS (nNOS). These results suggested that calycosin and formononetin act as endothelial BK_{Ca} activators for mediating endothelium-dependent vasodilation through enhancing endothelium hyperpolarization and NO production. Since activation of BK_{Ca} plays a role in improving behavioral and cognitive disorders, we suggested that these two isoflavonoids could provide beneficial effects to cognitive disorders through vascular regulation.

1. Introduction

Calycosin and formononetin (Figure 1) are two structurally similar isoflavonoids that are present abundantly in traditional Chinese medicine (TCM) such as Radix Astragali (Huang Qi) and phytoestrogenic herb including *Trifolium pretense* L. (red clover), and they have a long clinical history in treating various cardiovascular diseases [1, 2]. Previous studies have shown that calycosin and formononetin produced antihypertensive effects and improved endothelial and cardiovascular functions [3–6]. They have been shown

to display vasoactive effects in various vascular beds [3, 5–7]. In rat aorta, calycosin induced vasodilation mainly by inhibiting voltage-dependent Ca^{2+} channel (VDCC) in vascular smooth muscle [7], while formononetin caused vasodilation by releasing nitric oxide (NO) in endothelial cells, as well as by the activation of large-conductance Ca^{2+} -activated K^+ (BK_{Ca}) and ATP-sensitive potassium (K_{ATP}) channels in aortic smooth muscle cells [6]. In addition, these two isoflavonoids were reported to ameliorate cerebral ischemia and reperfusion injury by improving endothelial dysfunction [8, 9]. These observation led us to investigate the




FIGURE 1: Chemical structures of calycosin and formononetin.

pharmacological underlying mechanisms of calycosin and formononetin in small resistance arteries (internal diameter $\leq 300 \mu\text{m}$) and vascular endothelial cells.

Small resistance arteries are major sites of peripheral vascular resistance and are closely related to endothelial dysfunction and the pathogenesis of hypertension [10, 11]. Interestingly, it has been shown that endothelium-dependent hyperpolarization (EDH) was more pronounced in small resistance arteries than large conduit arteries such as aorta [10, 12]. In the vascular walls, calcium-activated potassium (K_{Ca}) channel is the main contributor for endothelium-derived hyperpolarizing factor- (EDHF-) mediated responses, which plays a crucial role in the regulation of vascular tone and the maintenance of systemic blood pressure [13, 14]. Recently, endothelial K_{Ca} channels have been used as new drug targets for cardiovascular diseases such as hypertension to stimulate EDHF and NO production to improve endothelial dysfunction [13, 15–17]. There are three types of K_{Ca} channels based on their conductances, including IK_{Ca} (intermediate conductance), SK_{Ca} (small conductance), and BK_{Ca} (large conductance) [18]. Although IK_{Ca} and SK_{Ca} are the major K_{Ca} channels present in the endothelial cells of arteries, BK_{Ca} channels have been identified in the endothelium of rat pulmonary and mesenteric arteries [19, 20] and cultured endothelial cells [21, 22]. It has been suggested that BK_{Ca} channel has a compensatory role for improving vasoreactivity in environment such as hypertension and cardiovascular diseases [23, 24]. In addition, it was shown that BK_{Ca} channel acts as a negative feedback mechanism for vascular dysfunction impaired by reactive oxygen species (ROS) and is overexpressed in diseases associated with endothelial dysfunction [25–27].

In the present study, we investigated the effects of calycosin and formononetin in rat mesenteric resistance arteries and their underlying mechanisms with a focus on endothelial K^+ channel. We demonstrated that calycosin and formononetin induced endothelium-dependent vasodilation through NO production and BK_{Ca} channel activation. We also showed that calycosin and formononetin increased NO production through endothelial nitric oxide synthase (eNOS) and neuronal nitric oxide synthase (nNOS) pathway and activated endothelial BK_{Ca} channels in human umbilical endothelial cells (HUVEC). Taken together, our study demonstrated that calycosin and formononetin are endothelial BK_{Ca} channel activators, suggesting a novel mechanism for vasodilation by these isoflavones, and they might be

potential for treating vascular and cerebrovascular diseases associated with endothelial dysfunction.

2. Material and Methods

2.1. Chemicals and Reagents. Calycosin and formononetin were purchased from Shanghai Forever Biotech (Shanghai, China), and they were dissolved in dimethyl sulphoxide (DMSO). All the chemicals were purchased from Sigma-Aldrich (St. Louis, MO, USA), while the cell culture reagents were from Gibco (Carlsbad, CA, USA). All the antibodies used for immunoblotting were purchased from Cell Signaling Technology (Danvers, MA, USA).

2.2. Animals. All the procedures were carried out according to the ethical guidelines of the Institute of Chinese Medical Sciences (ICMS), University of Macau, and NIH guide for the Care and Use of Laboratory Animals. Male Sprague-Dawley rats ($250 \pm 350 \text{ g}$) were killed by cervical dislocation.

2.3. Preparation of Resistance Mesenteric Arteries. The mesenteric arterial beds were dissected from rats immediately after cervical dislocation and were kept in cold Krebs–Henseleit buffer with the following compositions (mM): NaCl, 118; KCl, 4.7; MgSO_4 , 1.2; KH_2PO_4 1.2; NaHCO_3 , 25; CaCl_2 , 2.5; D-glucose, 5.5. The Krebs–Henseleit buffer was continuously gassed with a mixture of 95% O_2 /5% CO_2 to maintain a pH of 7.4. Small (the third branches of the mesenteric artery) mesenteric arteries were placed in a 4-chamber wire myograph (DMT, Tissue Bath System 700MO, Aarhus, Denmark) and were maintained at 37°C in Krebs–Henseleit buffer. The normalized protocol was used as described previously [28]. The tension was recorded by the PowerLab recording system (ADInstruments, Hastings, UK). In endothelium-denuded experiments, the endothelium was removed by rubbing the inner surface of the segment with human hair. In endothelium-intact experiments, the presence of a functional endothelium was examined by precontracting the arteries with methoxamine ($10 \mu\text{M}$) and then was relaxed by carbachol ($10 \mu\text{M}$), and this vasodilation was greater than 80%.

2.4. Myograph Experimental Protocol. After 30 min equilibration, arteries were firstly precontracted by methoxamine

(10 μM). Once the tension was stable, calycosin (1 nM–100 μM) and formononetin (1 nM–100 μM) were added cumulatively to produce a concentration-dependent response in endothelium-intact and endothelium-denuded mesenteric arteries. The effects of vasodilation in response to calycosin and formononetin were also examined with preincubation of indomethacin, N_ω -nitro-L-arginine methyl ester (L-NAME), tetraethylammonium chloride (TEACl), apamin, charybdotoxin (ChTX), iberiotoxin (IbTX), and glibenclamide for 30 min, before precontracting with methoxamine. In some experiments, the arteries were precontracted with high K^+ (60 mM KCl) Krebs–Henseleit buffer by replacing NaCl with KCl in the standard Krebs–Henseleit solution, as described previously [28].

2.5. Cell Culture. Human umbilical vein endothelial cells (HUVECs) were purchased from Life Technologies (Carlsbad, CA, USA) and were cultured in Hams F-12k nutrient medium supplemented with 15% fetal bovine serum, 1% penicillin-streptomycin, 100 mg/mL heparin, and 10 mg/mL endothelial cell growth supplement (Sigma-Aldrich, St. Louis, MO, USA) at 37°C under an atmosphere of 5% CO_2 in air. The cells were used in passages 2–6. Plated cells were allowed to adhere for 2 h before patch clamp experiments. For the experiments measuring NO production, and the expressions of different nitric oxide synthases (NOS), calycosin and formononetin were treated for 1 h in HUVEC.

2.6. NO Production Assay. NO was measured by a Nitric Oxide Assay Kit (Abcam, Cambridge, UK) following the manufacturer's instruction. Briefly, 75 μL of cell supernatants was mixed with 5 μL enzyme cofactor solution and 5 μL nitrate reductase. Following 2 h of incubation for converting nitrates to nitrites, 5 μL enhancer was added to each sample and incubated for 30 min. 5 μL DAN Probe was then added and incubated for 10 min. After that, 5 μL NaOH was added in the mixture for 10 min. The fluorescence intensity was detected by a microplate reader (SpectraMax M5, Molecular Devices, USA) using excitation at 360 nm and emission at 450 nm wavelengths.

2.7. Patch Clamp Experiments. All the cells were superfused with the extracellular solution with the following compositions (mM): NaCl, 140; KCl, 5.4; CaCl_2 , 1.8; MgCl_2 , 1; NaH_2PO_4 0.33; HEPES, 5; D-glucose, 5.5; pH 7.4. The recordings were made by using an Axopatch-200B amplifier, Digidata-1321 interface, and pClamp10.0 software (Axon Instruments, Foster City, CA). In the outward K^+ channel experiment, the pipettes (resistance 2–3 $\text{M}\Omega$) were filled with intracellular pipette solution (mM): KCl, 140; MgCl_2 , 2.5; D-glucose, 10; HEPES, 5; pH 7.2. In some experiments, $[\text{Ca}^{2+}]_i$ was adjusted to 250, 500, or 1000 nM by administering 6.7, 8.0, or 8.9 mM CaCl_2 to the intracellular pipette solution, respectively, and in the presence of 10 mM EGTA. The cells were clamped at a holding potential of -60 mV and various potential steps from -100 mV to $+100$ mV with 10 mV increments, and the currents were stimulated by a series of 250 ms.

2.8. $[\text{Ca}^{2+}]_i$ Measurements. The intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) was measured in single cells as previously described [29]. Cells were loaded with Fluo-4 AM (2 μM , Molecular Probes, US) in Tyrode solution containing 136.5 mM NaCl, 5.4 mM KCl, 0.53 mM MgCl_2 , 1.8 mM CaCl_2 , 0.33 mM NaH_2PO_4 , 5.5 mM glucose, and 5.5 mM HEPES (pH 7.4, adjusted with NaOH) for 30 min at 37°C. Fluo-4 fluorescence intensity (494 nm excitation; 506 nm emission) was sampled at 5 s intervals using a Cell^R system (MT20, Olympus, US).

2.9. Western Blot Analysis. After indicated treatment, the protein was extracted with ice-cold lysis buffer, and the concentrations of lysates were measured by the bicinchoninic acid kit (Pierce, US). 30 μg proteins were used and separated by 10% SDS-PAGE gels and were transferred onto the nitrocellulose membranes. Membranes were incubated with primary antibodies (eNOS, p-eNOS, nNOS, and iNOS antibody using 1/1000 dilution, whereas GAPDH antibody using 1/4000 dilution) overnight at 4°C, and secondary antibodies (anti-rabbit with 1/1000 dilution) for 1 hr, and blots were developed by enhanced chemiluminescence (GE Healthcare Life Sciences, UK) with an imaging system (Bio-Rad Laboratories, USA). GAPDH were used as housekeeping controls.

2.10. Data and Statistical Analysis. Data were expressed as mean \pm SEM. Vasodilation responses in each segment were expressed as a percentage of relaxation. The dose-response curves were fitted to a logistic equation described previously [30]. The maximum percentage of relaxation (R_{max}) and the concentration required to produce 50% of maximal response tone (EC_{50}) were calculated from the fitted curves. Currents curves were fitted by Boltzmann equations. The curve fitting and statistical analyzes were determined using GraphPad Prism 5 (San Diego, CA, USA). Significant differences were analyzed by *t*-test, one-way ANOVA followed by a Dunnett's test, or two-way ANOVA followed by a Bonferroni *post hoc* test. $P < 0.05$ was considered as significant.

3. Results

3.1. Effects of Endothelial Removal, L-NAME, and K^+ Channel Inhibitors on Vasodilation in Response to Calycosin. Calycosin induced dose-dependent vasodilation with methoxamine (10 μM) precontraction in endothelium-intact small mesenteric arteries (Figure 2(a), $\text{EC}_{50} = 7.7 \pm 0.05 \mu\text{M}$, $R_{\text{max}} = 87.9 \pm 2.0\%$; $n = 6$). The removal of endothelium significantly reduced this effect (Figure 2(a), $\text{EC}_{50} = 28.6 \pm 0.12 \mu\text{M}$, $R_{\text{max}} = 51.3 \pm 4.5\%$; $n = 6$). Next, we examined whether endothelium-derived NO was involved in this vasodilation. Preincubation with L-NAME (300 μM), a nitric oxide synthase inhibitor, partially reduced calycosin-induced vasodilation (Figure 2(b), $\text{EC}_{50} = 40.4 \pm 0.08 \mu\text{M}$, $R_{\text{max}} = 75.6 \pm 3.4\%$; $n = 5$). Notably, as shown in Figure 2(b), the inhibitory effects of endothelium denudation and L-NAME preincubation on calycosin-induced vasodilation were similar. However,




FIGURE 2: Concentration-response curves for calycosin-induced relaxation in the rat mesenteric arteries. (a) Vasorelaxation induced by calycosin with DMSO vehicle, or in the presence and absence of endothelium (** $P < 0.01$; ## $P < 0.01$). (b) Calycosin-induced vasorelaxation in the presence and absence of endothelium, or L-NAME (300 μM) pretreatment in endothelium-intact arteries (* $P < 0.05$; # $P < 0.05$ versus endothelium denuded). (c, d) Calycosin-induced vasorelaxation with pretreatments of either TEACl (3 mM; ** $P < 0.01$), glibenclamide (10 μM), apamin (50 nM) plus ChTX (50 nM), or IbTX (200 nM; ** $P < 0.01$). (e) Calycosin-induced vasorelaxation with IbTX (200 nM) pretreatment in endothelium-denuded arteries. (f) Calycosin-induced vasorelaxation with precontractions by methoxamine (10 μM) or KCl (60 mM; ** $P < 0.01$) in endothelium-intact arteries. Data were shown as mean \pm SEM. ChTX, charybdotoxin; IbTX, iberiotoxin; TEACl, tetraethylammonium chloride.

indomethacin (10 μM), cyclooxygenase inhibitor, did not affect the vasodilation by calycosin (data not shown).

Next, we examined whether K^+ channels were also involved in calycosin-induced vasodilation. With pretreatment of TEACl (3 mM), a nonspecific inhibitor of K^+ channels, the vasodilation effect was significantly reduced when compared to control (Figure 2(c), $\text{EC}_{50} = 25.1 \pm 0.12 \mu\text{M}$, $R_{\text{max}} = 58.3 \pm 5.1\%$; $n = 6$). Similarly, pretreatment with BK_{Ca} channel inhibitor, IbTX (200 nM), significantly reduced calycosin-induced vasodilation (Figure 2(d), $\text{EC}_{50} = 28.3 \pm 0.08 \mu\text{M}$, $R_{\text{max}} = 44.5 \pm 3.9\%$; $n = 6$). However, with pretreatment of K_{Ca} channel inhibitors, apamin (50 nM) plus

ChTX (50 nM), the vasodilation was reduced to a smaller extent (Figure 2(d), $\text{EC}_{50} = 20.7 \pm 0.12 \mu\text{M}$, $R_{\text{max}} = 56.0 \pm 4.3\%$; $n = 6$). Conversely, glibenclamide (10 μM), a K_{ATP} channel inhibitor, had no effect on calycosin-induced vasodilation (Figure 2(c), $\text{EC}_{50} = 8.9 \pm 0.08 \mu\text{M}$, $R_{\text{max}} = 76.4 \pm 6.2\%$; $n = 6$). Surprisingly, pretreatment with IbTX (200 nM) in endothelium-denuded arteries had no effect on calycosin-induced vasodilation (Figure 2(e)). Furthermore, calycosin-induced vasodilation was reduced with KCl (60 mM) precontraction compared to methoxamine precontraction in endothelium-intact arteries (Figure 2(f), $\text{EC}_{50} = 8.06 \pm 0.08 \mu\text{M}$, $R_{\text{max}} = 58.5 \pm 4.8\%$; $n = 5$). These data showed




FIGURE 3: Concentration–response curves for formononetin-induced relaxation in the rat mesenteric arteries. (a) Formononetin-induced vasorelaxation with DMSO vehicle, or in the presence and absence of endothelium (** $P < 0.01$; ## $P < 0.01$). (b) Formononetin-induced vasorelaxation in the presence and absence of endothelium, or L-NAME (300 μM) preincubation in endothelium-intact arteries (** $P < 0.01$; # $P < 0.05$ versus endothelium denuded). (c, d) Formononetin-induced vasorelaxation with pretreatments of either TEA (3 mM; ** $P < 0.01$), glibenclamide (10 μM ; ## $P < 0.01$), apamin (50 nM) plus ChTX (50 nM; ** $P < 0.01$), or IbTX (200 nM; ## $P < 0.01$). (e) Formononetin-induced vasorelaxation with IbTX (200 nM) preincubation in endothelium-denuded arteries. (f) Formononetin-induced vasorelaxation with precontractions of methoxamine (10 μM) or KCl in endothelium-intact arteries (60 mM; ** $P < 0.01$). Data were shown as mean \pm SEM. ChTX, charybdotoxin; IbTX, iberiotoxin; TEA, tetraethylammonium chloride.

that calycosin induced vasorelaxation via both endothelium-dependent and endothelium-independent pathways. More interestingly, the data also suggested that BK_{Ca} channels are closely related to the endothelium-dependent vasorelaxation.

3.2. Effects of Endothelial Removal, L-NAME, and K⁺ Channel Inhibitors on Vasodilation in Response to Formononetin. Formononetin also induced concentration-dependent vasodilation after methoxamine (10 μM) precontraction (Figure 3(a), $EC_{50} = 9.6 \pm 0.12 \mu\text{M}$, $R_{\text{max}} = 79.9 \pm 3.4\%$, $n = 6$), and removal of the endothelium (Figure 3(a), $EC_{50} = 13.9 \pm 0.09 \mu\text{M}$, $R_{\text{max}} = 63.6 \pm 3.2\%$, $n = 6$) or preincubation with L-NAME

(300 μM) (Figure 3(b), $EC_{50} = 42.8 \pm 0.30 \mu\text{M}$, $R_{\text{max}} = 56.3 \pm 4.5\%$, $n = 5$) significantly reduced this effect. On the other hand, indomethacin (10 μM) did not affect this vasodilation (data not shown).

As shown in Figure 3(c), formononetin-induced vasodilation was significantly inhibited with pretreatments of TEA (3 mM, $EC_{50} = 24.3 \pm 0.20 \mu\text{M}$, $R_{\text{max}} = 50.8 \pm 4.6\%$; $n = 6$) or glibenclamide (10 μM , $EC_{50} = 6.67 \pm 0.12 \mu\text{M}$, $R_{\text{max}} = 45.4 \pm 3.9\%$; $n = 7$). The vasodilation effect by formononetin was also reduced with pretreatment of IbTX (200 nM), or the combination of apamin plus ChTX (both 50 nM, Figure 3(d), IbTX: $EC_{50} = 23.7 \pm 0.18 \mu\text{M}$,

$R_{\max} = 43.6 \pm 4.2\%$; $n = 6$; $P < 0.01$; A+C: $EC_{50} = 27.8 \pm 0.17 \mu\text{M}$, $R_{\max} = 48.9 \pm 4.0\%$; $n = 6$). In addition, pretreatment with IbTX (200 nM) did not affect formononetin-induced vasodilation in endothelium-denuded arteries (Figure 3(e)). Formononetin-induced vasodilation was significantly reduced with KCl (60 mM) pre-contraction compared to methoxamine precontraction in endothelium-intact arteries (Figure 3(f), $EC_{50} = 4.38 \pm 0.05 \mu\text{M}$, $R_{\max} = 53.5 \pm 3.9\%$, $n = 5$). Similar to the effects of calycosin, these data showed that formononetin induced vasorelaxation via both endothelium-dependent and endothelium-independent pathways. More interestingly, the data also suggested that BK_{Ca} channels are closely related to the endothelium-dependent vasorelaxation.

3.3. Effect of Calycosin and Formononetin on NO Production and eNOS, iNOS, and nNOS Expression in HUVEC. In order to further determine whether these two isoflavonoids could regulate the production of NO and the expression of NOS in endothelial cells, HUVEC was employed as a cellular model. HUVEC is a widely *in vitro* cell model for the study of the regulation of endothelial function and the vascular diseases [31]. Figure 4(a) showed that calycosin increased NO production in a dose-dependent manner in HUVEC, and similar effect was also observed with formononetin (Figure 4(b)). There are three isoforms of NOS, including eNOS, inducible nitric oxide synthase (iNOS), and nNOS, which are responsible for the generation of NO in the vascular endothelium [32]. It has been reported that eNOS and nNOS are coexpressed in endothelial cells while iNOS is not [33]. Our results demonstrated that both calycosin (100 μM) and formononetin (100 μM) significantly induced the activation of eNOS (Figures 4(c) and 4(d)). In addition, both isoflavonoids also upregulated nNOS expression (Figures 4(c) and 4(d)). In contrast, iNOS expression was unaffected by neither calycosin nor formononetin (Figures 4(c) and 4(d)). These data suggested that calycosin and formononetin induced endothelium-dependent vasorelaxation via NO production through eNOS and nNOS.

3.4. Effect of Calycosin on Outward Currents in HUVEC. The activation of K_{Ca} is the major mechanism for EDH. We observed that exposure of calycosin induced dose-dependent outward currents in HUVEC recorded by whole cell patch clamp. The whole cell currents of HUVEC were recorded with 250 ms voltage steps between -100 mV and $+100$ mV from a holding potential of -60 mV. As shown in Figures 5(a) and 5(b), calycosin (10 nM–100 μM) increased dose-dependent outward currents in HUVEC ($n = 9$). At 100 μM calycosin, the current at $+100$ mV was significantly increased (Figures 5(a) and 5(c), 31.2 ± 1.7 pA/pF; $n = 7$) compared to control (9.9 ± 1.0 pA/pF), and this current was abolished by TEACL, a nonspecific inhibitor of K^+ channels (1 mM, Figures 5(c) and 5(e); 11.6 ± 1.5 pA/pF; $n = 6$).

Although IK_{Ca} and SK_{Ca} are the major K_{Ca} channels present in the endothelial cells, BK_{Ca} is also expressed moderately, and it was identified in HUVEC [22, 23]. We observed that K_{Ca} channels in calycosin induced outward

currents in HUVEC. An IK_{Ca} channel inhibitor, apamin (100 nM), had no effect on the outward currents induced by calycosin (Figure 5(e), $n = 6$). However, it was abolished by BK_{Ca} inhibitors, IbTX (200 nM, Figures 5(d) and 5(e); 13.8 ± 2.8 pA/pF; $n = 6$) and ChTX (200 nM, Figure 5(e); 14.9 ± 0.9 pA/pF; $n = 7$). In addition, by maintaining $[Ca^{2+}]_i$ at 250 nM, 500 nM, or 1000 nM, calycosin (100 μM) significantly increased the outward currents stimulated by single $+100$ mV step, compared with Ca^{2+} -free solution containing 10 mM EGTA (Figure 5(f), $n = 7-9$). These data strongly suggested that calycosin mainly activated endothelial BK_{Ca} channels but has minimal effects on IK_{Ca} or SK_{Ca} .

3.5. Effect of Formononetin on Outward Currents in HUVEC.







It was observed that formononetin (10 nM–100 μM) significantly increased the outward currents in a concentration-dependent manner in HUVEC, similar to the effects of calycosin (Figures 6(a) and 6(b), $n = 6-7$). At 100 μM , formononetin significantly increased the outward currents compared to control at $+100$ mV step (Figure 6(a), control: 1.4 ± 1.6 pA/pF, formononetin: 36.4 ± 3.1 pA/pF; $n = 6$). This current was abolished by TEACL (1 mM, Figures 6(c), and 6(e); 15.7 ± 3.6 pA/pF; $n = 6$), or IbTX (200 nM, Figures 6(d) and 6(e), 14.1 ± 0.9 pA/pF; $n = 5$), or ChTX (200 nM, Figure 6(e), 14.1 ± 0.9 pA/pF; $n = 5$). However, apamin (100 nM, Figure 6(e), 37.1 ± 3.2 pA/pF; $n = 6$) had no effect on these outward currents. Maintaining $[Ca^{2+}]_i$ at 250 nM, 500 nM, or 1000 nM, formononetin (100 μM) significantly increased the outward currents at $+100$ mV compared with Ca^{2+} -free solution containing 10 mM EGTA (Figure 6(f), $n = 6$). These data suggested that formononetin mainly activated endothelial BK_{Ca} channels but has minimal effects on IK_{Ca} or SK_{Ca} .

3.6. Effect of Calycosin and Formononetin on $[Ca^{2+}]_i$ in HUVEC.

Endothelial $[Ca^{2+}]_i$ elevation has been implicated in endothelium-mediated vasodilation [18] and is also needed for K_{Ca} activation [34]. Since calycosin and formononetin increased outward currents via endothelial K_{Ca} channel, we next examined their effects on $[Ca^{2+}]_i$ in HUVEC. As expected, we observed a rapid increase in $[Ca^{2+}]_i$ in response to calycosin (Figures 7(a) and 7(b)). Similar results also showed that formononetin evoked Ca^{2+} influx (Figures 7(a) and 7(b)). Both of these two drugs induced approximately 45% increases in $[Ca^{2+}]_i$ compared to untreated cells. Therefore, these data suggested that calycosin and formononetin activated endothelial BK_{Ca} channels probably by increasing $[Ca^{2+}]_i$.

4. Discussion

In the present study, we investigated the endothelial beneficial effects of calycosin and formononetin, two isoflavonoids isolated from the well-known antihypertensive herb, Radix Astragali, in rat small resistance arteries and HUVEC. The chemical structures of calycosin and formononetin are very similar, with an extra hydroxyl group at C-3 position of



(c)

FIGURE 4: Continued.




FIGURE 4: Calycosin and formononetin induced NO production via eNOS and nNOS pathways in HUVEC. (a, b) NO level was determined by a NO assay kit. HUVEC was incubated with DMSO, calycosin (1–100 μM), or formononetin (1–100 μM) for 1 h ($n = 5$). (c, d) Representative immunoblots and graphs for the protein expressions of eNOS, phosphorylation of eNOS, nNOS, iNOS, or GAPDH after (c) calycosin (1–100 μM) or (d) formononetin (1–100 μM) treatment for 1 h ($n = 3-4$). Data were shown as mean \pm SEM. * $P < 0.05$, *** $P < 0.001$ versus untreated cells.

the B-ring in calycosin [35]. Both isoflavonoids were shown to provide beneficial effects in vascular tone regulation and improvement in endothelial and cardiovascular dysfunction [3–6].

Here, we demonstrated that calycosin and formononetin produced very similar effects in rat small mesenteric arteries, that is, inducing vasorelaxation through endothelium-dependent and endothelium-independent pathways. We observed that relaxations elicited by both isoflavonoids could be reduced by endothelial disruption and were sensitive to L-NAME (inhibitor of NOS), apamin plus charybdotoxin (inhibitors of SK_{Ca} and IK_{Ca}), and iberiotoxin (IbTX)

(inhibitor of BK_{Ca}). Notably, the sensitivity to IbTX was only observed in endothelium-intact, but not in endothelium-denuded vessels, indicating the involvement of BK_{Ca} channels present in the endothelium. The relaxation elicited by both isoflavonoids was also reduced (to a lesser extent) when the arteries were contracted with high KCl (60 mM) depolarizing solution, indicating the inhibition of VDCC was also partly involved. Pretreatment with indomethacin to inhibit prostaglandin production did not affect the vasodilation, indicating that PGI_2 was not involved. Although calycosin and formononetin are very similar in structure, differential properties of their effects were also observed. Our data




FIGURE 5: Calycosin increased outward currents in HUVEC through BK_{Ca} channel. (a) Current-voltage (*I/V*) relationship in response to calycosin (1–100 μM). (b) Dose-response curve for whole cell recording of currents at +100 mV with different concentrations of calycosin (10 nM–100 μM). (c, d) Representative trace of currents that were recorded in response to calycosin in the absence or presence of (c) TEACl (1 mM) or (d) IbTX (200 nM). (e, f) Whole cell recording of currents at +100 mV in response to calycosin (100 μM) in the presence of (e) apamin (200 nM), IbTX (200 nM), ChTX (200 nM), or TEACl (1 mM; ****P* < 0.001 versus control; ##*P* < 0.01 and ###*P* < 0.001 versus calycosin-treated cells), or (f) with different [Ca²⁺]_i as indicated (***P* < 0.01 and ****P* < 0.001 versus calycosin-treated cells with free [Ca²⁺]_i). Data were shown as mean ± SEM. ChTX, charybdotoxin; IbTX, iberiotoxin; TEACl, tetraethylammonium chloride.

showed that relaxation induced by formononetin was sensitive to glibenclamide (K_{ATP} channel inhibitor), but not calycosin-induced relaxation. Besides, we also observed some discrepancies in the effects of calycosin and formononetin in blood vessels from various vascular beds. Previous study

intact aortic rings reported that the relaxation elicited by calycosin was endothelium-independent by acting as Ca²⁺ channel blocker [7]. However, here in rat mesenteric arteries we observed that calycosin-elicited relaxation is both endothelium-dependent and endothelium-independent. For




FIGURE 6: Formononetin increased outward currents in HUVEC through BK_{Ca} channel. (a) *I/V* graph in response to formononetin (1–100 μM). (b) Dose-response curve for whole cell recording of currents at +100 mV with different concentrations of formononetin (10 nM–100 μM). (c, d) Representative trace of currents that were recorded in response to formononetin, in the absence and presence of (c) TEACl (1 mM) or (d) IbTX (200 nM). (e, f) Whole cell recording of currents at +100 mV in response to formononetin (100 μM) in the presence of (e) apamin (200 nM), IbTX (200 nM), ChTX (200 nM), or TEACl (1 mM); *** P < 0.001 versus control; ** P < 0.01 and *** P < 0.001 versus formononetin-treated cells), or (f) with different $[\text{Ca}^{2+}]_i$ as indicated (** P < 0.001 versus formononetin-treated cells with free $[\text{Ca}^{2+}]_i$). Data were shown as mean \pm SEM. ChTX, charybdotoxin; IbTX, iberiotoxin; TEACl, tetraethylammonium chloride.

formononetin, results from previous studies in rat aortic rings showed that formononetin elicited relaxation through endothelium-dependent pathway involving NO synthesis, and endothelium-independent involving iberiotoxin-(IbTX-) sensitive BK_{Ca} channel, glibenclamide-sensitive

K_{ATP} channel, and the inhibition of VDCC [5, 6, 36]. Here in rat mesenteric arteries, we observed very similar effects of formononetin, but in our preparation the sensitivity to IbTX was only observed in endothelium-intact vessels, indicating a more important role of endothelial BK_{Ca} channels in the




FIGURE 7: Calycosin and formononetin-induced Ca^{2+} response in HUVEC. (a, b) HUVEC was loaded with Fluo-4 in Tyrode solution containing 2 mM Ca^{2+} . Representative graph and relative fluorescence intensity in intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$), evoked by calycosin (100 μ M) and formononetin (100 μ M) over the time course ($n = 3$). Data were shown as mean \pm SEM. * $P < 0.05$ versus control.

mesenteric resistance arteries. The discrepancies observed for calycosin and formononetin in rat aorta and mesenteric arteries may be explained by the differential physiological characteristics exhibited by conduit and resistance arteries for vascular homeostasis. In fact, it has been recognized that endothelium-dependent vasoactivities are more pronounced in small resistance arteries than in large conduit arteries such as aorta, and small resistance arteries are closely related to endothelial dysfunction [10, 12]. Thus, our results suggested that calycosin and formononetin could effectively promote endothelial functions in the small resistance arteries, and endothelial BK_{Ca} channels might have an important role.

BK_{Ca} channels are mainly expressed in vascular smooth muscle cells, but recent studies demonstrated that BK_{Ca}

channels are also present in vascular endothelium or freshly isolated endothelial cells from the small resistance arteries [19, 20]. When expressed in endothelial cells, BK_{Ca} channels are observed to regulate NO and EDH production, and their important roles have been implicated in disease conditions such as ischemic hypoxia [19, 23, 37, 38]. In line with this, we demonstrated that both calycosin and formononetin significantly increased NO production and upregulated the activities and expressions of eNOS and nNOS, without affecting iNOS in HUVEC. It has been suggested that endothelial BK_{Ca} channel might serve as a compensatory mechanism for improving vasoreactivity in disease environments such as hypertension and could be a potential therapeutic target for the regulation of blood pressure and flow through increased NO production vascular hyperpolarization [20, 39].

By using whole cell patch clamp, we further showed that IbTX-sensitive outward-rectifying currents were induced by exposure to calycosin or formononetin in a dose-dependent manner in HUVEC, an endothelial cell model commonly used for endothelial functions. Moreover, these currents were also sensitive to charybdotoxin (also an inhibitor at BK_{Ca} channel), but not apamin (selective SK_{Ca} channel inhibitor). Buffering intracellular Ca^{2+} by EGTA significantly reduced calycosin- and formononetin-induced outward currents, indicating that the activation of the K^+ channels was dependent on intracellular calcium which further indicated the characteristic of BK_{Ca} channel [40]. We further demonstrated that calycosin and formononetin could induced an $[Ca^{2+}]_i$ increase in HUVEC, providing more evidences that calycosin and formononetin activate endothelial BK_{Ca} channel by stimulating $[Ca^{2+}]_i$ increase.

Until recently, the emerging view of cerebrovascular dysregulation was implicated not only in cerebrovascular diseases, such as stroke, but also in neurodegeneration, like Alzheimer's disease (AD) [41]. Particularly, the inhibition of BK_{Ca} activity was observed in 3xTg AD model mice and might be involved in early dysfunction in the AD brain [42]. Several studies suggested that the activation of BK_{Ca} channels might be new therapeutic target for improving behavioral and cognitive disorders [43, 44]. Since our results demonstrated that calycosin and formononetin act as novel BK_{Ca} activators and also played a role in the regulation in small resistance arteries, we postulated these two isoflavonoids might have potential effects on cognitive disorders through the regulation of cerebral microcirculation by activating BK_{Ca} channels.

5. Conclusions

In summary, our findings demonstrated that calycosin and formononetin induced vasodilation in rat small mesenteric arteries involving both the endothelium and the vascular smooth muscle. The endothelium-dependent responses were associated with eNOS/nNOS-dependent NO production and endothelium hyperpolarization, possibly by directly activating endothelial BK_{Ca} channels. Therefore, we suggested that these isoflavonoids might provide potential therapeutic regiments to improve endothelial functions for treating

diseases related to abnormal vascular alteration, such as hypertension, cardiovascular diseases, and cerebrovascular-circulation related cognitive disorders such as stroke and vascular dementia.

Abbreviations

BK _{Ca} :	Large-conductance Ca ²⁺ -activated K ⁺
ChTX:	Charybdotoxin
EC ₅₀ :	Concentration producing 50% of the maximum effect
EDHF:	Endothelium-derived hyperpolarizing factor
Em:	Membrane potential
HUVEC:	Human umbilical vein endothelial cells
IbTX:	Iberiotoxin
K _{ATP} :	ATP-sensitive potassium channels
K _{Ca} :	Ca ²⁺ -activated K ⁺
L-NAME:	N _ω -nitro-L-arginine methyl ester
NO:	Nitric oxide
NOS:	Nitric oxide synthase
eNOS:	Endothelial nitric oxide synthase
iNOS:	Inducible nitric oxide synthase
nNOS:	Neuronal nitric oxide synthase
PGI ₂ :	Prostacyclin
TEACl:	Tetraethylammonium chloride
VDCC:	Voltage-dependent calcium channel.

Competing Interests

The authors report no conflict of interests.



Acknowledgments

This work was supported by grants from Science and Technology Development Fund of Macau SAR [FDCT 127/2014/A3], Research Committee at University of Macau [MYRG124(Y1-L3)-ICMS12-HPM], and National Natural Science Foundation of China [NSFC-81403139-H2809].

References

- [1] X.-L. Xu, H. Ji, S.-Y. Gu, Q. Shao, Q.-J. Huang, and Y.-P. Cheng, "Cardioprotective effects of Astragali Radix against isoproterenol-induced myocardial injury in rats and its possible mechanism," *Phytotherapy Research*, vol. 22, no. 3, pp. 389–394, 2008.
- [2] A. L. Miller, "Botanical influences on cardiovascular disease," *Alternative Medicine Review*, vol. 3, no. 6, pp. 422–431, 1998.
- [3] Y.-H. Jiang, W. Sun, W. Li et al., "Calycosin-7-O-β-D-glucoside promotes oxidative stress-induced cytoskeleton reorganization through integrin-linked kinase signaling pathway in vascular endothelial cells," *BMC Complementary and Alternative Medicine*, vol. 15, no. 1, article 315, 2015.
- [4] T. Sun, J. Wang, L.-H. Huang, and Y.-X. Cao, "Antihypertensive effect of formononetin through regulating the expressions of eNOS, 5-HT_{2A/1B} receptors and α₁-adrenoceptors in spontaneously rat arteries," *European Journal of Pharmacology*, vol. 699, no. 1–3, pp. 241–249, 2013.
- [5] T. Sun, R. Liu, and Y.-X. Cao, "Vasorelaxant and antihypertensive effects of formononetin through endothelium-dependent and -independent mechanisms," *Acta Pharmacologica Sinica*, vol. 32, no. 8, pp. 1009–1018, 2011.
- [6] J.-H. Wu, Q. Li, M.-Y. Wu et al., "Formononetin, an isoflavone, relaxes rat isolated aorta through endothelium-dependent and endothelium-independent pathways," *Journal of Nutritional Biochemistry*, vol. 21, no. 7, pp. 613–620, 2010.
- [7] X.-L. Wu, Y.-Y. Wang, J. Cheng, and Y.-Y. Zhao, "Calcium channel blocking activity of calycosin, a major active component of Astragali Radix, on rat aorta," *Acta Pharmacologica Sinica*, vol. 27, no. 8, pp. 1007–1012, 2006.
- [8] H. Zhu, L. Zou, J. Tian, F. Lin, J. He, and J. Hou, "Protective effects of sulphonated formononetin in a rat model of cerebral ischemia and reperfusion injury," *Planta Medica*, vol. 80, no. 4, pp. 262–268, 2014.
- [9] S. Fu, Y. Gu, J.-Q. Jiang et al., "Calycosin-7-O-β-D-glucoside regulates nitric oxide /caveolin-1/matrix metalloproteinases pathway and protects blood-brain barrier integrity in experimental cerebral ischemia-reperfusion injury," *Journal of Ethnopharmacology*, vol. 155, no. 1, pp. 692–701, 2014.
- [10] K.-T. Kang, "Endothelium-derived relaxing factors of small resistance arteries in hypertension," *Toxicological Research*, vol. 30, no. 3, pp. 141–148, 2014.
- [11] H. D. Intengan and E. L. Schiffrin, "Structure and mechanical properties of resistance arteries in hypertension: role of adhesion molecules and extracellular matrix determinants," *Hypertension*, vol. 36, no. 3, pp. 312–318, 2000.
- [12] C. J. Garland and K. A. Dora, "EDH: endothelium-dependent hyperpolarization and microvascular signalling," *Acta Physiologica*, 2016.
- [13] I. Grgic, B. P. Kaistha, J. Hoyer, and R. Köhler, "Endothelial Ca²⁺-activated K⁺ channels in normal and impaired EDHF-dilator responses-relevance to cardiovascular pathologies and drug discovery," *British Journal of Pharmacology*, vol. 157, no. 4, pp. 509–526, 2009.
- [14] C. G. Sobey, "Potassium channel function in vascular disease," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 21, no. 1, pp. 28–38, 2001.
- [15] R. Köhler, B. P. Kaistha, and H. Wulff, "Vascular K_{Ca}-channels as therapeutic targets in hypertension and restenosis disease," *Expert Opinion on Therapeutic Targets*, vol. 14, no. 2, pp. 143–155, 2010.
- [16] T. Dalsgaard, C. Kroigaard, and U. Simonsen, "Calcium-activated potassium channels—a therapeutic target for modulating nitric oxide in cardiovascular disease?" *Expert Opinion on Therapeutic Targets*, vol. 14, no. 8, pp. 825–837, 2010.
- [17] T. Dalsgaard, C. Kroigaard, M. Misfeldt, T. Bek, and U. Simonsen, "Openers of small conductance calcium-activated potassium channels selectively enhance NO-mediated bradykinin vasodilatation in porcine retinal arterioles," *British Journal of Pharmacology*, vol. 160, no. 6, pp. 1496–1508, 2010.
- [18] B. Nilius, F. Viana, and G. Droogmans, "Ion channels in vascular endothelium," *Annual Review of Physiology*, vol. 59, pp. 145–170, 1997.
- [19] O. Jackson-Weaver, J. M. Osmond, M. A. Riddle et al., "Hydrogen sulfide dilates rat mesenteric arteries by activating endothelial large-conductance Ca²⁺-activated K⁺ channels and smooth

- muscle Ca^{2+} sparks," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 304, no. 11, pp. H1446–H1454, 2013.
- [20] A. Vang, J. Mazer, B. Casserly, and G. Choudhary, "Activation of endothelial BKCa channels causes pulmonary vasodilation," *Vascular Pharmacology*, vol. 53, no. 3–4, pp. 122–129, 2010.
- [21] Y. C. Xu, G. P. H. Leung, P. Y. D. Wong, P. M. Vanhoutte, and R. Y. K. Man, "Kaempferol stimulates large conductance Ca^{2+} -activated K^+ (BKCa) channels in human umbilical vein endothelial cells via a cAMP/PKA-dependent pathway," *British Journal of Pharmacology*, vol. 154, no. 6, pp. 1247–1253, 2008.
- [22] M. Begg, F.-M. Mo, L. Offertaler et al., "G Protein-coupled endothelial receptor for atypical cannabinoid ligands modulates a Ca^{2+} -dependent K^+ Current," *Journal of Biological Chemistry*, vol. 278, no. 46, pp. 46188–46194, 2003.
- [23] M. Feletou, "Calcium-activated potassium channels and endothelial dysfunction: therapeutic options?" *British Journal of Pharmacology*, vol. 156, no. 4, pp. 545–562, 2009.
- [24] T. Chang, L. Wu, and R. Wang, "Altered expression of BK channel β_1 subunit in vascular tissues from spontaneously hypertensive rats," *American Journal of Hypertension*, vol. 19, no. 7, pp. 678–685, 2006.
- [25] J. L. Carvalho-de-Souza, W. A. Varanda, R. C. Tostes, and A. Z. Chignalia, "BK channels in cardiovascular diseases and aging," *Aging and Disease*, vol. 4, no. 1, pp. 38–49, 2013.
- [26] S. Brakemeier, I. Eichler, A. Knorr, T. Fassheber, R. Kohler, and J. Hoyer, "Modulation of Ca^{2+} -activated K^+ channel in renal artery endothelium in situ by nitric oxide and reactive oxygen species," *Kidney International*, vol. 64, no. 1, pp. 199–207, 2003.
- [27] Y. Liu and D. D. Gutterman, "The coronary circulation in diabetes: influence of reactive oxygen species on K^+ channel-mediated vasodilation," *Vascular Pharmacology*, vol. 38, no. 1, pp. 43–49, 2002.
- [28] R. White and C. R. Hiley, "A comparison of EDHF-mediated and anandamide-induced relaxations in the rat isolated mesenteric artery," *British Journal of Pharmacology*, vol. 122, no. 8, pp. 1573–1584, 1997.
- [29] B. Li, W. Jie, L. Huang et al., "Nuclear BK channels regulate gene expression via the control of nuclear calcium signaling," *Nature Neuroscience*, vol. 17, no. 8, pp. 1055–1063, 2014.
- [30] P. Tep-areenan, D. A. Kendall, and M. D. Randall, "Testosterone-induced vasorelaxation in the rat mesenteric arterial bed is mediated predominantly via potassium channels," *British Journal of Pharmacology*, vol. 135, no. 3, pp. 735–740, 2002.
- [31] H.-J. Park, Y. Zhang, S. P. Georgescu, K. L. Johnson, D. Kong, and J. B. Galper, "Human umbilical vein endothelial cells and human dermal microvascular endothelial cells offer new insights into the relationship between lipid metabolism and angiogenesis," *Stem Cell Reviews*, vol. 2, no. 2, pp. 93–102, 2006.
- [32] U. Forstermann and W. C. Sessa, "Nitric oxide synthases: regulation and function," *European Heart Journal*, vol. 33, no. 7, pp. 829–837, 2012.
- [33] T. Bachetti, L. Comini, S. Curello et al., "Co-expression and modulation of neuronal and endothelial nitric oxide synthase in human endothelial cells," *Journal of Molecular and Cellular Cardiology*, vol. 37, no. 5, pp. 939–945, 2004.
- [34] C. M. Fanger, S. Ghanshani, N. J. Logsdon et al., "Calmodulin mediates calcium-dependent activation of the intermediate conductance K_{Ca} channel, IKCa1 ," *The Journal of Biological Chemistry*, vol. 274, no. 9, pp. 5746–5754, 1999.
- [35] T. Wu, S. W. Annie Bligh, L.-H. Gu et al., "Simultaneous determination of six isoflavonoids in commercial Radix Astragali by HPLC-UV," *Fitoterapia*, vol. 76, no. 2, pp. 157–165, 2005.
- [36] Y. Zhao, B.-N. Chen, S.-B. Wang, S.-H. Wang, and G.-H. Du, "Vasorelaxant effect of formononetin in the rat thoracic aorta and its mechanisms," *Journal of Asian Natural Products Research*, vol. 14, no. 1, pp. 46–54, 2012.
- [37] A. M. Briones, A. S. Padilha, A. L. Cogolludo et al., "Activation of BKCa channels by nitric oxide prevents coronary artery endothelial dysfunction in ouabain-induced hypertensive rats," *Journal of Hypertension*, vol. 27, no. 1, pp. 83–91, 2009.
- [38] J. M. Hughes, M. A. Riddle, M. L. Paffett, L. V. Gonzalez Bosc, and B. R. Walker, "Novel role of endothelial BK Ca channels in altered vasoreactivity following hypoxia," *American Journal of Physiology—Heart and Circulatory Physiology*, vol. 299, no. 5, pp. H1439–H1450, 2010.
- [39] V. Calderone, A. Martelli, L. Testai, E. Martinotti, and M. C. Breschi, "Functional contribution of the endothelial component to the vasorelaxing effect of resveratrol and NS 1619, activators of the large-conductance calcium-activated potassium channels," *Naunyn-Schmiedeberg's Archives of Pharmacology*, vol. 375, no. 1, pp. 73–80, 2007.
- [40] J. Cui, D. H. Cox, and R. W. Aldrich, "Intrinsic voltage dependence and Ca^{2+} regulation of mslo large conductance Ca -activated K^+ channels," *Journal of General Physiology*, vol. 109, no. 5, pp. 647–673, 1997.
- [41] C. Iadecola, "Neurovascular regulation in the normal brain and in Alzheimer's disease," *Nature Reviews Neuroscience*, vol. 5, no. 5, pp. 347–360, 2004.
- [42] K. Yamamoto, Y. Ueta, L. Wang et al., "Suppression of a neocortical potassium channel activity by intracellular amyloid- β and its rescue with homer1a," *Journal of Neuroscience*, vol. 31, no. 31, pp. 11100–11109, 2011.
- [43] L. Wang, H. Kang, Y. Li et al., "Cognitive recovery by chronic activation of the large-conductance calcium-activated potassium channel in a mouse model of Alzheimer's disease," *Neuropharmacology*, vol. 92, pp. 8–15, 2015.
- [44] H. H. Dietrich, C. Xiang, B. H. Han, G. J. Zipfel, and D. M. Holtzman, "Soluble amyloid- β effect on cerebral arteriolar regulation and vascular cells," *Molecular Neurodegeneration*, vol. 5, no. 1, article 15, 2010.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

