

miR-200c Prevents TGF- β 1-Induced Epithelial-to-Mesenchymal Transition and Fibrogenesis in Mesothelial Cells by Targeting ZEB2 and Notch1

Jessica Y.S. Chu,¹ Mel K.M. Chau,¹ Caleb C.Y. Chan,¹ Andrew C.P. Tai,¹ Kwok Fan Cheung,¹ Tak Mao Chan,¹ and Susan Yung¹

¹Department of Medicine, The University of Hong Kong, Pok Fu Lam, Hong Kong

Peritoneal fibrosis and loss of transport function is a common complication contributing to adverse outcomes in patients on long-term peritoneal dialysis (PD). Epithelial-to-mesenchymal transition (EMT) in mesothelial cells is a salient feature, but its triggering mechanisms remain obscure. Dysregulation of microRNA (miR) expression is implicated in EMT and tissue fibrosis. We investigated the role of miR-200c in EMT and fibrogenesis in a murine PD model and in cultured peritoneal mesothelial cells. PD-fluid-treated mice showed peritoneal miR-200c expression reduced by 76.2% compared with PBS-treated mice, and this was accompanied by increased peritoneal α -smooth muscle actin, fibronectin, and collagen expression. PD fluid and TGF- β 1 both reduced miR-200c expression in cultured mesothelial cells, accompanied by downregulation of E-cadherin and decorin, and induction of fibronectin, collagen I and III, and transcription factors related to EMT. Decorin prevented the suppression of miR-200c by TGF- β 1. Lentivirus-mediated miR-200c overexpression prevented the induction of fibronectin, collagen I, and collagen III by TGF- β 1, independent of decorin, and partially prevented E-cadherin suppression by TGF- β 1. Target genes of miR-200c were identified as ZEB2 and Notch1. Our data demonstrate that miR-200c regulates EMT and fibrogenesis in mesothelial cells, and loss of peritoneal miR-200c contributes to PD-associated peritoneal fibrosis.

INTRODUCTION

Peritoneal dialysis (PD) accounts for approximately 15% of the global dialysis population, and its success depends on the structural and functional integrity of the peritoneal membrane.¹ Yet progressive peritoneal fibrosis and loss of dialytic function is near universal, especially in patients with long durations of PD, and results in inadequate dialysis or fluid removal or the uncommon but devastating complication of encapsulating peritoneal sclerosis.²⁻⁴ The normal peritoneal membrane is lined by a monolayer of mesothelial cells, under which is the submesothelium, which includes collagen fibrils, blood vessels, and a small number of fibroblasts. Peritoneal mesothelial cells provide the first line of defense against chemical and bacterial injury to the peritoneum and play an important role in reparative processes. Mesothelial cells are in perennial contact with peritoneal dialysate during

long-term PD. There is much evidence that unphysiological glucose-based PD fluids contribute to progressive peritoneal fibrosis and epithelial-to-mesenchymal transition (EMT) in mesothelial cells, accompanied by mesothelial denudation, increased matrix proteins in the submesothelium, and progressive subendothelial hyalinization of blood vessels with narrowing or obliteration of the vascular lumen.^{2,5,6} Also, TGF- β 1 has been demonstrated as a key mediator of tissue fibrosis and EMT.⁷⁻⁹ However, the mechanisms leading to abnormalities in peritoneal mesothelial cells and the peritoneal membrane remain to be fully elucidated.

MicroRNAs (miRs) are a family of endogenous, short non-coding RNAs that regulate gene expression at the post-transcriptional level, either through mRNA destabilization or translational inhibition of their specific targets.¹⁰ The human miRome contains more than 2,500 miRs, each having various expression patterns, ranging from ubiquitous to highly tissue specific.¹¹ Each miR can recognize over 100 target genes though sequence complementarity between the miR and binding sites in the 3' UTR of target mRNAs. Dysregulated miR expression has been observed in pathological conditions, such as cancer and tissue fibrosis, and several miRs have been implicated in the regulation of EMT transcription factors, resulting in repression or promotion of EMT and fibrosis. In this respect, miR-21 has been shown to promote peritoneal fibrosis,^{12,13} whereas miR-15a-5p, miR-30a, miR-129-5p, and miR-302c have been shown to suppress peritoneal fibrosis.¹⁴⁻¹⁷

The role of miR-200c in cancer and metastasis has been extensively studied, and the results showed that it controls EMT by targeting zinc finger E-box binding homeobox 1 (ZEB1), ZEB2, and Notch1, repressors of E-cadherin transcription that facilitate the dissolution of epithelial cell-cell contacts.¹⁸⁻²⁰ The aim of this study was to

Received 8 February 2019; accepted 12 May 2019;
<https://doi.org/10.1016/j.omtn.2019.05.008>.

Correspondence: Susan Yung, Department of Medicine, The University of Hong Kong, Pokfulam Road, Pok Fu Lam, Hong Kong.

E-mail: ssyung@hku.hk

Correspondence: Tak Mao Chan, Department of Medicine, The University of Hong Kong, Pokfulam Road, Pok Fu Lam, Hong Kong.

E-mail: dtmchan@hku.hk






Figure 1. Peritoneal Membrane Expression of miR-29a, -192, -200b, -200c, -324, and -377 and the Effect of Peritoneal Dialysate

(A) Mice were randomized to receive sterile PBS (○) or 4.25% glucose-based Dianeal PD fluid (+) (2 mL) twice daily for up to 30 days (n = 3 mice for days 1, 3, 5, 10, and 15; n = 6 for day 30). Peritoneal membrane expression of (A) miR-29a, (B) miR-192, (C) miR-200b, (D) miR-200c, (E) miR-324, and (F) miR-377 was determined by real-time PCR. For each miR, data were normalized to the level expressed on day 1 in PBS-treated mice. Mean Ct values were 22.2 and 22.6 for miR-29a and U6, respectively; 34.8 and 22.6 for miR-192 and U6, respectively; 32.9 and 22.7 for miR-200b and U6, respectively; 33.7 and 22.6 for miR-200c and U6, respectively; 29.2 and 22.6 for miR-324 and U6, respectively; and 35.2 and 22.5 for miR-377 and U6, respectively, for PBS-treated mice on day 1. *p < 0.05, **p < 0.01, and ***p < 0.001, PBS versus PD fluid for the same time point; ##p < 0.01 and ###p < 0.001, compared to day 1 for the same treatment. Data analyzed by using ANOVA.

protective effects on EMT and fibrogenesis in the peritoneum, mediated through specific gene targets.

RESULTS

PD Fluid Decreased Peritoneal miR-200c Expression

We examined peritoneal expression of 6 miRNos in a murine PD model. These miRNos were chosen based on their ability to regulate EMT in cancer cells or renal fibrosis, as reported in the literature.^{21–28} In PBS-treated mice, miR-29a, -192, -200b, -200c, -324, and -377 were all detected in peritoneal membrane specimens by real-time PCR, and the levels remained stable throughout the study period (Figure 1).

In mice treated with PD fluid, miR-192 was reduced by 29.0% after 1 day, whereas miR-200c was reduced by 76.2% after 3 days of PD (p < 0.05 and p < 0.001 respectively, compared with PBS-treated mice). The suppressive effect of PD fluid on miR-29a, -200b, -324, and -377 was evident after 5 days, with 35.1%, 44.0%, 44.3%, and 39.8% reduction, respectively (Figure 1). Repression of all six miRNos by PD fluid was sustained for 1 month. Since the reduction of miR-200c expression was most pronounced, we focused on miR-200c in subsequent studies. *In situ* hybridization showed a marked reduction in peritoneal miR-200c expression after 1 month of PD (Figure 2A). Peritoneal membrane thickening showed no significant difference between the treatment groups after 1 month (p = 0.06). In PBS-treated mice, peritoneal membrane specimens showed constitutive collagen expression and sparse cells in the submesothelium, and weak α -smooth muscle actin and fibronectin expression in the mesothelial but not submesothelial compartment. In PD-fluid-treated mice, fibronectin

investigate the role of miR-200c in peritoneal fibrosis and peritoneal mesothelial cell EMT, and also its target genes, in the context of PD. Importantly, our results show that conventional glucose-based PD fluid markedly reduced peritoneal miR-200c expression in C57BL/6N mice, and this was accompanied by increased α -smooth muscle actin, fibronectin, and collagen expression. PD fluid and TGF- β 1 both reduced miR-200c expression in cultured mesothelial cells, and this was accompanied by E-cadherin downregulation and induction of the EMT transcription factors, fibronectin, collagen I, and collagen III. We also demonstrated that lentivirus-mediated miR-200c overexpression in mesothelial cells attenuated TGF- β 1-induced EMT transcription factors and mesenchymal markers, and partially reversed the TGF- β 1 effect of reducing E-cadherin expression. Luciferase reporter assays demonstrated that miR-200c directly targeted ZEB2 and Notch1, and indirectly targeted Jagged2. Overall, our results imply that miR-200c has an important role in the pathogenesis of PD-associated peritoneal fibrosis, through its




Figure 2. Expression of miR-200c in the Peritoneal Membrane of C57BL/6N Mice

(A) Locked nucleic acid *in situ* hybridization showing miR-200c expression in peritoneal membrane specimens from mice treated with PBS or PD fluid for 1 month. Blue denotes miR-200c staining (arrow). (B) Representative images of H&E staining (a and b), α -smooth muscle actin (α -SMA, c and d), fibronectin (FN, e and f), and collagen (Masson's trichrome staining, g and h) expression in mice treated with PBS (a, c, e, and g) or PD fluid (b, d, f, and h). Arrows indicate increased α -SMA-positive cells in both the mesothelium and submesothelium and FN expression in the mesothelium in PD-fluid-treated mice. Peritoneal membrane thickness in H&E-stained specimens was determined in PBS- and PD-fluid-treated mice, by using Axiovision 4.7 software. Original magnification, $\times 200$ for H&E staining and miR-200c expression, and $\times 400$ for α -SMA, FN, and collagen expression ($n = 6$ for each group). Data analyzed with the Kolmogorov-Smirnov test, which assesses the whole distribution rather than the mean.

expression was increased in the mesothelium, and a marked increase in α -smooth-muscle-actin-positive cells showing an elongated appearance was noted in an intense collagen-rich submesothelium (Figure 2B).


Effect of Glucose-Based PD Fluid and TGF- β 1 on Mesothelial Cell miR-200c Expression

miR-200c was constitutively expressed in mesothelial cells. Incubation of mesothelial cells with 1.36% glucose-based (Dianeal®) PD fluid for 24 h decreased miR-200c expression by $21.8 \pm 4.5\%$

compared to serum-free medium (SFM) ($p < 0.001$) (Figure 3A). TGF- β 1 has been reported to contribute to PD-associated peritoneal fibrosis in mice.²⁹ Exposure of mesothelial cells to TGF- β 1 decreased miR-200c expression in a dose-dependent manner, with 40.7% reduction at 10 ng/mL TGF- β 1 compared with SFM ($p < 0.001$). Repression of miR-200c was accompanied by a significant decrease in E-cadherin and decorin expression and increased expression of mesenchymal markers ZEB2, Notch1, Jagged2, SNAIL, FSP-1, α -smooth muscle actin, fibronectin, and collagen I at transcription and/or translation (Figures 3, 4, 5, and 6). TGF- β 1 had no effect on vimentin expression (data not shown). The induction of EMT transcription factors and mesenchymal markers by TGF- β 1 was apparent after 24 h, but the effect of TGF- β 1 on phenotypic changes in mesothelial cells was observed only after 48 h (Figure 3J). Decorin has been shown to inhibit the profibrotic effects of TGF- β 1.³⁰ Incubating the cells with decorin prior to stimulation with TGF- β 1 attenuated the suppressive effect of TGF- β 1 on miR-200c expression (Figure 3B).

Overexpression of miR-200c Represses Mediators of EMT and Fibrogenesis in Mesothelial Cells

Mesothelial cells transfected with scrambled control hairpin (lenti-scramble) showed expression of miR-200c, E-cadherin, ZEB2, Notch1, Jagged2, SNAIL, fibronectin, collagen I, and collagen III similar to that of non-transfected cells (Figures 5 and 6). Primary human peritoneal mesothelial cells transfected with lentivector-based



(legend on next page)

miR precursor constructs (lenti-miR-200c) showed a 106.7 ± 10.1 - and 113.9 ± 4.6 -fold increase of miR-200c expression when compared with non-transfected and lenti-scramble-transfected cells, respectively ($p < 0.001$, for both; **Figure 5A**); and lenti-miR-200c increased miR-200c expression in Met-5A cells by 414.7 ± 65.5 - and 419.8 ± 74.5 -fold, respectively ($p < 0.001$, for both; **Figure 5B**). miR-200c overexpression had no effect on cell morphology (data not shown) or constitutive gene and protein expression of E-cadherin, decorin, EMT transcription factors, or fibronectin, but significantly decreased collagen I and III expression (**Figures 5 and 6; Table 1**). In cells stimulated with TGF- β 1, miR-200c overexpression attenuated TGF- β 1-induced ZEB2, Notch1, Jagged2, SNAIL, fibronectin, and collagen I and III expression and partially reversed the suppression of E-cadherin transcription but not translation, by TGF- β 1 (**Figures 5 and 6**).

miR-200c Directly Targets ZEB2 and Notch1 but not Jagged2 in Mesothelial Cells

To identify putative direct targets of miR-200c, we searched the TargetScan database (ver. 7.2; <http://www.targetscan.org/>) for miR-200c target sites in mRNA sequences of known repressors of E-cadherin. TargetScan predicted ZEB2, Jagged2, and Notch1 as likely target genes, with their 3' UTR containing 6, 1, and 1 binding sites, respectively, for miR-200c. To investigate targeting of these transcription factors by miR-200c, their miTarget 3' UTR miR target cloned downstream of a firefly luciferase reporter gene, were co-transfected with lenti-miR-200c into mesothelial cells. miR-200c overexpression resulted in a 74.0%, 12.8%, and 46.5% reduction in ZEB2, Notch1, and Jagged2 3' UTR-luciferase activity, respectively (**Figures 7A–7C**). To determine whether miR-200c regulated ZEB2, Notch1, and Jagged2 through direct interaction with its seed region within the 3' UTR region of these target genes, luciferase reporter pEZX-MT06 vectors containing 3' UTR miR targets (wild type) or their mutants were co-transfected with miR-200c mimic oligonucleotides or negative control (Scramble) into mesothelial cells. miR-200c mimic decreased luciferase activity for wild-type 3' UTR of ZEB2 and Notch1 (44.30% and 20.50% reduction, respectively), whereas mutations in the seed regions attenuated this decrease (**Figures 7D and 7E**). miR-200c mimic showed similar reduction in luciferase activity in cells transfected with wild-type and mutant 3' UTR of Jagged2, suggesting that the interaction of miR-200c with Jagged2 in mesothelial cells was indirect (**Figure 7F**).

DISCUSSION

Little is known about the expression and pathophysiological role of miRs in peritoneal mesothelial cells during PD. In this study, we

demonstrated a significant reduction in peritoneal miR-29a, miR-192, miR-200b, miR-200c, miR-324, and miR-377 expression in the murine PD model, when the peritoneal membrane was exposed to conventional glucose-based PD fluids. Peritoneal miR-200c expression was reduced by 80% after 3 days of PD, whereas reduced expression of the other miRs (by 30%–45%) occurred later, but all within 10 days of commencing PD fluid exposure. miR-200b and miR-200c are members of the miR-200 family, and both have been shown to preserve E-cadherin expression in Madin-Darby canine kidney (MDCK) cells, murine mammary gland epithelial cells, and human breast cancer cells.^{18,27,28} These miRs have been postulated to target various transcription factors related to EMT and mediators of fibrosis. The present report is the first to demonstrate the role of miR-200c in peritoneal fibrosis, and the results suggest that glucose-based PD fluid initiates fibrogenesis through repression of miR-200c. In our study, although we did not detect significant peritoneal membrane thickening after 1 month of PD, marked differences were noted in PD-fluid-treated mice, with an increased number of elongated α -smooth-muscle-actin-positive cells in the collagen-rich submesothelium and increased fibronectin expression in the mesothelium, the latter indicating that mesothelial cells play an important role in peritoneal fibrogenesis. That peritoneal membrane thickening was not apparent after 1 month of PD is in agreement with published data, although other studies have shown that peritoneal thickening can be induced after 1 month.^{31,32} The mouse strain, volume of PD fluid injected, and frequency of administration per day may account for such differences. It is possible that a longer PD duration would further exacerbate fibrotic processes and induce peritoneal membrane thickening.

For the studies focusing on miR-200c, we used two sources of peritoneal mesothelial cells: primary peritoneal mesothelial cells isolated from overnight dialysate effluent from PD patients and Met-5A cells, which are normal mesothelial cells transfected with the pRSV-T plasmid. Met-5A cells are often used in studies related to PD, and the data obtained from Met-5A cells showed concordance with that from primary mesothelial cells.^{33,34} Our results showed that glucose-based PD fluid and TGF- β 1 reduced miR-200c expression in cultured mesothelial cells and that this was accompanied by down-regulation of E-cadherin and decorin and induction of EMT transcription factors and α -smooth muscle actin, fibronectin, collagen I, and collagen III. Co-incubation with decorin attenuated the repression of miR-200c by TGF- β 1. Decorin is a dermatan sulfate proteoglycan that possesses anti-fibrotic properties through sequestration of TGF- β 1, thus inhibiting its biological activity.^{30,35} Decorin is the

Figure 3. Effect of Dianeal® or TGF- β 1 on Gene Expression of miR-200c and Mediators of EMT and Fibrosis in Mesothelial Cells

Real-time PCR results showing (A) the effect of Dianeal® and (B) exogenous TGF- β 1 on miR-200c expression. (A) The effect of spent dialysate from three different PD patients (P1, P2, and P3) on miR-200c expression in dialysate-derived human peritoneal mesothelial cells ($n = 3$). "Combined" represents the mean value \pm SD of P1, P2, and P3. (B) Co-incubation of Met-5A cells with TGF- β 1 (10 ng/mL) and decorin (300 ng/mL) prevented the suppression of miR-200c by TGF- β 1. Experiments were performed in duplicate at least three times, and data were analyzed by ANOVA. The effect of TGF- β 1 (10 ng/mL) on gene expression of (C) E-cadherin, (D) SNAIL, (E) fibroblast-specific protein-1 (FSP-1), (F) α -smooth muscle actin (α -SMA), (G) fibronectin, (H) collagen I, and (I) decorin in Met-5A cells. Experiments were performed in triplicate and repeated at least three times, and data were analyzed with the Mann-Whitney test. (J) Cell morphology of primary peritoneal mesothelial cells incubated with SFM or TGF- β 1 (10 ng/mL) for 48 h.




Figure 4. The Effect of TGF-β1 on Protein Expression of Mediators of EMT and Fibrosis in Mesothelial Cells

(A) Representative Western blots showing the effect of TGF-β1 on E-cadherin, SNAIL, FSP-1, α-smooth muscle actin (α-SMA), fibronectin, and collagen I expression in Met-5A cells. The intensity of the bands for (B) E-cadherin, (C) SNAIL, (D) FSP-1, (E) α-SMA, (F) fibronectin, and (G) collagen I was normalized to β-actin and expressed as relative expression compared to serum-free medium (SFM). (H) The effect of TGF-β1 on decorin secretion in mesothelial cells. *p < 0.05, **p < 0.01, and ***p < 0.001, with versus without TGF-β1. Data analyzed by using ANOVA.




Figure 5. miR-200c Overexpression in Mesothelial Cells and Mediators of EMT and Fibrosis


(A) Human peritoneal mesothelial cells (HPMCs) and (B) Met-5A cells showing miR-200c expression in non-transfected mesothelial cells ("C") or mesothelial cells transfected with scrambled control hairpin (lenti-scramble) or lentivector-based miR precursor construct expressing miR-200c (lenti-miR-200c) by real-time PCR. *** $p < 0.001$, lenti-miR-200c versus C or lenti-scramble. Data analyzed by ANOVA. Representative Westerns showing the effect of miR-200c overexpression on (C) E-cadherin, ZEB2, Notch1, Jagged2, and SNAIL expression in HPMCs and on (D) fibronectin and collagen I expression in Met-5A cells following incubation with SFM or TGF- β 1 (10 ng/mL) after 24 h. The intensity of the bands was normalized to β -actin and expressed as arbitrary densitometric scan (DU) (Table 1).

transcription markers or fibronectin. In contrast, cells with miR-200c overexpression showed less induction of ZEB2, Notch1, Jagged2, SNAIL, fibronectin, collagen I, and collagen III by TGF- β 1. miR-200c overexpression was associated with reduced suppression of gene expression, but not protein expression of E-cadherin by TGF- β 1. Possible explanations regarding the discrepancy between gene and protein expression of E-cadherin include variation in the duration of experiment, so that the effect of miR-200c becomes statistically significant, and also additional mechanisms that modulate the protein level of E-cadherin in addition to its gene expression. The impact of miR-200c on cellular changes following TGF- β 1 exposure was not mediated by decorin, as miR-200c overexpression had no effect on decorin expression.

Using luciferase reporter assays, we further demonstrated that miR-200c directly targeted the 3' UTR of ZEB2 and Notch1, whereas the effect of miR-200c on the 3' UTR of Jagged2 appeared to be indirect. The data on ZEB2 is consistent with previous findings in MDCK cells stimulated with TGF- β 1 and in murine NMuMG mammary epithelial cells.^{27,28} It has been suggested that miRs with perfect or near-perfect complementarity can induce mRNA degradation, whereas imperfect complementarity is associated with translational repression.²⁸ In this respect, the perfect complementarity between the seed sequence of miR-200c and the 3' UTR of ZEB2 suggests the former mechanism of regulating gene expression. The ability of miR-200c to target E-cadherin repressors ZEB1 and -2 is well established in a number of epithelial-derived cancer cell types.^{18–20,38} Independent researchers have also identified Notch1 as a target of miR-200c in pancreatic cancer cells and endothelial cells,^{20,39} Notch with its ligand Jagged has been reported to promote EMT through

predominant proteoglycan secreted by cultured human peritoneal mesothelial cells.³⁶ Reduced peritoneal decorin expression has been observed in patients on long-term PD,³⁷ and dialysate decorin level declines with increasing time on PD (unpublished data).

TGF- β 1 induced mesenchymal markers and transcription factors of EMT in mesothelial cells after 24 h of exposure, but phenotypic changes were evident only after 48 h, showing a loss of epithelial morphology, cell detachment, and acquisition of a fibroblastic appearance. In the absence of TGF- β 1, lentivector-mediated miR-200c overexpression did not affect constitutive expression of EMT



(legend on next page)

Table 1. Effect of miR-200c Overexpression on Mediators of EMT and Fibrosis in Mesothelial Cells

Marker	SFM			TGF-β1		
	Non-transfected Mesothelial Cells	Mesothelial Cells Transfected with Lenti-Scramble	Mesothelial Cells Transfected with Lenti-miR-200c	Non-transfected Mesothelial Cells	Mesothelial Cells Transfected with Lenti-Scramble	Mesothelial Cells Transfected with Lenti-miR-200c
E-cadherin	1.01 ± 0.02	0.96 ± 0.07	1.09 ± 0.07	0.45 ± 0.07***	0.35 ± 0.06**	0.42 ± 0.04***
ZEB2	1.06 ± 0.07	0.99 ± 0.1	1.08 ± 0.06	2.57 ± 0.18****	2.8 ± 0.21****	0.72 ± 0.06
Notch1	ND	ND	ND	2.30 ± 0.12****	2.20 ± 0.15****	1.06 ± 0.07
Jagged2	ND	ND	ND	1.06 ± 0.06****	0.97 ± 0.02****	0.18 ± 0.9
SNAIL	ND	ND	ND	0.27 ± 0.03****	0.28 ± 0.04****	0.03 ± 0.03
Fibronectin	1.0 ± 0.01	1.07 ± 0.09	0.96 ± 0.04	2.87 ± 0.14****	2.33 ± 0.17****	0.94 ± 0.13
Collagen I	1.37 ± 0.19###	1.26 ± 0.15###	0.06 ± 0.06	3.17 ± 0.18****	3.17 ± 0.18****	ND

Data expressed as mean ± SEM from 3 separate experiments. ND, not detected. ***p < 0.001, SFM versus TGF-β1 for the same transfection; ###p < 0.01, non-transfected or lenti-scramble versus lenti-miR-200c for the same stimulation. Data analyzed by ANOVA.

SNAIL2-induced repression of E-cadherin,⁴⁰ whereas ZEB expression follows SNAIL activation and is associated with repression of epithelial markers and polarity genes, and activation of mesenchymal genes.⁴¹ Our results showed that repression of ZEB2 by miR-200c was accompanied by downregulation of fibronectin, collagen, SNAIL, Jagged2, and Notch1 expression in mesothelial cells. How miR-200c regulates these changes and whether our findings can be extrapolated to experimental models of PD warrant further investigation. Outside the context of peritoneal fibrosis, there is evidence that fibronectin 1 mRNA is a target of miR-200c in breast and endometrial cancer cell lines.⁴² Fibronectin is one of the first matrix proteins to be deposited during tissue fibrosis, and it serves as a provisional scaffold for the deposition of other extracellular matrix components, such as collagen and laminin.⁴³ Reducing fibronectin deposition by peritoneal mesothelial cells may have therapeutic implications on peritoneal submesothelial thickening commonly observed in patients on PD.

Other investigators have reported miR-200c expression in dialysate-derived mesothelial cells, which decreased with increasing time on PD,⁴⁴ but in another study, investigators using a hybridization array found low signal intensity of miR-200c expression in omentum-derived mesothelial cells that did not change upon stimulation with TGF-β1.¹² The discrepant findings may be related to the source of mesothelial cells used in the experiments, the dose of TGF-β1, and the methods used to detect miRs. miR-200a, a member of the miR-200 family, has also been detected in mesothelial cells and negatively regulates TGF-β1-induced EMT by targeting ZEB1 and -2.⁴⁵ In addition to mesothelial cells, other cell types, such as fibroblasts and endothelial cells, may also participate in peritoneal fibrosis during PD through myofibroblast activation and EMT,

respectively. The role of miRs in these processes requires further investigation.

To summarize, we have identified six miRs that demonstrate a significant preventive effect on EMT and fibrogenesis in peritoneal mesothelial cells subjected to chemical and metabolic injury inflicted by PD fluids that is mediated through TGF-β1. The expression of these miRs is markedly reduced within 1 week of commencing PD in the animal model tested, with the most profound reduction observed with miR-200c, which targets ZEB2 and Notch1 in mesothelial cells. Mechanistic studies showed that reduction of miR-200c was associated with decreased E-cadherin and decorin expression, and increased expression of EMT transcription factors and mesenchymal markers. The data provides evidence that loss of miR-200c is an important contributor to peritoneal fibrosis in PD, and results from miR-200c overexpression experiments suggest that replenishment of miR-200c could be a means to tackle this common PD complication which leads to adverse patient outcomes, but for which there is a total lack of therapeutic options.


MATERIALS AND METHODS

Reagents, Chemicals, and Assay Kits

Tissue culture flasks were purchased from Falcon (Becton-Dickinson, Gene Company, Hong Kong), and culture medium (Medium-199) and supplements, Trizol reagent, Lipofectamine, LTX reagent, and PLUS reagent were purchased from Life Technologies (Thermo Fisher Scientific, Hong Kong). Locked nucleic acid (LNA) probes were purchased from Exiqon (Genetimes ExCell International Holdings, Hong Kong). Digoxigenin (DIG) labeling kits and nitro blue tetrazolium chloride/5-bromo-4-chloro-3-indolyl-phosphate (NBT/BCIP) solution were purchased from Roche Diagnostics (Hong

Figure 6. Effect of miR-200c Overexpression on Gene Expression of Transcription Factors Related to EMT, Fibronectin, Collagen I, and Collagen III

Non-transfected Met-5A cells (C) or Met-5A cells transfected with lenti-scramble (S) or lenti-miR-200c (M) were incubated with serum-free medium (SFM) or TGF-β1 (10 ng/mL) for 24 h, and gene expression of (A) E-cadherin, (B) ZEB2, (C) Notch1, (D) Jagged2, (E) SNAIL, (F) fibronectin, (G) collagen I, and (H) collagen III were determined by real-time PCR. **p < 0.01 and ***p < 0.001, SFM versus TGF-β1 for the same transfection; §p < 0.05, §§p < 0.01, and §§§p < 0.001, non-transfected cells versus lenti-miR-200c; ###p < 0.01 and ####p < 0.001, lenti-scramble versus lenti-miR-200c. Data analyzed by using ANOVA.



(legend on next page)

Kong). Pre-hybridization buffer, PerfectHyb Plus Hybridization buffer, mouse anti-human fibronectin, mouse anti-human α -smooth muscle actin, and mouse anti-human β -actin antibodies were purchased from Sigma Aldrich Co. (Tin Hang Technology, Hong Kong). Met-5A cells were purchased from the American Type Culture Collection (Tin Hang Technology, Hong Kong). Lentivector-based miR precursor constructs expressing miR-200c (PMIRH200cPA-1) and scrambled control hairpin (PMIRH000PA-1), the pPACKH1 lentivector packaging kit, PureFection reagent, PEG-it Virus Precipitation Solution, Global UltraRapid Titer kits, and Transdux reagent were purchased from System Biosciences (Genetix ExCell International Holdings, Hong Kong). miTarget 3' UTR miR target clones for ZEB2 (HmiT054256-MT01), Notch1 (HmiT011873-MT01), and Jagged2 (HmiT067341-MT01), pEZX-MT06 vector, Luc-Pair miR Luciferase Assay kits, and the Dual Luciferase Reporter Assay System were purchased from GeneCopoeia (ATCG, Hong Kong). For mutation studies, wild-type and mutant 3' UTR miR target clones for ZEB2, Notch1, and Jagged2 were purchased from Beijing Genomics Institute (Hong Kong). Hsa-miR-200c-3p mimic and negative control (Scramble) were purchased from GenePharma (Shanghai, China). Taqman MicroRNA Reverse Transcription kits, Taqman microRNA assay, and Taqman probes for E-cadherin (Hs01023894_m1), ZEB2 (Hs00207691_m1), Notch1 (Hs01062011_m1), Jagged2 (Hs00171432_m1), SNAIL (Hs00195591_m1), FSP-1 (Hs00243202_m1), α -smooth muscle actin (Hs00426835_g1), fibronectin (Hs01549976_m1), collagen I (Hs00164004_m1), collagen III (Hs00943809_m1), and decorin (Hs00754870_s1) were purchased from Applied Biosystems (Thermo Fisher Scientific, Hong Kong). RNeasy mini kits were purchased from QIAGEN (Hong Kong). DC Protein Assay kits were purchased from Bio-Rad Pacific (Hong Kong). Goat-anti-human collagen I and collagen III were purchased from Southern Biotech (ATCG, Hong Kong). TGF- β 1 and decorin were purchased from R&D Systems (ATCG, Hong Kong). Rabbit anti-human SNAIL and rabbit anti-human Jagged2 antibodies were purchased from Cell Signaling Technology (Hong Kong). Rabbit anti-human FSP-1, mouse anti-human E-cadherin, and rabbit anti-human ZEB2 antibodies were purchased from Novus Biologicals (Gene Company, Hong Kong). Glucose-based (Dianeal®) PD fluid was purchased from Baxter Healthcare (Hong Kong).

Animal Studies

All animal experiments were approved by the Institutional Committee on the Use of Live Animals in Teaching and Research. Eight-week-old male C57BL/6N mice were purchased from the Laboratory

Animal Unit and housed in a specific pathogen-free animal facility at the University of Hong Kong. Mice were kept under normal housing conditions in a 12-h night and day cycle. Water and standard chow were available *ad libitum*. At 8 weeks of age, the mice were randomized to receive either sterile PBS or 4.25% glucose-based PD fluid (Dianeal® 2 mL) twice daily by intra-peritoneal injection for 1, 3, 5, 10, 15, and 30 days ($n = 3-6$ per time point per group), after which time, the peritoneal membrane was excised for paraffin embedding, *in situ* hybridization, and total RNA extraction.

Peritoneal membrane histology obtained from PBS- and PD-fluid-treated mice was assessed by H&E, Masson's trichrome, and cytochemical staining for α -smooth muscle actin and fibronectin. Peritoneal membrane thickening was determined at six different locations along the length of each specimen using Axiovision software (Carl Zeiss Far East, Hong Kong).

LNA *In Situ* Hybridization

Paraffin-embedded peritoneal specimens (5 μ m) from control and PD-fluid-treated mice were used to determine miR-200c expression, using LNA *in situ* hybridization with a DIG-labeled LNA probe complementary to the miR-200c mature miR sequence. Briefly, deparaffinized sections were permeabilized by incubation with proteinase K (15 μ g/mL) for 2 min, washed three times with diethyl pyrocarbonate (DEPC)-treated PBS, and acetylated for 10 min in 0.1 M triethanolamine (pH 8), containing 0.25% acetic anhydride. Sections were washed three times with DEPC-treated PBS, incubated in pre-hybridization buffer for 1 h at 42°C, and hybridized in PerfectHyb Plus Hybridization buffer containing DIG-labeled LNA probes (0.5 μ M) overnight at 42°C. Sections were subsequently washed twice with 2 \times saline sodium citrate (SSC) buffer for 30 min at 42°C, twice with 1 \times SSC buffer for 10 min at 42°C, and twice with 0.5 \times SSC buffer for 10 min at 42°C. Sections were blocked with 3% fetal bovine serum (FBS) in 0.1 M Tris (pH 7.5), containing, 0.15 M NaCl and 0.1% Tween-20 (NTT) for 1 h and incubated with anti-DIG antibody (dilution 1:500) in NTT overnight at 4°C. The slides were washed three times with NTT followed by three washes for 5 min each with 0.1 M Tris (pH 9.5), containing 50 mM MgCl₂, 0.1 M NaCl, and 0.2% Tween-20, and the sections were developed by incubation with NBT/BCIP solution over 3 days.

Collection of Spent PD Fluid

Overnight PD fluid (1.35% glucose-based Dianeal®) were collected from three stable PD patients who had been free of peritonitis for

Figure 7. Luciferase Reporter Assays Identify Target Genes of miR-200c

miR-200c overexpression reduced luciferase activity of (A) ZEB2, (B) Jagged2, and (C) Notch1. C: non-transfected Met-5A cells, S: Met-5A cells transfected with lenti-scramble, M: Met-5A transfected with lenti-miR-200c. * $p < 0.05$ and *** $p < 0.001$, non-transfected cells versus cells transfected with lenti-miR-200c; ### $p < 0.001$, lenti-scramble versus lenti-miR-200c. Red line highlights the position of the miR-200c seed sequences in the 3' UTR of ZEB2, Notch1, and Jagged2. Luciferase reporter assays showing direct binding of miR-200c to the wild-type but not mutant sequences within the 3' UTR seed regions of (D) ZEB2 and (E) Notch1. (F) Luciferase reporter assay demonstrated that miR-200c did not directly bind to the seed region of Jagged2. Each study was performed in triplicate and repeated at least three times. The blue line in panels (D–F) highlights the putative miR-200c binding sites that were deleted in mutant-type target gene 3' UTR for ZEB2, Notch1, and Jagged2. * $p < 0.05$ and *** $p < 0.001$, scrambled versus miR-200c mimic; # $p < 0.05$ and ## $p < 0.01$, wild-type versus mutant for miR-200c mimic. Data analyzed by using ANOVA.

at least 6 months. PD fluid was centrifuged at 3,000 rpm at 4°C for 10 min, and the supernatant was collected, filter sterilized, and stored at -80°C until analysis. The collection of PD fluid was approved by the Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster. All patients were recruited from the PD Unit at Queen Mary Hospital.

Cell Culture

Immortalized human mesothelial cells (Met-5A cells) were cultured in Medium-199 supplemented with penicillin (100 IU/mL), streptomycin (100 mg/mL), insulin (5 µg/mL), transferrin (5 µg/mL), and 10% FBS (Complete Medium). Primary human peritoneal mesothelial cells were isolated from overnight 1.36%-glucose-based Dianeal® PD fluid obtained from patients who had been on PD for less than 2 years. Briefly, cells were centrifuged at 1,000 × g for 5 min, and pelleted cells seeded in 25 cm² flasks in Complete Medium. Only primary mesothelial cells with a polygonal, cobblestone appearance of the second passage were used, and mixed cell populations or senescent mesothelial cells were discarded. Similarities between primary peritoneal mesothelial cells with a polygonal appearance, and Met-5A cells with regard to morphology and function have been demonstrated previously.^{33,34} All studies were conducted in confluent, growth arrested human peritoneal mesothelial cells and Met-5A cells. To determine the effect of 1.36% glucose-based Dianeal® PD fluid on miR-200c expression, confluent human peritoneal mesothelial cells were incubated with serum-free medium (SFM) or autologous PD fluid for 24 h. To determine the effect of TGF-β1 on cell morphology, confluent human peritoneal mesothelial cells were incubated with SFM or TGF-β1 (10 ng/mL) for up to 48 h. To determine the effect of TGF-β1 on the expression of miR-200c and mediators of EMT and fibrosis, Met-5A cells were incubated with SFM or TGF-β1 (0.1–10.0 ng/mL) for 24 h after which the culture medium was decanted and cells lysed with Trizol or 25 mM sodium acetate (pH 6.0), containing 4 M urea, 1% Triton X-100, and a cocktail of proteinase inhibitors for the isolation of total RNA or protein respectively.^{8,46,47} Met-5A and human peritoneal mesothelial cells will be referred to as mesothelial cells in this study.

Overexpression of miR-200c in Mesothelial Cells

Lenti-miR-200c and lenti-scramble vectors were generated by System Biosciences and contained an expression module for the copGFP fluorescent marker gene that enabled the detection of cells with positive transfection or transduction. Met-5A cells were transfected with lenti-miR-200c or lenti-scramble vectors (15 µg) using Lipofectamine LTX Reagent and PLUS Reagent, according to the manufacturer's instructions. Human peritoneal mesothelial cells were transduced with lenti-miR-200c and lenti-scramble vectors packaged in pseudovirus particles (1 × 10⁴ lentivirus transducing units) according to the manufacturer's instructions. After 5 h, the medium was replaced with fresh Complete Medium. Transfection efficiency was monitored by detection of the GFP reporter gene by fluorescence microscopy and by real-time PCR. Cell lysates or total RNA were collected after 2 days.

miR-200c Target Analysis and Luciferase Reporter Assay

Using TargetScan version 7.2 (<http://www.targetscan.org/>), ZEB2 (target gene accession: NM_001171653.1), Notch1 (NM_017617.5), and Jagged2 (NM_002226.4) were predicted to be target genes of miR-200c. Mesothelial cells were transfected with miTarget ZEB2, Notch1, or Jagged2 3' UTR constructs cloned downstream of the secreted Gaussia luciferase gene (1 µg), with or without lenti-scramble or lenti-miR-200c (1.4 µg), using Lipofectamine 2000 according to the manufacturer's instructions. These miTarget constructs also contained the secreted embryonic alkaline phosphatase gene driven by a cytomegalovirus (CMV) promoter used as the internal control. After 5 h, the medium was replaced with Complete Medium and the cells incubated for 48 h, after which the supernatants were collected for analysis of secreted Gaussia luciferase and secreted embryonic alkaline phosphatase activities using the Luc-Pair miR Luciferase Assay kit according to the manufacturer's instructions.

In a separate study, wild-type and mutant 3' UTR of ZEB2, Notch1, and Jagged2 were cloned into the *EcoRI* and *XhoI* sites downstream of *Renilla* luciferase in a CMV-driven firefly/*Renilla* Duo-Luciferase reporter vector (pEZX-MT06). To investigate whether miR-200c directly targeted the 3' UTR of ZEB2, Notch1, and Jagged2, 85%–95% confluent mesothelial cells were co-transfected with pEZX-MT06 reporter plasmids (0.4 µg) and either miR-200c mimic or negative control (0.4 µg) using Lipofectamine 2000 for 48 h, after which time, luciferase activities were determined using the Dual Luciferase Reporter Assay System. For ZEB2 mutant, seed sequences at positions 455–462, 813–820, 898–905, 1,029–1,036, and 1,363–1,369 were deleted. For Notch1 and Jagged2 mutants, seed sequences at positions 736–743 and 777–783, respectively, were deleted.

Real-Time PCR

Total RNA from peritoneal membrane specimens or mesothelial cells was extracted by using Trizol reagent according to the manufacturer's instructions, and dissolved in RNase-free water. The yield and purity of the total RNA was determined with the Nanodrop system. To determine the expression of miR-29a, -192, -200b, -200c, -324, and -377, total RNA (50 ng) was reverse transcribed with Taqman MicroRNA Reverse Transcription kits, according to the manufacturer's instructions followed by real-time PCR, using Taqman microRNA assays. miRs expression was normalized to endogenous U6 snRNA expression.

To determine mRNA expression of mediators of EMT and fibrosis in mesothelial cells, total mRNA was extracted with RNeasy mini kits according to the manufacturer's instructions. Two micrograms of total RNA was reverse transcribed into cDNA with Moloney murine leukemia virus (MMLV) reverse transcriptase, by using the random hexamer method.⁴⁸ The effect of TGF-β1 and miR-200c overexpression on E-cadherin, SNAIL, Notch1, ZEB2, Jagged2, FSP-1, α-smooth muscle actin, fibronectin, collagen I, collagen III, and decorin was assessed by quantitative real-time PCR on a Lightcycler 480 II real-time PCR system (Roche Diagnostics, DKSH Hong Kong, Hong Kong). All

samples were analyzed in triplicate, and mRNA expression of the aforementioned mediators of fibrosis was calculated using the delta Ct ($2^{-\Delta\Delta Ct}$) method, normalized to GAPDH.

Western Blot

Whole-cell lysates (20 μ g total protein) were electrophoresed under denaturing conditions on 8% polyacrylamide gels to determine fibronectin and collagen I expression; on 12% polyacrylamide gels to determine E-cadherin, ZEB2, Notch1, Jagged2, vimentin, α -smooth muscle actin, and β -actin expression; and on 15% polyacrylamide gels to determine SNAIL expression. Proteins were transferred onto nitrocellulose membranes and immunoblotted with the relevant primary antibodies followed by the addition of secondary antibodies, as previously described.^{48,49} Bands were visualized with enhanced chemiluminescence (ECL), semi-quantitated by densitometry using ImageJ (NIH, USA), and expressed as arbitrary densitometric units (DU). All proteins of interest were normalized to β -actin.

Measurement of Decorin in Culture Supernatant

Decorin secretion in mesothelial cells cultured under control and experimental conditions was measured in culture supernatant using decorin DuoSet ELISA (R&D Systems, Bio-Techne Hong Kong, Hong Kong). The detection range was 31.2–2,000 pg/mL.

Statistical Analysis

Data are presented as means \pm SEM of results from at least three separate experiments, unless otherwise mentioned. Data were analyzed by ANOVA with Bonferroni's multiple-comparison post hoc test, Mann-Whitney or Kolmogorov-Smirnov test as appropriate, using GraphPad Prism version 6 for Windows, (GraphPad Software, San Diego, CA, USA). A two-tailed $p < 0.05$ was considered statistically significant.

AUTHOR CONTRIBUTIONS

Conceptualization: S.Y. and T.M.C.; Methodology: S.Y., J.Y.S.C., and A.C.P.T.; Investigation: J.Y.S.C., M.K.M.C., C.C.Y.C., A.C.P.T., and K.F.C.; Validation: J.Y.S.C., A.C.P.T., and S.Y.; Formal Analysis: S.Y., J.Y.S.C., and A.C.P.T.; Writing – Original Draft: S.Y. and J.Y.S.C.; Writing – Review & Editing: S.Y. and T.M.C.; Funding Acquisition: S.Y. and T.M.C.; and Supervision: S.Y. and T.M.C.

CONFLICTS OF INTEREST

The authors declare no competing interests.

ACKNOWLEDGMENTS

This study was supported by the RGC General Research Fund (HKU 7848/12M and 17105915), HKU Seed Fund for Basic Research (201211159081), the Stanley Ho Alumni Challenge, the Department of Medicine Academic Activities Fund, and kind donations from Mr. C.S. Yung, Mr. S. Ho, and the Hui Hoy & Chow Sin Lan Charity Fund and the family of Mr. Hui Ming. S.Y. is supported by the Endowment Fund established for the Yu Chiu Kwong Professorship in Medicine awarded to T.M.C., and the Wai Hung Charitable Foundation Limited. We would like to thank all PD patients for partici-

pating in this study. We are most grateful to the nurses at the Renal Unit, Queen Mary Hospital, Miss Vicky Ho, and Miss Cindy Tang for coordinating the collection of PD fluid. We would also like to thank Dr. Mandy K.M. Kam and Dr. Ming Shum Yip for technical assistance.

REFERENCES

1. Cho, Y., and Johnson, D.W. (2014). Peritoneal dialysis-related peritonitis: towards improving evidence, practices, and outcomes. *Am. J. Kidney Dis.* 64, 278–289.
2. Williams, J.D., Craig, K.J., Topley, N., Von Ruhland, C., Fallon, M., Newman, G.R., Mackenzie, R.K., and Williams, G.T.; Peritoneal Biopsy Study Group (2002). Morphologic changes in the peritoneal membrane of patients with renal disease. *J. Am. Soc. Nephrol.* 13, 470–479.
3. Davies, S.J., Bryan, J., Phillips, L., and Russell, G.I. (1996). Longitudinal changes in peritoneal kinetics: the effects of peritoneal dialysis and peritonitis. *Nephrol. Dial. Transplant.* 11, 498–506.
4. Korte, M.R., Sampimon, D.E., Betjes, M.G., and Krediet, R.T. (2011). Encapsulating peritoneal sclerosis: the state of affairs. *Nat. Rev. Nephrol.* 7, 528–538.
5. Honda, K., Nitta, K., Horita, S., Yumura, W., and Nihei, H. (1996). Morphological changes in the peritoneal vasculature of patients on CAPD with ultrafiltration failure. *Nephron* 72, 171–176.
6. Yáñez-Mó, M., Lara-Pezzi, E., Selgas, R., Ramírez-Huesca, M., Domínguez-Jiménez, C., Jiménez-Heffernan, J.A., Aguilera, A., Sánchez-Tomero, J.A., Bajo, M.A., Alvarez, V., et al. (2003). Peritoneal dialysis and epithelial-to-mesenchymal transition of mesothelial cells. *N. Engl. J. Med.* 348, 403–413.
7. Ha, H., Yu, M.R., and Lee, H.B. (2001). High glucose-induced PKC activation mediates TGF- β 1 and fibronectin synthesis by peritoneal mesothelial cells. *Kidney Int.* 59, 463–470.
8. Chan, T.M., Leung, J.K., Tsang, R.C., Liu, Z.H., Li, L.S., and Yung, S. (2003). Emodin ameliorates glucose-induced matrix synthesis in human peritoneal mesothelial cells. *Kidney Int.* 64, 519–533.
9. Yang, A.H., Chen, J.Y., and Lin, J.K. (2003). Myofibroblastic conversion of mesothelial cells. *Kidney Int.* 63, 1530–1539.
10. Bartel, D.P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116, 281–297.
11. Kozomara, A., and Griffiths-Jones, S. (2014). miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res.* 42, D68–D73.
12. Lopez-Anton, M., Lambie, M., Lopez-Cabrera, M., Schmitt, C.P., Ruiz-Carpio, V., Bartosova, M., Schaefer, B., Davies, S., Stone, T., Jenkins, R., et al. (2017). miR-21 Promotes Fibrogenesis in Peritoneal Dialysis. *Am. J. Pathol.* 187, 1537–1550.
13. Gao, Q., Xu, L., Yang, Q., and Guan, T.J. (2019). MicroRNA-21 contributes to high glucose-induced fibrosis in peritoneal mesothelial cells in rat models by activation of the Ras-MAPK signaling pathway via Sprouty-1. *J. Cell. Physiol.* 234, 5915–5925.
14. Shang, J., He, Q., Chen, Y., Yu, D., Sun, L., Cheng, G., Liu, D., Xiao, J., and Zhao, Z. (2019). miR-15a-5p suppresses inflammation and fibrosis of peritoneal mesothelial cells induced by peritoneal dialysis via targeting VEGFA. *J. Cell. Physiol.* 234, 9746–9755.
15. Zhou, Q., Yang, M., Lan, H., and Yu, X. (2013). miR-30a negatively regulates TGF- β 1-induced epithelial-mesenchymal transition and peritoneal fibrosis by targeting Snail. *Am. J. Pathol.* 183, 808–819.
16. Xiao, L., Zhou, X., Liu, F., Hu, C., Zhu, X., Luo, Y., Wang, M., Xu, X., Yang, S., Kanwar, Y.S., and Sun, L. (2015). MicroRNA-129-5p modulates epithelial-to-mesenchymal transition by targeting SIP1 and SOX4 during peritoneal dialysis. *Lab. Invest.* 95, 817–832.
17. Li, X., Liu, H., Sun, L., Zhou, X., Yuan, X., Chen, Y., Liu, F., Liu, Y., and Xiao, L. (2019). MicroRNA-302c modulates peritoneal dialysis-associated fibrosis by targeting connective tissue growth factor. *J. Cell. Mol. Med.* 23, 2372–2383.
18. Bracken, C.P., Gregory, P.A., Kolesnikoff, N., Bert, A.G., Wang, J., Shannon, M.F., and Goodall, G.J. (2008). A double-negative feedback loop between ZEB1-SIP1 and the microRNA-200 family regulates epithelial-mesenchymal transition. *Cancer Res.* 68, 7846–7854.

19. Hur, K., Toiyama, Y., Takahashi, M., Balaguer, F., Nagasaka, T., Koike, J., Hemmi, H., Koi, M., Boland, C.R., and Goel, A. (2013). MicroRNA-200c modulates epithelial-to-mesenchymal transition (EMT) in human colorectal cancer metastasis. *Gut* 62, 1315–1326.
20. Huang, C.C., Lin, C.M., Huang, Y.J., Wei, L., Ting, L.L., Kuo, C.C., Hsu, C., Chiou, J.F., Wu, A.T.H., and Lee, W.H. (2017). Garcinol downregulates Notch1 signaling via modulating miR-200c and suppresses oncogenic properties of PANC-1 cancer stem-like cells. *Biotechnol. Appl. Biochem.* 64, 165–173.
21. Kwon, J.J., Willy, J.A., Quirin, K.A., Wek, R.C., Korc, M., Yin, X.M., and Kota, J. (2016). Novel role of miR-29a in pancreatic cancer autophagy and its therapeutic potential. *Oncotarget* 7, 71635–71650.
22. Ye, C., Hu, Y., and Wang, J. (2019). MicroRNA-377 Targets Zinc Finger E-box-Binding Homeobox 2 to Inhibit Cell Proliferation and Invasion of Cervical Cancer. *Oncol. Res.* 27, 183–192.
23. Cao, L., Xie, B., Yang, X., Liang, H., Jiang, X., Zhang, D., Xue, P., Chen, D., and Shao, Z. (2015). MiR-324-5p Suppresses Hepatocellular Carcinoma Cell Invasion by Counteracting ECM Degradation through Post-Transcriptionally Downregulating ETS1 and SP1. *PLoS ONE* 10, e0133074.
24. Du, B., Ma, L.M., Huang, M.B., Zhou, H., Huang, H.L., Shao, P., Chen, Y.Q., and Qu, L.H. (2010). High glucose down-regulates miR-29a to increase collagen IV production in HK-2 cells. *FEBS Lett.* 584, 811–816.
25. Wang, B., Komers, R., Carew, R., Winbanks, C.E., Xu, B., Herman-Edelstein, M., Koh, P., Thomas, M., Jandeleit-Dahm, K., Gregorevic, P., et al. (2012). Suppression of microRNA-29 expression by TGF- β 1 promotes collagen expression and renal fibrosis. *J. Am. Soc. Nephrol.* 23, 252–265.
26. Wang, B., Herman-Edelstein, M., Koh, P., Burns, W., Jandeleit-Dahm, K., Watson, A., Saleem, M., Goodall, G.J., Twigg, S.M., Cooper, M.E., and Kanharidis, P. (2010). E-cadherin expression is regulated by miR-192/215 by a mechanism that is independent of the profibrotic effects of transforming growth factor-beta. *Diabetes* 59, 1794–1802.
27. Gregory, P.A., Bert, A.G., Paterson, E.L., Barry, S.C., Tsykin, A., Farshid, G., Vadas, M.A., Khew-Goodall, Y., and Goodall, G.J. (2008). The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1. *Nat. Cell Biol.* 10, 593–601.
28. Korpala, M., Lee, E.S., Hu, G., and Kang, Y. (2008). The miR-200 family inhibits epithelial-mesenchymal transition and cancer cell migration by direct targeting of E-cadherin transcriptional repressors ZEB1 and ZEB2. *J. Biol. Chem.* 283, 14910–14914.
29. Loureiro, J., Aguilera, A., Selgas, R., Sandoval, P., Albar-Vizcaino, P., Pérez-Lozano, M.L., Ruiz-Carpio, V., Majano, P.L., Lamas, S., Rodríguez-Pascual, F., et al. (2011). Blocking TGF- β 1 protects the peritoneal membrane from dialysate-induced damage. *J. Am. Soc. Nephrol.* 22, 1682–1695.
30. Border, W.A., Noble, N.A., Yamamoto, T., Harper, J.R., Yamaguchi, Yu., Pierschbacher, M.D., and Ruoslahti, E. (1992). Natural inhibitor of transforming growth factor-beta protects against scarring in experimental kidney disease. *Nature* 360, 361–364.
31. Morishita, Y., Yoshizawa, H., Watanabe, M., Imai, R., Imai, T., Hirahara, I., Akimoto, T., Ookawara, S., Muto, S., and Nagata, D. (2016). MicroRNA expression profiling in peritoneal fibrosis. *Transl. Res.* 169, 47–66.
32. Yu, J.W., Duan, W.J., Huang, X.R., Meng, X.M., Yu, X.Q., and Lan, H.Y. (2014). MicroRNA-29b inhibits peritoneal fibrosis in a mouse model of peritoneal dialysis. *Lab. Invest.* 94, 978–990.
33. Rampino, T., Cancarini, G., Gregorini, M., Guallini, P., Maggio, M., Ranghino, A., Soccio, G., and Dal Canton, A. (2001). Hepatocyte growth factor/scatter factor released during peritonitis is active on mesothelial cells. *Am. J. Pathol.* 159, 1275–1285.
34. Bidmon, B., Endemann, M., Arbeiter, K., Ruffingshofer, D., Regele, H., Herkner, K., Eickelberg, O., and Aufricht, C. (2004). Overexpression of HSP-72 confers cytoprotection in experimental peritoneal dialysis. *Kidney Int.* 66, 2300–2307.
35. Yamaguchi, Y., Mann, D.M., and Ruoslahti, E. (1990). Negative regulation of transforming growth factor-beta by the proteoglycan decorin. *Nature* 346, 281–284.
36. Yung, S., Thomas, G.J., Stylianou, E., Williams, J.D., Coles, G.A., and Davies, M. (1995). Source of peritoneal proteoglycans. Human peritoneal mesothelial cells synthesize and secrete mainly small dermatan sulfate proteoglycans. *Am. J. Pathol.* 146, 520–529.
37. Osada, S., Hamada, C., Shimaoka, T., Kaneko, K., Horikoshi, S., and Tomino, Y. (2009). Alterations in proteoglycan components and histopathology of the peritoneum in uraemic and peritoneal dialysis (PD) patients. *Nephrol. Dial. Transplant.* 24, 3504–3512.
38. Li, X., Roslan, S., Johnstone, C.N., Wright, J.A., Bracken, C.P., Anderson, M., Bert, A.G., Selth, L.A., Anderson, R.L., Goodall, G.J., et al. (2014). MiR-200 can repress breast cancer metastasis through ZEB1-independent but moesin-dependent pathways. *Oncogene* 33, 4077–4088.
39. Zhang, Y., Quan, Q., and Jin, X. (2017). miR-200c serves an important role in H5V endothelial cells in high glucose by targeting Notch1. *Mol. Med. Rep.* 16, 2149–2155.
40. Leong, K.G., Niessen, K., Kulic, I., Raouf, A., Eaves, C., Pollet, I., and Karsan, A. (2007). Jagged1-mediated Notch activation induces epithelial-to-mesenchymal transition through Slug-induced repression of E-cadherin. *J. Exp. Med.* 204, 2935–2948.
41. Lamouille, S., Xu, J., and Derynck, R. (2014). Molecular mechanisms of epithelial-mesenchymal transition. *Nat. Rev. Mol. Cell Biol.* 15, 178–196.
42. Howe, E.N., Cochrane, D.R., and Richer, J.K. (2011). Targets of miR-200c mediate suppression of cell motility and anoikis resistance. *Breast Cancer Res.* 13, R45.
43. Dixon, A.J., Burns, J., Dunnill, M.S., and McGee, J.O. (1980). Distribution of fibronectin in normal and diseased human kidneys. *J. Clin. Pathol.* 33, 1021–1028.
44. Zhang, L., Liu, F., Peng, Y., Sun, L., and Chen, G. (2013). Changes in expression of four molecular marker proteins and one microRNA in mesothelial cells of the peritoneal dialysate effluent fluid of peritoneal dialysis patients. *Exp. Ther. Med.* 6, 1189–1193.
45. Guo, R., Hao, G., Bao, Y., Xiao, J., Zhan, X., Shi, X., Luo, L., Zhou, J., Chen, Q., and Wei, X. (2018). MiR-200a negatively regulates TGF- β 1-induced epithelial-mesenchymal transition of peritoneal mesothelial cells by targeting ZEB1/2 expression. *Am. J. Physiol. Renal Physiol.* 314, F1087–F1095.
46. Yung, S., Ng, C.Y., Au, K.Y., Cheung, K.F., Zhang, Q., Zhang, C., Yap, D.Y., Chau, M.K., and Chan, T.M. (2017). Binding of anti-dsDNA antibodies to proximal tubular epithelial cells contributes to renal tubulointerstitial inflammation. *Clin. Sci. (Lond.)* 131, 49–67.
47. Yung, S., Ng, C.Y., Ho, S.K., Cheung, K.F., Chan, K.W., Zhang, Q., Chau, M.K., and Chan, T.M. (2015). Anti-dsDNA antibody induces soluble fibronectin secretion by proximal renal tubular epithelial cells and downstream increase of TGF- β 1 and collagen synthesis. *J. Autoimmun.* 58, 111–122.
48. Yung, S., Cheung, K.F., Zhang, Q., and Chan, T.M. (2010). Anti-dsDNA antibodies bind to mesangial annexin II in lupus nephritis. *J. Am. Soc. Nephrol.* 21, 1912–1927.
49. Yung, S., Zhang, Q., Zhang, C.Z., Chan, K.W., Lui, S.L., and Chan, T.M. (2009). Anti-DNA antibody induction of protein kinase C phosphorylation and fibronectin synthesis in human and murine lupus and the effect of mycophenolic acid. *Arthritis Rheum.* 60, 2071–2082.