

Research Paper

†Authors contributed equally

Cite this article: Zhang Y, Xu S-Q, Liu W, Fung WK, Zhou J-Y (2020). A robust test for X-chromosome genetic association accounting for X-chromosome inactivation and imprinting. *Genetics Research* **102**, e2, 1–12. <https://doi.org/10.1017/S0016672320000026>

Received: 2 October 2019

Revised: 17 January 2020

Accepted: 26 February 2020

Keywords:

association test; imprinting effects; inactivation; X chromosome

Author for correspondence:

Professor Ji-Yuan Zhou,
E-mail: zhoujiyuan5460@hotmail.com

A robust test for X-chromosome genetic association accounting for X-chromosome inactivation and imprinting

Yu Zhang^{1,†}, Si-Qi Xu^{1,2,†}, Wei Liu¹, Wing Kam Fung² and Ji-Yuan Zhou¹ 

¹State Key Laboratory of Organ Failure Research, Ministry of Education, and Guangdong Provincial Key Laboratory of Tropical Disease Research, Department of Biostatistics, School of Public Health, Southern Medical University, Guangzhou, China and ²Department of Statistics and Actuarial Science, The University of Hong Kong, Hong Kong, China

Abstract

The X chromosome is known to play an important role in many sex-specific diseases. However, only a few single-nucleotide polymorphisms on the X chromosome have been found to be associated with diseases. Compared to the autosomes, conducting association tests on the X chromosome is more intractable due to the difference in the number of X chromosomes between females and males. On the other hand, X-chromosome inactivation takes place in female mammals, which is a phenomenon in which the expression of one copy of two X chromosomes in females is silenced in order to achieve the same gene expression level as that in males. In addition, imprinting effects may be related to certain diseases. Currently, there are some existing approaches taking X-chromosome inactivation into account when testing for associations on the X chromosome. However, none of them allows for imprinting effects. Therefore, in this paper, we propose a robust test, Z_{XCI} , which accounts for both X-chromosome inactivation and imprinting effects without requiring specifying the genetic models in advance. Simulation studies are conducted in order to investigate the validity and performance of Z_{XCI} under various scenarios of different parameter values. The simulation results show that Z_{XCI} controls the type I error rate well when there is no association. Furthermore, with regards to power, Z_{XCI} is robust in all of the situations considered and generally outperforms most of the existing methods in the presence of imprinting effects, especially under complete imprinting effects.

1. Introduction

The X chromosome has been found to play an important role in many complex diseases (Ober *et al.*, 2008; Wise *et al.*, 2013). However, the development of methods for detecting associations with X-linked markers has lagged behind that for autosomal markers due to the complexity of the inheritance patterns of the X chromosome (Wise *et al.*, 2013; Schurz *et al.*, 2019). One primary characteristic of the X chromosome in mammals is that females have two copies of the X chromosome while males only have one, which increases the difficulty of X-linked association studies (Clayton, 2009; Ziegler, 2009; Loley *et al.*, 2011). In addition, the phenomenon of X-chromosome inactivation (XCI) in females may constitute a risk factor for diseases, which is defined as the expression silencing of one of the two copies of the X chromosome in females. Thus, the X-chromosome gene dosage in female XX cells equals that in male XY cells, namely dosage compensation (Chow *et al.*, 2005; Payer & Lee, 2008; Pessia *et al.*, 2012). As such, the genetic effect of homozygous females can be regarded as the same as that of hemizygous males under XCI. It has been reported that most of the genes on the X chromosome are subject to XCI, while only about 15% of X-linked genes escape from inactivation (XCI-E) (Carrel & Willard, 2005). Random X-chromosome inactivation (XCI-R) is the general process of XCI by which one of the two copies of the X chromosome in each cell is randomly inactivated. But the XCI patterns in some females may become skewed from that of the XCI-R in an age- and tissue-dependent manner, and the same allele can be inactivated in more than 75% of cells in some cases (Migeon, 1998; Minks *et al.*, 2008; Starmer & Magnuson, 2009; Wang *et al.*, 2014), which is denoted by XCI-S for convenience.

At present, there are some association tests available for single-nucleotide polymorphisms (SNPs) on the X chromosome. Zheng *et al.* (2007) proposed six methods for testing associations on the X chromosome by combining the genetic effects in females and males. Among them, the allele-based tests Z_A and Z_{mfA} require the assumption of a Hardy–Weinberg equilibrium (HWE), while the genotype-based methods Z_C and Z_{mfG} are robust to departures from a HWE. Furthermore, note that all four methods mentioned

© The Author(s), 2020. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

above rely on the assumption that females and males have the same risk alleles. Thus, two other methods (\tilde{Z}_{mfA} and \tilde{Z}_{mfG}) were developed and are applicable to the situation in which females and males have different risk alleles. On the other hand, the six methods of Zheng *et al.* (2007) only consider the information on XCI-E and do not take account of XCI, which may lead to loss of power if XCI is present. Clayton (2008) was the first to suggest that XCI should be considered in X-chromosome association studies. Clayton's methods (T_A and T_{AD}) are equivalent to the score tests of generalized linear models accounting for XCI-R and give the same codes for homozygous females and hemizygous males. When the allele frequencies of the same allele differ between the sexes, the test statistics T_A^s , T_{AD}^s and S_A , stratified by sex, have been proposed by Loley *et al.* (2011) and König *et al.* (2014). In addition, a software toolset XWAS (Gao *et al.*, 2015) includes four tests (FM_{01} , FM_{02} , FM_F and FM_S) based on logistic regressions. However, those approaches only consider XCI-R and ignore XCI-S. In order to simultaneously incorporate three biological patterns on the X chromosome (XCI-E, XCI-R and XCI-S), Wang *et al.* (2014) developed a maximum likelihood ratio method. However, this method is time-consuming because it is a permutation-based procedure for obtaining an empirical P -value. Meanwhile, Chen *et al.* (2017) proposed a robust method (*Xcat*) based on a generalized genetic model with the approximate P -value being easily obtained. Recently, Wang *et al.* (2019) proposed a robust test, Z_{max} , by taking account of different dosage compensation patterns, which requires neither the assumption of a HWE nor the specification of underlying genetic models.

Imprinting is an epigenetic phenomenon that results in the differential expression of paternal and maternal alleles (Falls *et al.*, 1999). Researchers have found evidence for the existence of imprinting effects on some diseases, such as Angelman, Beckwith–Wiedemann and Prader–Willi syndromes (Falls *et al.*, 1999; Dong *et al.*, 2005; Ziegler & König, 2006; Wallace *et al.*, 2010). On the other hand, it is likely that imprinted genes on the X chromosome are crucial to some diseases, such as Turner's syndrome (Donnelly *et al.*, 2002; Loesch *et al.*, 2005). For some sex-specific diseases, such as autism, alleles on the paternal chromosome seem to be preferentially expressed, which is likely to explain why females are always less susceptible than males (Skuse, 2000). Imprinting is generally detected through testing for parent-of-origin effects (Hager *et al.*, 2008). Thus, we use the term 'parent-of-origin effects' instead of 'imprinting effects' in the following sections. However, there is no method available for taking parent-of-origin effects into account when conducting association tests on the X chromosome.

Therefore, in this paper, we propose a robust method, Z_{XCI} , which is an extension of *Xcat* to the generalized linear model simultaneously accounting for imprinting and three biological patterns (XCI-E, XCI-R and XCI-S) into X-chromosome association tests without the need to specify the genetic models on the X chromosome. We investigate the performance of the proposed method and compare it with several existing tests through extensive simulation studies. Simulation results show that the proposed method controls the size well under all of the scenarios considered when there is no association. Moreover, with regards to power, Z_{XCI} is robust in all of the situations considered and generally outperforms most of the existing methods in the presence of imprinting effects, especially under complete imprinting effects.

2. Materials and methods

For a candidate SNP on the X chromosome with the mutant allele A and the normal allele a , there are four ordered genotypes for female offspring: a/a , a/A , A/a and A/A , where the left (right) allele of the slash is paternal (maternal). To distinguish the parent of origin of the mutant allele A in heterozygous female offspring, the information on their parental genotypes is required. With regards to male offspring, there are only two kinds of genotypes, a and A , which are maternal. Thus, we do not need to collect their parental genotypes. Assume that G_{f1} and G_{f2} are the numbers of allele A on the paternal and maternal X chromosomes in female offspring, respectively, and G_m is the number of allele A on the X chromosome in male offspring. The values of G_{f1} , G_{f2} and G_m for different genotypes in the offspring generation are shown in Table 1. The disease status of an individual (female or male) in the offspring generation is denoted by Y with 1 (0) representing being affected (unaffected). In this paper, an affected daughter together with her parents is called a case–parent trio and an unaffected daughter together with her parents is considered as a control–parent trio (Deng & Chen, 2001; Li *et al.*, 2016). Table 2 gives the genotype counts for the female offspring, where n_f is the total number of daughter–parent trios consisting of r_f case–parent trios and s_f control–parent trios. The genotype counts for the male offspring are also listed in Table 2, where n_m is the total number of males including r_m cases and s_m controls. As such, there are $n_r = r_f + r_m$ cases and $n_s = s_f + s_m$ controls in total. Therefore, the sample size is $N = n_r + n_s = n_f + n_m$. Let ϕ_{f0} , ϕ_{f01} , ϕ_{f10} and ϕ_{f2} be the penetrances of genotypes a/a , a/A , A/a and A/A in female offspring, respectively, and let ϕ_{m0} and ϕ_{m1} be the penetrances of genotypes a and A in male offspring, respectively. To test the association between the disease status Y and the SNP under study, we make the following two assumptions, just like *Xcat* (Chen *et al.*, 2017): (1) in the presence of association between the disease and the SNP, the generalized genetic model is assumed to hold in female offspring with ordered penetrances, either increasing ($\phi_{f0} \leq \phi_{f01}$, $\phi_{f10} \leq \phi_{f2}$) or decreasing ($\phi_{f0} \geq \phi_{f01}$, $\phi_{f10} \geq \phi_{f2}$); and (2) the mutant allele in female offspring is the same as that in male offspring.

A logistic regression model is proposed to describe the association between the disease and the SNP in female offspring:

$$\begin{aligned} \text{Logit}(\Pr(Y = 1 | G_{f1}, G_{f2}, \mathbf{X}_f)) \\ = \beta_{f0} + \beta_{f1}G_{f1} + \beta_{f2}G_{f2} + \beta_{f3}G_{f1}G_{f2}, \quad (1) \\ + \mathbf{b}_f^T \mathbf{X}_f \end{aligned}$$

where β_{f0} is the intercept, β_{f1} , β_{f2} and β_{f3} are the respective regression coefficients for G_{f1} , G_{f2} and the interaction term $G_{f1}G_{f2}$, \mathbf{X}_f is a vector of covariates and \mathbf{b}_f is a vector of the regression coefficients for \mathbf{X}_f . The estimates of these coefficients can be obtained with the iteratively reweighted least squares method (Wood, 2006) using the *glm* function in R language (<http://www.r-project.org>). The null hypothesis of no association between the disease and the SNP in female offspring is $H_{f0}: \beta_{f1} = \beta_{f2} = \beta_{f3} = 0$. If at least one of these equations is not satisfied, then the association exists, which indicates the alternative hypothesis (H_{f1}). Logit ($\Pr(Y = 1 | G_{f1}, G_{f2}, \mathbf{X}_f)$) outcomes for different genotypes in female offspring are presented in the fourth column of Table 1. Thus, under H_{f1} , the parent-of-origin effects at the SNP locus can be

Table 1. Values of G_{f1} , G_{f2} and G_m for different genotypes in the offspring generation.

Genotype	Female			Male	
	G_{f1}	G_{f2}	$Logit(Pr(Y=1 G_{f1}, G_{f2}, \mathbf{X}_f))$	Genotype	G_m
a/a	0	0	$\beta_{f0} + \mathbf{b}_f^T \mathbf{X}_f$	a	0
a/A	0	1	$\beta_{f0} + \beta_{f2} + \mathbf{b}_f^T \mathbf{X}_f$	A	1
A/a	1	0	$\beta_{f0} + \beta_{f1} + \mathbf{b}_f^T \mathbf{X}_f$		
A/A	1	1	$\beta_{f0} + \beta_{f1} + \beta_{f2} + \beta_{f3} + \mathbf{b}_f^T \mathbf{X}_f$		

expressed by:

$$\frac{Logit(Pr(Y = 1|a/A, \mathbf{X}_f)) - Logit(Pr(Y = 1|A/a, \mathbf{X}_f))}{Logit(Pr(Y = 1|A/A, \mathbf{X}_f)) - Logit(Pr(Y = 1|a/a, \mathbf{X}_f))} = \frac{\beta_{f2} - \beta_{f1}}{\beta_{f1} + \beta_{f2} + \beta_{f3}}, \tag{2}$$

when \mathbf{X}_f is fixed at the same level. For example, $\beta_{f1} = \beta_{f2}$ represents no parent-of-origin effects, while $\beta_{f2} = \beta_{f3} = 0$ denotes complete maternal parent-of-origin effect and $\beta_{f1} = \beta_{f3} = 0$ indicates complete paternal parent-of-origin effect. Moreover, we can use

$$\gamma = 2 \times \frac{[Logit(Pr(Y = 1|a/A, \mathbf{X}_f)) + Logit(Pr(Y = 1|A/a, \mathbf{X}_f)) - 2 \times Logit(Pr(Y = 1|a/a, \mathbf{X}_f))]/2}{Logit(Pr(Y = 1|A/A, \mathbf{X}_f)) - Logit(Pr(Y = 1|a/a, \mathbf{X}_f))} = \frac{\beta_{f1} + \beta_{f2}}{\beta_{f1} + \beta_{f2} + \beta_{f3}}, \tag{3}$$

to measure the degree of inactivation under XCI in a similar way to Wang *et al.* (2019). On the other hand, the difference between β_{f3} and 0 can be interpreted as the deviation of the genetic model from the additive one under XCI-E. To be specific, Table 3 gives the explanations of the regression coefficients for several situations of XCI and XCI-E under no parent-of-origin effects ($\beta_{f1} = \beta_{f2} = \beta$). $\beta_{f1} = \beta_{f2} = -\beta_{f3}$ means XCI-S with $\gamma = 2$ representing 100% of the cells having the mutant allele active or a dominant model under XCI-E. $\beta_{f1} = \beta_{f2} = \beta$ and $\beta_{f3} = -\frac{2}{3}\beta$ stand for XCI-S with $\gamma = 1.5$, where 75% of the cells have the mutant allele active. $\beta_{f1} = \beta_{f2} \neq 0$ and $\beta_{f3} = 0$ correspond to XCI-R with $\gamma = 1$ or an additive model under XCI-E. $\beta_{f1} = \beta_{f2} = \beta$ and $\beta_{f3} = 2\beta$ imply XCI-S with $\gamma = 0.5$, where 25% of the cells have the mutant allele active. $\beta_{f1} = \beta_{f2} = 0$ and $\beta_{f3} \neq 0$ indicate XCI-S with $\gamma = 0$ representing that 100% of the cells have the normal allele active or a recessive model under XCI-E. However, in the presence of parent-of-origin effects, the explanation of the regression coefficients is more complicated, since parent-of-origin effects may contribute to the XCI. For example, $\beta_{f1} = 0.5$ and $\beta_{f2} = \beta_{f3} = 0$ are indicative of the complete maternal parent-of-origin effect, whereas γ is obtained to be 1 (suggesting XCI-R) in this case. Therefore, XCI-R may be also caused by the complete maternal parent-of-origin effect.

Recall that when the disease is associated with the SNP, the generalized genetic model with ordered penetrances is assumed

to hold in female offspring. As such, we have

$$Logit(Pr(Y = 1|a/a, \mathbf{X}_f)) \leq Logit(Pr(Y = 1|a/A, \mathbf{X}_f)) \leq Logit(Pr(Y = 1|A/A, \mathbf{X}_f))$$

and

$$Logit(Pr(Y = 1|a/a, \mathbf{X}_f)) \leq Logit(Pr(Y = 1|A/a, \mathbf{X}_f)) \leq Logit(Pr(Y = 1|A/A, \mathbf{X}_f))$$

which are equivalent to $0 \leq \beta_{f1} \leq \beta_{f1} + \beta_{f2} + \beta_{f3}$ and $0 \leq \beta_{f2} \leq \beta_{f1} + \beta_{f2} + \beta_{f3}$, respectively, with at least one inequality being strict. Adding these two inequalities together, we get $0 \leq \beta_{f1} + \beta_{f2} \leq 2(\beta_{f1} + \beta_{f2} + \beta_{f3})$ and thus $\beta_{f1} + \beta_{f2} + 2\beta_{f3} \geq 0$. Therefore, the alternative hypothesis becomes $H_{f1} : \beta_{f1} \geq 0, \beta_{f2} \geq 0, \beta_{f1} + \beta_{f2} + 2\beta_{f3} \geq 0$, with at least one inequality being strict, which can be expressed in matrix form as follows:

$$\mathbf{C}\boldsymbol{\beta}_f = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{pmatrix} \begin{pmatrix} \beta_{f1} \\ \beta_{f2} \\ \beta_{f3} \end{pmatrix} \geq \mathbf{0}, \tag{4}$$

where $\mathbf{C} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{pmatrix}$, $\boldsymbol{\beta}_f = \begin{pmatrix} \beta_{f1} \\ \beta_{f2} \\ \beta_{f3} \end{pmatrix}$, and $\mathbf{0}$ is a vector with all of the elements being 0. To test for the association, we first consider the following test statistics:

$$\mathbf{Z} = (Z_1, Z_2, Z_3)^T = (\hat{\mathbf{C}}\hat{\boldsymbol{\beta}}_f)^T \mathbf{I}^{-\frac{1}{2}} \hat{\boldsymbol{\beta}}_f, \tag{5}$$

where $\hat{\boldsymbol{\beta}}_f = (\hat{\beta}_{f1}, \hat{\beta}_{f2}, \hat{\beta}_{f3})^T$ with $\hat{\beta}_{f1}$, $\hat{\beta}_{f2}$ and $\hat{\beta}_{f3}$ being the maximum likelihood estimates of β_{f1} , β_{f2} and β_{f3} , respectively. $\hat{\mathbf{I}}$ is the empirical Fisher's information matrix (Wood, 2006).

Under the null hypothesis of no association, Z_1 , Z_2 and Z_3 are independent of one another and asymptotically have standard normal distributions. Note that $\mathbf{C}\boldsymbol{\beta}_f \geq \mathbf{0}$ leads to $\mathbf{Z} \geq \mathbf{0}$ under H_{f1} , and we thus only calculate the right-sided P -values for Z_1 , Z_2 and Z_3 , respectively. Then, we combine them using the Fisher's method (Fisher, 1954). Thus, the test statistic for female offspring can be constructed as:

$$Q_f^R = -2 \ln(\Phi(-Z_1)\Phi(-Z_2)\Phi(-Z_3)), \tag{6}$$

where $\Phi(\cdot)$ is the cumulative distribution function of the standard normal distribution. Under the null hypothesis, Q_f^R has an asymptotic χ^2 distribution with degrees of freedom (df) being 6. As

Table 2. Genotype counts for the single-nucleotide polymorphism on the X chromosome stratified by sex in the offspring generation.

Group	Female					Male		
	<i>a/a</i>	<i>a/A</i>	<i>A/a</i>	<i>A/A</i>	Total	<i>a</i>	<i>A</i>	Total
Case	$r_{a/a}$	$r_{a/A}$	$r_{A/a}$	$r_{A/A}$	r_f	r_a	r_A	r_m
Control	$s_{a/a}$	$s_{a/A}$	$s_{A/a}$	$s_{A/A}$	s_f	s_a	s_A	s_m
Total	$n_{a/a}$	$n_{a/A}$	$n_{A/a}$	$n_{A/A}$	n_f	n_a	n_A	n_m

Table 3. Explanation of the regression coefficients under no parent-of-origin effects.

Coefficients	γ	XCI	XCI-E
$\beta_{f1} = \beta_{f2} = -\beta_{f3}$	2	XCI-S (100% of the cells have the mutant allele active)	Dominant model
$\beta_{f1} = \beta_{f2} = \beta, \beta_{f3} = -\frac{2}{3}\beta$	1.5	XCI-S (75% of the cells have the mutant allele active)	—
$\beta_{f1} = \beta_{f2} \neq 0, \beta_{f3} = 0$	1	XCI-R (random XCI)	Additive model
$\beta_{f1} = \beta_{f2} = \beta, \beta_{f3} = 2\beta$	0.5	XCI-S (25% of the cells have the mutant allele active)	—
$\beta_{f1} = \beta_{f2} = 0, \beta_{f3} \neq 0$	0	XCI-S (100% of the cells have the normal allele active)	Recessive model

XCI = X-chromosome inactivation.

such, the P -value of Q_f^R is $P_f^R = 1 - \chi_6^2(Q_f^R)$, where $\chi_6^2(\cdot)$ is the cumulative distribution function of the χ^2 distribution with df being 6.

For male offspring, we model the relationship between the disease and the SNP using a logistic regression as:

$$\text{Logit}(\text{Pr}(Y = 1 | G_m, \mathbf{X}_m)) = \beta_{m0} + \beta_m G_m + \mathbf{b}_m^T \mathbf{X}_m, \quad (7)$$

where β_{m0} is the intercept, β_m is the regression coefficient for G_m , \mathbf{X}_m is a vector of covariates and \mathbf{b}_m is a vector of the regression coefficients for \mathbf{X}_m . When there is no association between the disease and the SNP, the null hypothesis for male offspring is H_{m0} : $\beta_m = 0$. Then, the test statistic for male offspring is

$$Z_m = \frac{\hat{\beta}_m}{S_{\hat{\beta}_m}}, \quad (8)$$

where $\hat{\beta}_m$ is the maximum likelihood estimate of β_m and $S_{\hat{\beta}_m}$ is the standard error of $\hat{\beta}_m$. Z_m follows a standard normal distribution under H_{m0} . When there are no covariates, Eq. (8) can be simplified to

$$Z_m = \frac{n_m^{1/2}(r_m s_a - s_m r_a)}{(n_a n_A r_m s_m)^{1/2}} \quad (9)$$

as in Zheng *et al.* (2007) and Chen *et al.* (2017).

For combining the test statistics of female and male offspring, we need to turn the P -value for female offspring (P_f^R) into a Z -score, which is $Z_f^R = -\Phi^{-1}(P_f^R)$. Then, under the assumption that the mutant allele in female offspring is the same as that in male offspring, the combined test statistics Z^R can be constructed as follows:

$$Z^R = \sqrt{\frac{n_f}{n_f + n_m}} Z_f^R + \sqrt{\frac{n_m}{n_f + n_m}} Z_m, \quad (10)$$

where Z_f^R and Z_m are weighted by their respective proportions of the sample size. Under the overall null hypothesis that there is no association between the disease and the SNP in both female and male offspring ($H_0: \mathbf{C}\boldsymbol{\beta}_f = \mathbf{0}$ and $\beta_m = 0$), Z^R is asymptotically distributed as $N(0, 1)$. Since the mutant allele is assumed to be A , with the overall one-sided alternative hypothesis $H_1: \mathbf{C}\boldsymbol{\beta}_f \geq \mathbf{0}$ (with at least one inequality being strict) or $\beta_m > 0$, we only need to calculate the right-sided P -value of Z^R when the mutant allele is known in advance.

So far, we have only considered the situation when the mutant allele is A . When the mutant allele is a , the overall alternative hypothesis turns to be $H_1: \mathbf{C}\boldsymbol{\beta}_f \leq \mathbf{0}$ (with at least one inequality being strict) or $\beta_m < 0$. Therefore, the corresponding test statistic for female offspring is $Q_f^L = -2 \ln(\Phi(Z_1)\Phi(Z_2)\Phi(Z_3))$, which combines the left-sided P -values of Z_1, Z_2 and Z_3 , and the P -value of Q_f^L is $P_f^L = 1 - \chi_6^2(Q_f^L)$. Again, we combine the transformed Z -score ($Z_f^L = -\Phi^{-1}(P_f^L)$) for female offspring and Z_m for male offspring to obtain the overall test statistic as:

$$Z^L = \sqrt{\frac{n_f}{n_f + n_m}} Z_f^L + \sqrt{\frac{n_m}{n_f + n_m}} (-Z_m). \quad (11)$$

Z^L is asymptotically distributed as $N(0, 1)$ under the overall null hypothesis. With this H_1 , just like Z^R , only the right-sided P -value of Z^L is needed when the mutant allele is known to be a in advance.

However, we generally have no information on the mutant allele before conducting the association studies. In this case, we propose the test statistic as:

$$Z_{XCII} = \max(Z^L, Z^R). \quad (12)$$

Although Z^L and Z^R are obviously dependent on each other, note that the components of $\mathbf{Z}_t = (Z_1, Z_2, Z_3, Z_m)^T$ are independent of each other, and the functions $-Z^L$ and Z^R of \mathbf{Z}_t are non-decreasing functions. Thus, the P -value of Z_{XCII} can be

approximately bounded by

$$2\xi - \xi^2 \leq Pr(Z_{XCI} > z) \leq 2\xi, \tag{13}$$

where $\xi = 1 - \Phi(z)$ according to Owen (2009) and Esary *et al.* (1967). Therefore, we can simply get the approximated *P*-value of Z_{XCI} by 2ξ .

3. Simulation study

3.1. Settings

We conduct a simulation study to investigate the size and power of the proposed Z_{XCI} method and compare it with the existing ones. Notice that in Zheng *et al.* (2007), and \tilde{Z}_{mfA} and \tilde{Z}_{mfG} are less powerful than the other four test statistics (Z_A , Z_C , Z_{mfA} and Z_{mfG}) under the assumption that the mutant allele in females is the same as that in males. Thus, in this simulation study, \tilde{Z}_{mfA} and \tilde{Z}_{mfG} are excluded. T_A^s and FM_S are also excluded because they are asymptotically equivalent to Z_C (Loley *et al.*, 2011) and Z_{mfG} (Zheng *et al.*, 2007; Gao *et al.*, 2015; Wang *et al.*, 2019), respectively. On the other hand, the permutation-based method in Wang *et al.* (2014) is excluded due to the intensive computations involved. Finally, we choose 14 methods (Z_{XCI} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C , Z_{mfG} , T_A , T_{AD} , T_{AD}^s , FM_{01} , FM_F , Z_{mfA} and Z_A) for the comparison. The references for the selected methods are listed in Table S1.

Note that most of the methods we compare do not consider the covariates, such as $Xcat$, S_A , Z_C , Z_{mfG} , T_A , T_{AD} , Z_{mfA} and Z_A . Thus, we do not include any covariate for simplicity in this simulation study and directly generate the genotype counts in Table 2. Let p_f and p_m denote the frequencies of the mutant allele *A* for females and males in the parental generation, respectively. Under random mating, the genotype frequencies of *a/a*, *a/A*, *A/a* and *A/A* for female offspring are $g_{f0} = (1 - p_m)(1 - p_f)$, $g_{f01} = (1 - p_m)p_f$, $g_{f10} = p_m(1 - p_f)$ and $g_{f2} = p_m p_f$, respectively, and the genotype frequencies of *a* and *A* for male offspring are $g_{m0} = 1 - p_f$ and $g_{m1} = p_f$, respectively. Note that if random mating holds in the parental generation, HWE holds in the offspring generation only under the assumption that the frequency of the same allele in females and that of males are equal (Puig *et al.*, 2017). On the other hand, we consider the situation where $p_f = p_m = p$ but HWE does not hold in the female offspring. The corresponding frequencies of the four genotypes are $g_{f0} = (1 - p)^2 + \rho p(1 - p)$, $g_{f01} = (1 - p)p(1 - p)$, $g_{f10} = (1 - \rho)p(1 - p)$ and $g_{f2} = p^2 + \rho p(1 - p)$, respectively, when the inbreeding coefficient $\rho \neq 0$. Furthermore, the genotype frequencies for male offspring are $g_{m0} = 1 - p$ and $g_{m1} = p$, respectively.

Note that the relationships among the penetrances and the regression coefficients are $\frac{\phi_{f01}(1-\phi_{f0})}{(1-\phi_{f01})\phi_{f0}} = e^{\beta_{f2}}$, $\frac{\phi_{f10}(1-\phi_{f0})}{(1-\phi_{f10})\phi_{f0}} = e^{\beta_{f1}}$ and $\frac{\phi_{f2}(1-\phi_{f0})}{(1-\phi_{f2})\phi_{f0}} = e^{\beta_{f1}+\beta_{f2}+\beta_{f3}}$ for *a/A*, *A/a* and *A/A*, respectively, for female offspring and $\frac{\phi_{m1}(1-\phi_{m0})}{(1-\phi_{m1})\phi_{m0}} = e^{\beta_m}$ for male offspring. Thus, genotype counts for female offspring in Table 2 can be generated according to a quadrinomial distribution with probabilities $(\frac{g_{f0}\phi_{f0}}{\phi_f}, \frac{g_{f01}\phi_{f01}}{\phi_f}, \frac{g_{f10}\phi_{f10}}{\phi_f}, \frac{g_{f2}\phi_{f2}}{\phi_f})$ for cases and $(\frac{g_{f0}(1-\phi_{f0})}{1-\phi_f}, \frac{g_{f01}(1-\phi_{f01})}{1-\phi_f}, \frac{g_{f10}(1-\phi_{f10})}{1-\phi_f}, \frac{g_{f2}(1-\phi_{f2})}{1-\phi_f})$ for controls, where $\phi_f = g_{f0}\phi_{f0} + g_{f01}\phi_{f01} + g_{f10}\phi_{f10} + g_{f2}\phi_{f2}$ is the disease prevalence of females. Similarly, we can obtain genotype counts for male offspring through a binomial

distribution with probabilities $(\frac{g_{m0}\phi_{m0}}{\phi_m}, \frac{g_{m1}\phi_{m1}}{\phi_m})$ for cases and $(\frac{g_{m0}(1-\phi_{m0})}{1-\phi_m}, \frac{g_{m1}(1-\phi_{m1})}{1-\phi_m})$ for controls, where $\phi_m = g_{m0}\phi_{m0} + g_{m1}\phi_{m1}$ is the disease prevalence of males.

We consider various simulation settings. (p_f, p_m) is taken to be (0.15, 0.25), (0.20, 0.20), (0.25, 0.15), (0.25, 0.35), (0.30, 0.30) and (0.35, 0.25). Then, under random mating, the corresponding allele frequencies for females and males in the offspring generation are (0.20, 0.15), (0.20, 0.20), (0.20, 0.25), (0.30, 0.25), (0.30, 0.30) and (0.30, 0.35), respectively. When $p_f = p_m = p = 0.2$ and 0.3, we set $\rho = -0.05$ and $\rho = 0.05$ for simulating the departure from HWE. ϕ_{f0} and ϕ_{m0} are set to be 0.120. For simulating the size, let all of the other penetrances be 0.120. When XCI exists, we suppose $\phi_{f2} = \phi_{m1} = 0.240$. The values of γ under XCI with different values of ϕ_{f01} and ϕ_{f10} are shown in Table S2. To investigate the power, we first consider the situations where there are both XCI and parent-of-origin effects: (1) $(\phi_{f01}, \phi_{f10}) = (0.120, 0.240)$ (XCI with $\gamma = 1$ and complete maternal parent-of-origin effect); (2) $(\phi_{f01}, \phi_{f10}) = (0.192, 0.216)$ (XCI with $\gamma = 1.499$ and incomplete maternal parent-of-origin effect); (3) $(\phi_{f01}, \phi_{f10}) = (0.144, 0.204)$ (XCI with $\gamma = 1.001$ and incomplete maternal parent-of-origin effect); (4) $(\phi_{f01}, \phi_{f10}) = (0.132, 0.156)$ (XCI with $\gamma = 0.492$ and incomplete maternal parent-of-origin effect); (5) $(\phi_{f01}, \phi_{f10}) = (0.240, 0.120)$ (XCI with $\gamma = 1$ and complete paternal parent-of-origin effect); (6) $(\phi_{f01}, \phi_{f10}) = (0.216, 0.192)$ (XCI with $\gamma = 1.499$ and incomplete paternal parent-of-origin effect); (7) $(\phi_{f01}, \phi_{f10}) = (0.204, 0.144)$ (XCI with $\gamma = 1.001$ and incomplete paternal parent-of-origin effect); and (8) $(\phi_{f01}, \phi_{f10}) = (0.156, 0.132)$ (XCI with $\gamma = 0.492$ and incomplete paternal parent-of-origin effect). Next, we take account of the scenarios where XCI exists but there are no parent-of-origin effects with $\phi_{f01} = \phi_{f10} = \phi$: (1) $\phi = 0.240$ (XCI with $\gamma = 2$); (2) $\phi = 0.204$ (XCI with $\gamma = 1.503$); (3) $\phi = 0.168$ (XCI with $\gamma = 0.935$); (4) $\phi = 0.144$ (XCI with $\gamma = 0.500$); and (5) $\phi = 0.120$ (XCI with $\gamma = 0$). Furthermore, we consider the situation where there is neither XCI nor parent-of-origin effects, which is $(\phi_{f01}, \phi_{f10}, \phi_{f2}, \phi_{m1}) = (0.180, 0.180, 0.240, 0.180)$. The sample size *N* for each replication is selected to be 1000, including $n_r = 500$ cases and $n_s = 500$ controls. To investigate the effect of sex ratio, we fix the sex ratio in the control group as $s_f:s_m = 1:1$, while it varies in the case group as $r_f:r_m = 3:2, 1:1$ and 2:3. We use the significance level $\alpha = 10^{-5}$, and the number of replications is fixed to be 10^6 and 10^4 for estimating the size and power, respectively. The definitions of these parameters and the detailed biological meanings of the situations we consider are provided in Tables S3 and S4, respectively.

3.2. Size

Table 4 gives the estimated sizes of Z_{XCI} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C , Z_{mfG} , T_A , T_{AD} , T_{AD}^s , FM_{01} , FM_F , Z_{mfA} and Z_A under different simulation settings when random mating holds in the parental generation. From Table 4, we can see that Z_{XCI} , Z_{max} , $Xcat$, FM_{02} , Z_C , T_{AD}^s , FM_{01} , FM_F , Z_{mfA} and Z_A generally control the size well, except that some of them produce a slightly conservative size under some situations. The sizes of S_A and Z_{mfG} are inflated when $(p_f, p_m) = (0.35, 0.25)$ and the sex ratio is 3:2, and they stay close to the nominal level 10^{-5} for all of the other situations. T_A and T_{AD} can have inflated size when (p_f, p_m) is equal to (0.25, 0.15) and (0.35, 0.25), which may be caused by the different allele frequencies between females and males in the offspring

Table 4. Estimated size ($\times 10^{-5}$) under random mating at significance level $\alpha = 10^{-5}$ based on 10^6 replicates.^a

p_F	p_M	Sex ratio	Z_{XCII}	Z_{max}	X_{cat}	S_A	FM_{02}	Z_C	Z_{mFG}	T_A	T_{AD}	T_{AD}^S	FM_{01}	FM_F	Z_{mFA}	Z_A	
0.15	0.25	3:2	0.3	0.7	0.8	1.0	0.7	1.2	0.8	0.4	0.7	1.0	0.6	0.5	1.0	0.9	
		1:1	1.0	1.0	1.1	1.1	0.6	1.0	1.4	0.4	0.9	0.9	1.1	0.6	1.0	1.1	
		2:3	0.2	0.8	0.4	0.6	0.5	0.9	0.4	0.4	0.6	0.5	0.6	0.4	0.5	0.8	
0.20	0.20	3:2	0.4	0.8	0.9	1.3	0.5	1.1	1.2	1.0	1.2	1.1	0.9	0.5	1.1	0.8	
		1:1	0.7	0.8	0.6	0.8	0.8	0.8	0.5	0.9	0.8	1.2	0.7	0.5	0.4	0.8	0.8
		2:3	0.5	0.9	0.3	0.8	1.0	0.8	1.0	1.0	1.2	0.9	0.7	0.8	0.5	0.9	0.9
0.25	0.15	3:2	0.3	0.7	0.2	0.8	0.9	0.7	0.8	2.9	1.5	0.6	0.6	0.4	0.5	1.2	
		1:1	0.2	0.5	0.7	1.1	0.3	0.8	0.9	1.5	0.7	0.6	0.7	0.8	0.5	0.6	
		2:3	0.7	0.9	0.9	0.8	0.9	1.1	1.1	2.5	2.1	0.6	0.8	0.6	0.9	1.2	
0.25	0.35	3:2	1.4	1.0	1.1	1.1	0.9	1.2	1.1	0.9	0.9	1.3	0.9	1.1	1.1	1.6	
		1:1	0.5	1.0	1.0	0.8	0.8	0.7	0.9	0.6	0.7	0.9	0.9	0.7	1.4	1.3	
		2:3	0.7	0.7	0.7	0.4	1.0	1.1	0.9	1.2	0.9	0.6	0.4	0.6	0.6	1.2	
0.30	0.30	3:2	1.0	1.0	0.8	1.0	1.2	1.0	1.3	1.0	0.5	0.7	0.8	0.4	0.9	0.7	
		1:1	0.6	0.7	0.3	0.6	0.7	0.4	1.0	1.1	0.2	0.6	0.7	0.2	0.8	0.9	
		2:3	0.7	0.9	0.8	1.2	0.6	0.7	0.8	0.6	0.9	0.2	1.0	0.3	1.1	1.0	
0.35	0.25	3:2	1.2	1.6	1.2	1.7	1.6	0.9	1.7	2.1	1.7	0.4	1.1	0.8	1.1	1.3	
		1:1	1.3	0.9	0.7	1.4	1.2	0.9	1.6	2.1	0.8	0.4	1.2	0.7	1.2	1.2	
		2:3	0.4	0.8	0.5	0.5	1.4	0.7	0.7	1.7	1.1	0.4	0.6	0.5	0.6	1.1	

^aNumbers that are outside of the 95% confidence interval (0.38×10^{-5} , 1.62×10^{-5}) are highlighted in bold.




Fig. 1. Estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} against sex ratio ($r_f:r_m=3:2, 1:1$ and $2:3$) under random mating when there is X-chromosome inactivation with $\gamma=1$ and complete maternal parent-of-origin effects. The simulation is based on 10,000 replicates with $N=1000$, $\phi_{f0}=\phi_{m0}=\phi_{f01}=0.120$ and $\phi_{f10}=\phi_{f2}=\phi_{m1}=0.240$. (a) $p_f=0.15, p_m=0.25$. (b) $p_f=0.20, p_m=0.20$. (c) $p_f=0.25, p_m=0.15$. (d) $p_f=0.25, p_m=0.35$. (e) $p_f=0.30, p_m=0.30$. (f) $p_f=0.35, p_m=0.25$.

generation. However, they have a well-controlled size under the other situations. Table S5 reports the estimated sizes of different methods when $p_F=p_M=p$ but HWE does not hold in female offspring. In addition, Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C , Z_{mfG} , T_A , T_{AD} , T_{AD}^s , FM_{01} , FM_F , Z_{mfA} and Z_A generally control the size well. Z_{mfA} and Z_A can have inflated size when $\rho=0.05$ and $p=0.30$ since the allele-based test relies on the assumption of HWE in females.

3.3. Power

To clearly illustrate the power results, we show the estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} with relatively better performance in Figures 1–6 and Figures S1–S22, and those of T_A , T_{AD} , T_{AD}^s , FM_{01} , FM_F , Z_{mfA} and Z_A with inflated size or lower powers are displayed in Figures S23–S50. Figure 1 gives the estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} against sex ratio under random mating when there is XCI with $\gamma=1$ and complete maternal parent-of-origin effect. It is shown in Figure 1 that Z_{XCII} has the highest power among all seven methods. The powers of Z_{max} , FM_{02} and Z_{mfG} are similar to each other and are generally higher than those of $Xcat$, S_A and Z_C . On the other hand, the powers are influenced by the sex ratio. When the proportion of males in the case group gets larger ($r_f:r_m$ changing from 3:2 to 2:3), the power of Z_{XCII} becomes smaller in Figure 1(a), while it remains nearly unchanged in the other subplots of Figure 1, and the powers of Z_{max} , $Xcat$, FM_{02} , Z_C and Z_{mfG} are almost unchanged in Figure 1(a), while they are larger in the other subplots. However, with the number of males in the case group, S_A is less powerful. It is also found

that all of the methods have higher powers with increasing allele frequency (comparing the first row with the second row). Figure 2 displays the corresponding estimated powers when there is XCI with $\gamma=1.001$ and incomplete maternal parent-of-origin effect. From Figure 2, we can see that the powers of Z_{XCII} , Z_{max} , FM_{02} and Z_{mfG} are very close to each other, which are generally larger than those of $Xcat$, S_A and Z_C . Compared to Figure 1, the effect of the sex ratio on Z_{XCII} is greater as the power of Z_{XCII} increases with larger male proportion in the case group in the second and third columns of Figure 2.

When there are XCI and no parent-of-origin effects under random mating, the estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} with $\gamma=2, 0.935$ and 0 are shown in Figures 3–5, respectively. From Figure 3, Z_{mfG} has the highest power in the first row of Figure 3, while Z_{XCII} is the most powerful in the second row. In fact, the powers of Z_{XCII} , Z_{max} , $Xcat$ and Z_{mfG} are very close to each other, which are larger than those of FM_{02} and Z_C . S_A has relatively good performance in the first row of Figure 3, while it performs worse in the second row. In Figure 4, we find that Z_{XCII} generally has higher power than $Xcat$, S_A and Z_C , although it has less power than Z_{max} , FM_{02} and Z_{mfG} . $Xcat$ is always the most powerful in all of the subplots of Figure 5. In the first row of Figure 5, Z_{XCII} , Z_{max} , FM_{02} and Z_C have similar powers, which perform much better than S_A and Z_{mfG} . In the second row of Figure 5, Z_{XCII} is more powerful than the other five methods, except for $Xcat$. Furthermore, by comparing Figures 3–5, we find that the powers get larger with increasing γ -value. By comparing Figure 1 (complete maternal parent-of-origin effect), Figure 2 (incomplete maternal




Fig. 2. Estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} against sex ratio ($r_f:r_m=3:2, 1:1$ and $2:3$) under random mating when there is X-chromosome inactivation with $\gamma=1.001$ and incomplete maternal parent-of-origin effects. The simulation is based on 10,000 replicates with $N=1000$, $\phi_{r0}=\phi_{m0}=0.120$, $\phi_{r01}=0.144$, $\phi_{r10}=0.204$ and $\phi_{r2}=\phi_{m1}=0.240$. (a) $p_r=0.15$, $p_u=0.25$. (b) $p_r=0.20$, $p_u=0.20$. (c) $p_r=0.25$, $p_u=0.15$. (d) $p_r=0.25$, $p_u=0.35$. (e) $p_r=0.30$, $p_u=0.30$. (f) $p_r=0.35$, $p_u=0.25$.




Fig. 3. Estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} against sex ratio ($r_f:r_m=3:2, 1:1$ and $2:3$) under random mating when there is X-chromosome inactivation with $\gamma=2$ and no parent-of-origin effects. The simulation is based on 10,000 replicates with $N=1000$, $\phi_{r0}=\phi_{m0}=0.120$ and $\phi_{r01}=\phi_{r10}=\phi_{r2}=\phi_{m1}=0.240$. (a) $p_r=0.15$, $p_u=0.25$. (b) $p_r=0.20$, $p_u=0.20$. (c) $p_r=0.25$, $p_u=0.15$. (d) $p_r=0.25$, $p_u=0.35$. (e) $p_r=0.30$, $p_u=0.30$. (f) $p_r=0.35$, $p_u=0.25$.




Fig. 4. Estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} against sex ratio ($r_f:r_m = 3:2, 1:1$ and $2:3$) under random mating when there is X-chromosome inactivation with $\gamma = 0.935$ and no parent-of-origin effects. The simulation is based on 10,000 replicates with $N = 1000$, $\phi_{f0} = \phi_{m0} = 0.120$, $\phi_{f01} = \phi_{f10} = 0.168$ and $\phi_{f2} = \phi_{m1} = 0.240$. (a) $p_f = 0.15, p_u = 0.25$. (b) $p_f = 0.20, p_u = 0.20$. (c) $p_f = 0.25, p_u = 0.15$. (d) $p_f = 0.25, p_u = 0.35$. (e) $p_f = 0.30, p_u = 0.30$. (f) $p_f = 0.35, p_u = 0.25$.




Fig. 5. Estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} against sex ratio ($r_f:r_m = 3:2, 1:1$ and $2:3$) under random mating when there is X-chromosome inactivation with $\gamma = 0$ and no parent-of-origin effects. The simulation is based on 10,000 replicates with $N = 1000$, $\phi_{f0} = \phi_{m0} = \phi_{f01} = \phi_{f10} = 0.120$ and $\phi_{f2} = \phi_{m1} = 0.240$. (a) $p_f = 0.15, p_u = 0.25$. (b) $p_f = 0.20, p_u = 0.20$. (c) $p_f = 0.25, p_u = 0.15$. (d) $p_f = 0.25, p_u = 0.35$. (e) $p_f = 0.30, p_u = 0.30$. (f) $p_f = 0.35, p_u = 0.25$.




Fig. 6. Estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} against sex ratio ($r_f:r_m=3:2, 1:1$ and $2:3$) under random mating when there is neither X-chromosome inactivation nor parent-of-origin effects. The simulation is based on 10,000 replicates with $N=1000$, $\phi_{f0}=\phi_{m0}=0.120$, $\phi_{f01}=\phi_{f10}=\phi_{m1}=\phi_{m10}=0.180$ and $\phi_{f2}=\phi_{m2}=0.240$. (a) $p_f=0.15, p_u=0.25$. (b) $p_f=0.20, p_u=0.20$. (c) $p_f=0.25, p_u=0.15$. (d) $p_f=0.25, p_u=0.35$. (e) $p_f=0.30, p_u=0.30$. (f) $p_f=0.35, p_u=0.25$.

parent-of-origin effect) and Figure 4 (no parent-of-origin effects) with γ being fixed close to 1 (XCI-R), the power of Z_{XCII} becomes smaller and smaller. Figure 6 plots the estimated powers of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} against the sex ratio under random mating when there is neither XCI nor parent-of-origin effects. Z_{XCII} has similar power to $Xcat$ and FM_{02} in most situations. Z_{max} , S_A and Z_{mfG} always outperform the other methods, while the power of Z_C is always the lowest among those methods. The relatively low power of Z_{XCII} is due to no XCI and no parent-of-origin effects.

The power results of Z_{XCII} , Z_{max} , $Xcat$, S_A , FM_{02} , Z_C and Z_{mfG} with $\gamma=1.499$ and 0.492 under random mating and incomplete maternal parent-of-origin effect are given in Figures S1 and S2, respectively. When there are no parent-of-origin effects, Figures S3 and S4 plot the estimated powers under XCI with $\gamma=1.503$ and 0.500 , respectively. The powers of these seven methods under random mating and paternal parent-of-origin effects are shown in Figures S5–S8. The results are similar to those under maternal parent-of-origin effects, except that the powers of Z_{XCII} seem to be more strongly affected by the difference between p_f and p_m under paternal parent-of-origin effects. For example, the difference in power between Figure S5(c) and Figure S5(a) is much larger than that between Figure 1(c) and Figure 1(a).

Figures S9–S22 present the powers under the simulation settings where $p_f=p_m=p$ but HWE does not hold in female offspring. The left column of each figure represents the powers when $\rho=-0.05$, while the right column denotes the powers when $\rho=0.05$. When comparing the two columns of each figure with the middle column in the corresponding figure under random mating ($\rho=0$), we find that the powers with $\rho=-0.05, 0$

and 0.05 have similar trends, while the powers slightly increase as ρ changes from -0.05 to 0.05 . This is probably due to the increase of genotype frequency of A/A . Finally, Figures S23–S50 display the powers of the other seven methods ($T_A, T_{AD}, T_{AD}^s, FM_{01}, FM_F, Z_{mfA}$ and Z_A), which control the size less well or have relatively low powers.

4. Discussion

In this paper, we propose a robust test, Z_{XCII} , for testing associations between certain diseases and an X-linked SNP by simultaneously accounting for XCI and parent-of-origin effects. Our proposed method is an extension of $Xcat$ for the situation where parent-of-origin effects have influence on the process of XCI. Two reasonable assumptions are made for Z_{XCII} , just like $Xcat$ (Chen *et al.*, 2017): the generalized genetic model is hypothesized for female offspring and the mutant allele in female offspring is the same as that in male offspring. A good feature of the proposed method that should be emphasized is that there is no need to specify the patterns of XCI or parent-of-origin effects. The simulation studies are conducted in order to investigate the validity and performance of Z_{XCII} under various scenarios of parameter values. The simulation results demonstrate that Z_{XCII} is robust in all of the situations considered. It controls the size well and generally outperforms most of the 13 existing methods in power in the presence of parent-of-origin effects, especially complete parent-of-origin effects, although it suffers from slight loss in power when there are no parent-of-origin effects. Thus, the proposed method is a preferred choice when we are not sure whether or not there are parent-of-origin effects in practice.

It should be noted that Z_{XCI} is an extension of $Xcat$. We first use the Fisher's method to combine Z_1 , Z_2 and Z_3 in female offspring (denoted by Z_f) and then obtain the proposed Z_{XCI} by weighting Z_f in female offspring and Z_m in male offspring, while $Xcat$ applies the Fisher's method directly to incorporate the test statistics for females and males (Chen *et al.*, 2017). In fact, we have used the other methods to directly combine the test statistics for females and males, such as Fisher's approach used in Chen *et al.* (2017) and Stouffer's method (Owen, 2009). However, we find that Z_{XCI} is optimal for most of the situations considered. On the other hand, compared to $Xcat$, the regression-based method allows us to adjust for covariates, which is another potential advantage of the proposed method. According to the simulation results (omitted here for brevity), we also found that Z_{XCI} and other methods are not applicable to the association study for rare alleles. We may need to use the SKAT (Wu *et al.*, 2011) or the extensions of SKAT (Larson *et al.*, 2019) for dealing with this situation, which will be our subsequent work. In addition, note that the proposed Z_{XCI} is only suitable for qualitative traits. If we want to analyse quantitative traits in future, we will need to change the logistic regression to multiple linear regression and conduct simulations to compare it with existing methods for quantitative traits. Finally, just like Wang *et al.* (2014), in order to simplify our model, we assumed that XCI-E is regarded as a binary variable to distinguish whether or not XCI is present. However, many genes have been observed to be of 'variable escape', with the levels of escape varying between individuals, cells and tissues or over time. How to consider these variable levels of XCI-E in our model will be our future work.

Supplementary material. For supplementary material accompanying this paper visit <https://doi.org/10.1017/S0016672320000026>.

Author contributions. Yu Zhang helped design the study, drafted the article and conducted the simulation study. Si-Qi Xu helped design the study and drafted the article. Wei Liu revised the article critically. Wing Kam Fung reviewed the whole paper and revised the article. Ji-Yuan Zhou helped design the study, supervised the field activities and directed their implementation, including quality assurance and control. All authors read and approved this version of the manuscript.

Acknowledgements. The authors thank the two reviewers for their helpful comments that greatly improve the presentation of this paper.

Financial support. This work was supported by the National Natural Science Foundation of China (grant number 81773544) and the Hong Kong RGC GRF grant (grant number 17302919).

Conflict of interest. None.

Ethical standards. None.

References

- Carrel L. and Willard H.F. (2005). X-inactivation profile reveals extensive variability in X-linked gene expression in females. *Nature* **434**(7031), 400–404.
- Chen Z., Ng H.K.T., Li J., Liu Q. and Huang H. (2017). Detecting associated single-nucleotide polymorphisms on the X chromosome in case control genome-wide association studies. *Statistical Methods in Medical Research* **26**(2), 567–582.
- Chow J.C., Yen Z., Ziesche S. M. and Brown C.J. (2005). Silencing of the mammalian X chromosome. *Annual Review of Genomics and Human Genetics* **6**, 69–92.
- Clayton D. (2008). Testing for association on the X chromosome. *Biostatistics* **9**(4), 593–600.
- Clayton D.G. (2009). Sex chromosomes and genetic association studies. *Genome Medicine* **1**(11), 110.
- Deng H.W. and Chen W.M. (2001). The power of the transmission disequilibrium test (TDT) with both case–parent and control–parent trios. *Genetics Research* **78**(3), 289–302.
- Dong C., Li W.D., Geller F. *et al.* (2005). Possible genomic imprinting of three human obesity-related genetic loci. *American Journal of Human Genetics* **76**(3), 427–437.
- Donnelly S.L., Wolpert C., Menold M.M. *et al.* (2002). Female with autistic disorder and monosomy X (Turner syndrome): parent-of-origin effect of the X chromosome. *American Journal of Medical Genetics* **96**(3), 312–316.
- Esary J.D., Proschan F. and Walkup D.W. (1967). Association of random variables, with applications. *Annals of Mathematical Statistics* **38**(5), 1466–1474.
- Falls J.G., Pulford D.J., Wylie A.A. and Jirtle R.L. (1999). Genomic imprinting: implications for human disease. *American Journal of Pathology* **154**(3), 635–647.
- Fisher R.A. (1954). *Statistical Methods for Research Workers*. 12th ed. Edinburgh, UK: Oliver and Boyd.
- Gao F., Chang D., Biddanda A. *et al.* (2015). XWAS: a software toolset for genetic data analysis and association studies of the X chromosome. *Journal of Heredity* **106**(5), 666–671.
- Hager R., Cheverud J.M. and Wolf J.B. (2008). Maternal effects as the cause of parent-of-origin effects that mimic genomic imprinting. *Genetics* **178**(3), 1755–1762.
- König I.R., Loley C., Erdmann J. and Ziegler A. (2014). How to include chromosome X in your genome-wide association study. *Genetic Epidemiology* **38**(2), 97–103.
- Larson N.B., Chen J. and Schaid D.J. (2019). A review of kernel methods for genetic association studies. *Genetic Epidemiology* **43**(2), 122–136.
- Li M., Li J., He Z. *et al.* (2016). Testing allele transmission of an SNP set using a family based generalized genetic random field method. *Genetic Epidemiology* **40**(4), 345–351.
- Loesch D.Z., Quang Minh B., Wendy K. *et al.* (2005). Effect of Turner's syndrome and X-linked imprinting on cognitive status: analysis based on pedigree data. *Brain Development* **27**(7), 494–503.
- Loley C., Ziegler A. and König I.R. (2011). Association tests for X-chromosomal markers – a comparison of different test statistics. *Human Heredity* **71**(1), 23–36.
- Migeon B. (1998). Non-random X chromosome inactivation in mammalian cells. *Cytogenetics and Cell Genetics* **80**, 142–148.
- Minks J., Robinson W.P. and Brown C.J. (2008). A skewed view of X chromosome inactivation. *Journal of Clinical Investigation* **118**(1), 20–23.
- Ober C., Loisel D.A. and Gilad Y. (2008). Sex-specific genetic architecture of human disease. *Nature Reviews Genetics* **9**(12), 911–922.
- Owen A.B. (2009). Karl Pearson's meta-analysis revisited. *Annals of Statistics* **37**(6B), 3867–3892.
- Payer B. and Lee J.T. (2008). X chromosome dosage compensation: how mammals keep the balance. *Annual Review of Genetics* **42**(1), 733–772.
- Pessia E., Makino T., Bailly-Bechet M., McLysaght A. and Marais G.A.B. (2012). Mammalian X chromosome inactivation evolved as a dosage-compensation mechanism for dosage-sensitive genes on the X chromosome. *Proceedings of the National Academy of Sciences of the United States of America* **109**(14), 5346–5351.
- Puig X., Ginebra J. and Graffelman J. (2017). A Bayesian test for Hardy–Weinberg equilibrium of biallelic X-chromosomal markers. *Heredity* **119**(4), 226–236.
- Schurz H., Salie M., Tromp G., Hoal E.G., Kinnear C.J. and Möller M. (2019). The X chromosome and sex-specific effects in infectious disease susceptibility. *Human Genomics* **13**(1), 2.
- Skuse D.H. (2000). Imprinting, the X-chromosome, and the male brain: explaining sex differences in the liability to autism. *Pediatric Research* **47**(1), 9–16.
- Starmer J. and Magnuson T. (2009). A new model for random X chromosome inactivation. *Development* **136**(1), 1–10.
- Wallace C., Smyth D.J., Maisuria-Armer M., Walker N.M., Todd J.A. and Clayton D. G. (2010). The imprinted *DLK1-MEG3* gene region on chromosome 14q32.2 alters susceptibility to type 1 diabetes. *Nature Genetics* **42**(1), 68–71.

- Wang J., Yu R. and Shete S.** (2014). X-chromosome genetic association test accounting for X-inactivation, skewed X-inactivation, and escape from X-inactivation. *Genetic Epidemiology* **38**(6), 483–493.
- Wang P., Xu S.Q., Wang B.Q., Fung W.K. and Zhou J.Y.** (2019). A robust and powerful test for case–control genetic association study on X chromosome. *Statistical Methods in Medical Research* **28**(10–11), 3260–3272.
- Wang P., Zhang Y., Wang B.Q. et al.** (2019). A statistical measure for the skewness of X chromosome inactivation based on case–control design. *BMC Bioinformatics* **20**(1), 11.
- Wise A.L., Gyi L. and Manolio T.A.** (2013). eXclusion: toward integrating the X chromosome in genome-wide association analyses. *American Journal of Human Genetics* **92**(5), 643–647.
- Wood S.N.** (2006). *Generalized Additive Models: An Introduction with R. 1st ed.* London, UK: Chapman & Hall Ltd.
- Wu M.C., Lee S., Cai T., Li Y., Boehnke M. and Lin X.** (2011). Rare-variant association testing for sequencing data with the sequence kernel association test. *American Journal of Human Genetics* **89**(1), 82–93.
- Zheng G., Joo J., Zhang C. and Geller N.L.** (2007). Testing association for markers on the X chromosome. *Genetic Epidemiology* **31**(8), 834–843.
- Ziegler A.** (2009). Genome-wide association studies: quality control and population-based measures. *Genetic Epidemiology* **33**(S1), S45–S50.
- Ziegler A. and König I.R.** (2006). *A Statistical Approach to Genetic Epidemiology: Concepts and Applications. 1st ed.* Weinheim, Germany: Wiley-VCH.