

## Research Article

# *Candida albicans* Secreted Aspartic Protease 7 is Essential for Damage of Human Oral Epithelial Cells

Yang HP<sup>1,2</sup>, Tsang PCS<sup>3</sup>, Pow EHN<sup>4</sup>, Lam OLT<sup>5</sup> and Tsang PWK<sup>6,7\*</sup>

<sup>1</sup>The Affiliated Stomatology Hospital, Zhejiang University School of Medicine, China

<sup>2</sup>Key Laboratory of Oral Biomedical Research of Zhejiang Province, Zhejiang University School of Medicine, China

<sup>3</sup>Department of Operative Dentistry, Faculty of Dentistry, The University of Hong Kong, Hong Kong, China

<sup>4</sup>Department of Prosthodontics, Faculty of Dentistry, The University of Hong Kong, Hong Kong, China

<sup>5</sup>George and Fay Yee Centre for Healthcare Innovation, University of Manitoba, Canada

<sup>6</sup>School of General Education and Languages, Technological and Higher Education Institute of Hong Kong, Hong Kong, China

<sup>7</sup>Centre for Interdisciplinary Research on Food By-products Utilization, Technological and Higher Education Institute of Hong Kong, Hong Kong, China

\*Corresponding author: Paul Wai Kei Tsang, Technological and Higher Education Institute of Hong Kong, Hong Kong, China

Received: June 02, 2021; Accepted: July 19, 2021;

Published: July 26, 2021

## Abstract

**Aims:** *Candida albicans* is an important human fungal pathogen in clinical settings. It possesses a wide spectrum of virulence traits, including but not limited to the production of Secreted Aspartic Proteases (SAPs), to invade host cells under predisposing conditions. The aims of the present study were to investigate the functional role of *C. albicans* SAP7 in invasion ability.

**Methods:** The present study was carried out to construct *C. albicans* *sap7Δ/Δ* mutant strain using a PCR-based gene disruption method. The behaviors of this SAP7 knockout strain was evaluated and compared with the wild type and SAP7 complemented strains between human oral epithelial cells with respect to endocytosis, invasion, and tissue damage.

**Results:** Compared with the wild type *C. albicans* strain, a 52% reduction in the endocytosis of the *sap7Δ/Δ* mutant strain by oral epithelial cells was observed, as well as a 25% attenuation of internalization, and a 27% reduction of tissue damage ( $P < 0.05$ ).

**Conclusion:** Our data clearly demonstrates that *C. albicans* SAP7 contributes to tissue invasion into human oral epithelial cells which warrant further investigations as potential targets for antifungal interventions.

**Keywords:** *Candida albicans*; Secreted aspartic protease; Oral epithelial cells; Endocytosis; Tissue damage

## Introduction

*Candida* species are the major cause of opportunistic fungal infections in humans. They exist as harmless commensal in immunocompetent individuals and inhabit in human oral cavity, gastrointestinal and urogenital tracts [1]. However, under predisposing conditions, when normal microbiota is disturbed or immune system is compromised, *Candida* fungi transform into lethal microbes and cause superficial infections or deep-seated, systemic candidiasis [2,3]. In fact, candidiasis ranks fourth among the leading types of nosocomial infections in clinical sectors with high mortality and morbidity rates ranging from 40-60% in spite of careful management of antifungal interventions [4].

It has been considered epithelial tissues are the first line of defense of humans against pathogenic invasion and manifestations. In *Candida* fungi, two distinguished but complementary pathways of invasion to human oral epithelial cells exist: induced endocytosis and active penetration. The induced endocytosis of *C. albicans* is triggered by hyphae that the host cells produce pseudopods and engulf the microorganisms [5,6]. Two *Candida* invasins have been found: Agglutinin-like sequence 3 (Als3) and heat shock protein 1 (Ssa1) found on the surface of hyphae play a key role in induced endocytosis via binding to specific host ligands (E-cadherin on epithelial cells or N-cadherin on endothelial cells) [7]. However, the molecular network and potential cross-talk between these two mechanisms are still ill-defined. Furthermore, the contribution of hydrolytic activity to *Candida* invasion and interaction with host cells is largely unknown.

Secreted aspartic protease (Sap) family, consisting of 10 Saps, is the major hydrolytic enzyme produced by *C. albicans*. It is believed that Saps contribute both to increased endocytosis and active penetration. Previous studies showed that pepstatin A (Sap inhibitor) was able to block *Candida* induced endocytosis in its early stage of invasion, and reduce active penetration when combined with cytochalasin D (cytD) and butanedione monoxime (endocytosis inhibitors) [8]. Besides, significant reduction of induced endocytosis was evident in *sap1-3Δ/Δ* and *sap4-6Δ/Δ* knockout mutants, suggesting the key involvement of these Saps in this process. On the contrary, several studies showed a limited role of Saps 1-6 in *Candida* invasion in a reconstituted model of human epithelia as well as in a mice model [9,10]. Thus, further investigations are warranted to decipher into the functional of different Saps in *Candida* pathobiology.

By now, limited knowledge is available for *C. albicans* Sap7 for its functional significance in pathogenesis, especially in *in vivo* systems [11]. Here, we reported an investigation of *C. albicans* Sap7 in the invasion process of human oral epithelial cells.

## Materials and Methods

### *Candida* strains and culture conditions

*C. albicans* SC5314 was used as the wild type strain in this study for comparison. *C. albicans* *sap7Δ/Δ* mutant strain was constructed by Ura-blaster method [12] and the genotype is  $\Delta sap7::hisG/\Delta sap7::hisG-URA3-hisG$ , which was confirmed by colony PCR with gene-specific primers. All fungal strains were prepared in liquid YPD (1% yeast extract, 2% bactopectone, 2% D-glucose) medium on a

rotary shaker at 30°C overnight. Cell concentrations were adjusted by hemocytometer counting in serum-free DMEM medium immediately prior to the experiments.

### Oral epithelia cells

The OKF6/TERT2 cell line (passage 18) was a gift from J. Rheinwald's Laboratory (Harvard University, Boston, MA) [13]. The OKF6/TERT2 cells were cultured in the humidified incubator containing 5% CO<sub>2</sub> at 37°C. For a standard experiment, 5×10<sup>4</sup> cells/ml of OKF6/TERT2 cells were seeded into μ-Slide 8 well plates (ibidi, Germany) and cultured for 2-3 days post-seeding until 95% confluence.

### Endocytosis assay

The number of fungal cells that were associated with and/or endocytosed by the OKF6/TERT2 cells was determined using a standard differential fluorescence assay with minor modifications [14]. Briefly, 95% confluence OKF6/TERT2 cells were infected with 1×10<sup>5</sup> fungal cells in serum-free DMEM medium. After 3 hrs incubation at 37°C, the cells were gently rinsed by Hank's Balanced Salt Solution (HBSS) twice to remove any nonadherent cells, and fixed with 3% paraformaldehyde. The adherent but not endocytosed fungal cells were stained by a polyclonal rabbit anti-*Candida* antibody conjugated with red fluorescent Alexa Fluor 568 for 1 h. Afterwards, the OKF6/TERT2 cells were permeabilized with 0.5% Triton X-100 in PBS for 30 min. All cell-associated fungal cells (including both adherent and endocytosed) were labelled with anti-*Candida* antibody conjugated with green fluorescent Alexa Fluor 488, and observed under fluorescence microscope. The number of endocytosed fungal cells was determined by subtracting the number of adherent fungal cells (red fluorescing) from the number of cell-associated fungal cells (green fluorescing). Fungal cells which were partially internalized were regarded as endocytosed cells. In each well, at least 100 fungal cells were examined.

### Invasion assay

The number of fungal cells that was active penetrated to oral epithelial cells was determined as described previously [14]. Briefly, the OKF6/TERT2 cells were cultured as described above. To block the induced endocytosis, the OKF6/TERT2 cells were pre-treated with 0.5 μM cyt D for 30 min (this concentration of cyt D was found efficient to stop the rearrangement of actin cytoskeleton without affecting cell viability [8]) and co-incubated throughout the whole infection period. Thus, 1×10<sup>5</sup> fungal cells in serum-free DMEM medium were allowed to infect the OKF6/TERT2 cells for 3 h. Afterwards, non-adhesive fungal cells were gently removed by HBSS washings, the OKF6/TERT2 cells were fixed as described above. All adherent fungal cells were stained with red-fluorescent Alexa Fluor 568. After further rinsing, the OKF6/TERT2 cells were permeabilized, and associated fungal cells were stained with green-fluorescent Alexa Fluor 488 as described above. The percentage of invaded fungal cells was determined by the number of internalized cells (including partially internalized cells) by the number of all the adherent cells. At least 100 fungal cells were examined.

### Damage assay

Tissue damage of the OKF6/TERT2 cells was determined by measuring the amount of released Lactate Dehydrogenase (LDH) into

the surrounding medium upon invasion by *C. albicans* SC5314 and *C. albicans sap7Δ/Δ* mutant strain using CytoTox 96 nonradioactive cytotoxicity assay [15].

### Statistical analysis


All experiments were performed in three separate occasions, and triplicate was performed in each occasion. The difference between the wild type *C. albicans* SC5314 and the *C. albicans sap7Δ/Δ* mutant strain was statistically assessed by Student's *t*-test. A P value <0.05 was considered statistically significant.

## Results and Discussion


The present study was a continuation of our ongoing research on the elucidation of the functional significance of *C. albicans* Saps in its pathobiology. Our previous study clearly revealed, for the first time, the potential role of *C. albicans* Sap9 in serum induced-hyphal formation and interaction with human oral epithelial cells [15], and laid a foundation for further investigations of ill-defined Saps, including Sap7. Previous studies showed a correlation between Sap7 and virulence using an intravenous candidiasis model [8], and was proved to be insensitive to pepstatin A [16]. Here, we investigated the functional significance of Sap7 in *C. albicans* invasion to human oral epithelial cells using a *C. albicans sap7Δ/Δ* mutant strain and compared it with the wild type counterpart.

### Sap7 plays a key role in *C. albicans* induced endocytosis, active penetration, and cell damage in human oral epithelial cells

Induced endocytosis and active penetration are two distinct routes for microorganism's invasion into non-phagocytic host cells such as oral epithelial cells and endothelial cells. The ability of the *C. albicans sap7Δ/Δ* mutant strain in induced endocytosis was found reduced in all the fields examined (Figure 1). A reduction of 52%



**Figure 1:** Fluorescence microscopic analysis of the interaction between *Candida* cells and human oral epithelial cells. The endocytosed cells were labelled as red fluorescence while the associated cells were labelled as green fluorescence.



**Figure 2:** Analyses of the functional role of *C. albicans* Sap7 in (A) endocytosis; (B) invasion; and (C) LDH activity upon co-incubation with the human oral epithelial cells ( $P < 0.05$ ).

( $P < 0.05$ ) of induced endocytosis was observed in the *C. albicans sap7Δ/Δ* mutant strain on OKF6/TERT2 cells (Figure 2A); the number of cell-associated *C. albicans sap7Δ/Δ* mutant strain was 34% less than the wild type *C. albicans* SC5314, although the results were not statistically significant. It is envisaged that Sap7 facilitates *Candida* induced endocytosis via possible cellular interactions with other invasins on hyphae and/or other host cell receptors which warrant further investigations.

To investigate the role of Sap7 in active penetration, we co-incubated the fungal cells with the human oral epithelial cells in the presence of the specific microfilament inhibitor cytD (0.5  $\mu\text{m}$ ) in an attempt to selectively block all induced endocytosis. The proportion of the *C. albicans sap7Δ/Δ* mutant strain that penetrated into the OKF6/TERT2 cells was significantly smaller than (25%;  $P < 0.05$ ) that of the wild type *C. albicans* SC5314 (Figure 2B), suggesting that Sap7 might play a key role in *C. albicans* active penetration in human oral epithelial cells.

We measured the amount of released LDH into co-culture supernatant to investigate quantitatively the degree of tissue damage caused by the fungal cells. The ability of the *C. albicans sap7Δ/Δ* mutant strain to infect the OKF6/TERT2 cells was moderately reduced by 27% ( $P < 0.05$ ) when compared with the wild type *C. albicans* SC5314, suggesting that the knockout of *Sap7* gene led to a weakened invasion power to the human oral epithelial cells (Figure 2C).

## Conclusion

Taken together, our present study clearly indicated that the *C. albicans sap7Δ/Δ* mutant strain exhibits attenuated capacity to adhering to and endocytosed by the human oral epithelial cells,

and therefore suggesting that *C. albicans* Sap7 contributes to tissue invasion process and leads to host cell damage. As little is known about the cellular pathways/interacting molecules involved in *C. albicans* Sap7 pathogenesis, further investigations using proteomic study are warranted.

## Funding

The present study was supported by the funding of the University of Hong Kong (No.: 201007176095).

## References

- Mohr A, Simon M, Joha T, Hanses F, Salzberger B, Hitzentichler F. Epidemiology of candidemia and impact of infectious disease consultation on survival and care. *Infection* 2020; 48: 275-284.
- Lim CSY, Rosli R, Seow HF, Chong PP. *Candida* and invasive candidiasis: back to basics. *Eur J Clin Microbiol Infect Dis*. 2012; 31: 21-31.
- Sardi JCO, Scorzoni L, Bernardi T, Fusco-Almeida AM, Mendes Giannini MJ. *Candida* species: current epidemiology, pathogenicity, biofilm formation, natural antifungal products and new therapeutic options. *J Med Microbiol*. 2013; 62: 10-24.
- Antinori S, Milazzo L, Sollima S, Galli M, Corbellino M. Candidemia and invasive candidiasis in adults: a narrative review. *Eur J Intern Med*. 2016; 34: 21-28.
- Friedman DZP, Schwartz IS. Emerging fungal infections: new patients, new patterns, and new pathogens. *J Fungi (Basel)*. 2019; 5: 67.
- Sheppard DC, Filler SG. Host cell invasion by medically important fungi. *Cold Spring Harb Perspect Med*. 2014; 5: a019687.
- Yang W, Yan L, Wu C, Zhao X, Tang J. Fungal invasion of epithelial cells. *Microbiol Res*. 2014; 169: 803-811.
- Dalle F, Wachtler B, L'Ollivier C, Holland G, Bannert N, Wilson D, et al. Cellular interactions of *Candida albicans* with human oral epithelial cells and enterocytes. *Cell Microbiol*. 2010; 12: 248-271.

9. Lermann U, Morschhauser J. Secreted aspartic proteases are not required for invasion of reconstituted human epithelia by *Candida albicans*. *Microbiology*. 2008; 154: 3281-3295.
10. Correia A, Lermann U, Teixeira L, Cerca F, Botelho S, da Costa RM, et al. Limited role of secreted aspartyl proteinases Sap1 to Sap6 in *Candida albicans* virulence and host immune response in murine hematogenously disseminated candidiasis. *Infect Immun*. 2010; 78: 4829-4849.
11. Naglik JR, Moyes D, Makwana J, Kanzaria P, Tsihlaki E, Weindl G, et al. Quantitative expression of the *Candida albicans* secreted aspartyl proteinase gene family in human oral and vaginal candidiasis. *Microbiology*. 2008; 154: 3266-3280.
12. Noble SM, Johnson AD. Strains and strategies for large-scale gene deletion studies of the diploid human fungal pathogen *Candida albicans*. *Eukaryot Cell*. 2005; 4: 298-309.
13. Dickson MA, Hahn WC, Ino Y, Ronfard V, Wu JY, Weinberg RA, et al. Human keratinocytes that express hTERT and also bypass a p16(INK4a)-enforced mechanism that limits life span become immortal yet retain normal growth and differentiation characteristics. *Mol Cell Biol*. 2000; 20: 1436-1447.
14. Yang H, Tsang PCS, Pow EHN, Lam OLT, Tsang PWK. Potential role of *Candida albicans* secreted aspartic protease 9 in serum induced-hyphal formation and interaction with oral epithelial cells. *Microb Pathog*. 2020; 139: 103896.
15. Tsang PWK, Fong WP, Samaranyake LP. *Candida albicans* orf19.3727 encodes phytase activity and is essential for human tissue damage. *PLoS One*. 2017; 12: e0189219.
16. Aoki W, Kitahara N, Miura N, Morisaka H, Yamamoto Y, Kuroda K, et al. *Candida albicans* possesses Sap7 as a pepstatin A-insensitive secreted aspartic protease. *PLoS One* 2012; 7: e32513.