



Editorial: Acute Promyelocytic Leukemia – Towards A Chemotherapy-Free Approach to Cure in All Patients

Harinder Gill^{1*}, Yok-Lam Kwong¹ and Farhad Ravandi²

¹ Department of Medicine, The University of Hong Kong, Hong Kong, Hong Kong SAR, China, ² Department of Leukemia, University of Texas MD Anderson Cancer Center, Houston, TX, United States

Keywords: acute promyelocytic leukemia, arsenic trioxide, chemotherapy-free, tetraarsenic tetrasulfide, oral arsenic trioxide

Editorial on the Research Topic

Acute Promyelocytic Leukemia - Towards a Chemotherapy-Free Approach to Cure in All Patients

Acute promyelocytic leukemia (APL) is a subtype of acute myeloid leukemia (AML) characterized by t(15;17)(q24;21) and the fusion gene *PML-RARA*. With optimal supportive care and frontline use of all-trans retinoic acid (ATRA) and chemotherapy, first complete remission (CR1) rates of more than 90% and long-term survival of more 80% can be achieved. Regimens that include As₂O₃, ATRA and chemotherapy result in a CR rare of 90-100% and OS between 86-97% (1–3). Frontline treatment of APL has evolved rapidly. An emerging theme is the incorporation of As₂O₃ early in the treatment algorithm, starting from induction to consolidation.

Various forms of arsenic were used in China for over 5000 years. Arsenic first appeared in Western Medicine in the eighteenth century. In hematology, oral arsenic was first reported in the treatment of chronic myeloid leukemia from the 1860s to 1920s in Germany and Boston City (4). This treatment was phased out following World War II with the development of alkylating chemotherapy and radiotherapy. Intravenous pure As₂O₃ solution was first used in Harbin, China in 1973. Its mechanism of action, pharmacokinetics and clinical efficacy was extensively published in 1996. In this Research Topic collection, Kumana et al. gave a historical account on the development of pure oral arsenic trioxide that was invented and patented in Hong Kong. With memories of the Fowler's solution, an oral As₂O₃ formulation or the "modern" liquor arsenicalis was revived in 1998 in Hong Kong. In Hong Kong, China, an oral preparation of As₂O₃ (oral-As₂O₃) was developed, and was shown to be efficacious for APL in first relapse (R1), inducing second complete remission (CR2) in more than 90% of patients (5). Further, in an effort to prevent relapses, oral-As₂O₃ was used during induction and maintenance of CR1 (6, 7). This strategy resulted in favorable overall-survival (OS) and leukemia-free-survival (LFS). This 1mg/ml oral-As₂O₃ solution has a bioavailability comparable with that of i.v. As₂O₃ (8). Oral arsenic trioxide (Arsenol[®]) from Hong Kong was also the first oral preparation of pure arsenic trioxide produced under the Good Manufacturing Practice (GMP) standards. On the other hand, Realgar-Indigo naturalis formula (RIF) was developed in the 1980s entirely based on traditional Chinese Medicine (TCM) concepts and comprises realgar, *Indigo naturalis*, *Salvia miltiorrhiza* and *Radix pseudostellariae* (9). Tetraarsenic tetrasulfide (As₄S₄), indirubin and tanshinone IIA are the

OPEN ACCESS

Edited and reviewed by:

Alessandro Isidori,
AORMN Hospital, Italy

*Correspondence:

Harinder Gill
gillhsh@hku.hk

Specialty section:

This article was submitted to
Hematologic Malignancies,
a section of the journal
Frontiers in Oncology

Received: 08 December 2021

Accepted: 27 December 2021

Published: 20 January 2022

Citation:

Gill H, Kwong Y-L and
Ravandi F (2022) Editorial:
Acute Promyelocytic Leukemia –
Towards A Chemotherapy-Free
Approach to Cure in All Patients.
Front. Oncol. 11:831308.
doi: 10.3389/fonc.2021.831308

active ingredients of RIF with *in-vitro* synergism (9). In this Research Topic, Zhu and Lou et al. described the all oral, chemotherapy-free model in the frontline management of APL highlighting the applications of RIF and summarized the clinical data and excellent outcome of patients treated with RIF-based induction. Zhu also highlighted a major obstacle to cure in APL, that is, early deaths. In addition, multi-centre clinical trials in patients treated with ATRA, arsenic trioxide and anthracyclines reported a relatively low incidence of early deaths of 3-10% (1, 2). Xu and Huang discuss the evolution of various therapeutic approaches from hospital-based induction and consolidation to home-based oral As₂O₃-based therapy especially during consolidation and maintenance. Other oral formulations of As₂O₃ are also being investigated with the hope of eventually developing an all oral, effective regimen in standard-risk APL (10).

With regard to the therapeutic application of intravenous arsenic trioxide, Russell and Dillon summarize the United Kingdom's NCRI AML17 experience using the attenuated arsenic trioxide regimen in newly diagnosed APL. The treatment of "high-risk" APL remains a topic of contention as As₂O₃ is not yet approved for this indication. An important exploratory study by the MD Anderson Cancer Center using ATRA-As₂O₃ with or without gemtuzumab ozogamycin (GO) suggested that an essentially chemotherapy-free regimen might be feasible for the upfront treatment of APL (11). The NCRI AML17 trial built on these findings to investigate the de-intensification of treatment by randomizing patients irrespective of their risk status between As₂O₃-ATRA and the ATRA-idarubicin (AIDA) regimen. The AML17 trial included a total of 57 high-risk patients and their overall survival at 4 years was not significantly different from standard risk patients, being

95% in standard-risk compared with 87% in high-risk patients. Of the 28 high-risk patients in the As₂O₃ arm of AML17 who received the planned induction of ATRA, As₂O₃ and GO, the 4-year survival was 89% (12).

Last, but not least, Sanz et al. reappraised the role of hematopoietic stem cell transplantation (HSCT) in the current era of arsenic trioxide. As₂O₃-based regimens are currently regarded as the first option for relapsed APL. The selection of the most appropriate post-remission treatment option for patients in second CR (CR2), including the use of HSCT, depends on several variables, such as pre-transplant molecular status, duration of first remission, age, and donor availability. Despite modest evidence from retrospective studies, autologous HSCT has been the preferred option for consolidation in patients achieving molecularly negative CR2. The suitability of HSCT as compared with other non-HSCT alternatives has recently engendered much interest, thereby necessitating prospective controlled studies to define the role of HSCT in APL.

With the above backdrop, this special Research Topic provides an overview on the biology of APL and highlights perspectives on the past, present, and future treatment approaches in APL. This topic plays an important role in addressing the development of As₂O₃-based therapy in improving the outcome of APL, from once the most fatal to currently the most curable leukemia.

AUTHOR CONTRIBUTIONS

HG wrote and approved the manuscript. All authors contributed to the article and approved the submitted version.

REFERENCES

- Lo-Coco F, Avvisati G, Vignetti M, Thiede C, Orlando SM, Iacobelli S, et al. Retinoic Acid and Arsenic Trioxide for Acute Promyelocytic Leukemia. *N Engl J Med* (2013) 369(2):111–21. doi: 10.1056/NEJMoa1300874
- Iland HJ, Bradstock K, Supple SG, Catalano A, Collins M, Hertzberg M, et al. All-Trans-Retinoic Acid, Idarubicin, and IV Arsenic Trioxide as Initial Therapy in Acute Promyelocytic Leukemia (APML4). *Blood* (2012) 120(8):1570–80; quiz 1752. doi: 10.1182/blood-2012-02-410746
- Abaza Y, Kantarjian H, Garcia-Manero G, Estey E, Borthakur G, Jabbour E, et al. Long-Term Outcome of Acute Promyelocytic Leukemia Treated With All-Trans-Retinoic Acid, Arsenic Trioxide, and Gemtuzumab. *Blood* (2017) 129(10):1275–83. doi: 10.1182/blood-2016-09-736686
- Kwong YL, Todd D. Delicious Poison: Arsenic Trioxide for the Treatment of Leukemia. *Blood* (1997) 89(9):3487–8. doi: 10.1182/blood.V89.9.3487
- Gill H, Yim R, Lee HKK, Mak V, Lin SY, Kho B, et al. Long-Term Outcome of Relapsed Acute Promyelocytic Leukemia Treated With Oral Arsenic Trioxide-Based Reinduction and Maintenance Regimens: A 15-Year Prospective Study. *Cancer* (2018) 124(11):2316–26. doi: 10.1002/cncr.31327
- Gill H, Kumana CR, Yim R, Hwang YY, Chan TSY, Yip SF, et al. Oral Arsenic Trioxide Incorporation Into Frontline Treatment With All-Trans Retinoic Acid and Chemotherapy in Newly Diagnosed Acute Promyelocytic Leukemia: A 5-Year Prospective Study. *Cancer* (2019) 125(17):3001–12. doi: 10.1002/cncr.32180
- Gill HS, Yim R, Kumana CR, Tse E, Kwong YL. Oral Arsenic Trioxide, All-Trans Retinoic Acid, and Ascorbic Acid Maintenance After First Complete Remission in Acute Promyelocytic Leukemia: Long-Term Results and Unique Prognostic Indicators. *Cancer* (2020) 126(14):3244–54. doi: 10.1002/cncr.32937
- Kumana CR, Au WY, Lee NS, Kou M, Mak RW, Lam CW, et al. Systemic Availability of Arsenic From Oral Arsenic-Trioxide Used to Treat Patients With Hematological Malignancies. *Eur J Clin Pharmacol* (2002) 58(8):521–6. doi: 10.1007/s00228-002-0514-x
- Wang L, Zhou GB, Liu P, Song JH, Liang Y, Yan XJ, et al. Dissection of Mechanisms of Chinese Medicinal Formula Realgar-Indigo Naturalis as an Effective Treatment for Promyelocytic Leukemia. *Proc Natl Acad Sci USA* (2008) 105(12):4826–31. doi: 10.1073/pnas.0712365105
- Ravandi F, Koumenis I, Johri A, Tallman M, Roboz GJ, Strickland S, et al. Oral Arsenic Trioxide ORH-2014 Pharmacokinetic and Safety Profile in Patients With Advanced Hematologic Disorders. *Haematologica* (2020) 105(6):1567–74. doi: 10.3324/haematol.2019.229583
- Ravandi F, Estey E, Jones D, Faderl S, O'Brien S, Fiorentino J, et al. Effective Treatment of Acute Promyelocytic Leukemia With All-Trans-Retinoic Acid, Arsenic Trioxide, and Gemtuzumab Ozogamicin. *J Clin Oncol* (2009) 27(4):504–10. doi: 10.1200/JCO.2008.18.6130
- Burnett AK, Russell NH, Hills RK, Bowen D, Kell J, Knapper S, et al. Arsenic Trioxide and All-Trans Retinoic Acid Treatment for Acute Promyelocytic Leukaemia in All Risk Groups (AML17): Results of a Randomised, Controlled, Phase 3 Trial. *Lancet Oncol* (2015) 16(13):1295–305. doi: 10.1016/S1470-2045(15)00193-X

Conflict of Interest: The University of Hong Kong currently holds two United States (US) patents (7,521,071 B2 and 8,906,422 B2), one Japanese patent (4786341) and one European patent (EP 1562616 B1) for the use of oral arsenic

trioxide in the treatment of leukemias and lymphomas. HG is employed by the University of Hong Kong.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in

this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Gill, Kwong and Ravandi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.