

Article

Matrix-Tilted Archimedean Copulas

Marius Hofert ^{1,*}  and Johanna F. Ziegel ² 

¹ Department of Statistics and Actuarial Science, University of Waterloo, 200 University Avenue West, Waterloo, ON N2L 3G1, Canada

² Institute of Mathematical Statistics and Actuarial Science, University of Bern, Alpeneggstrasse 22, 3012 Bern, Switzerland; johanna.ziegel@stat.unibe.ch

* Correspondence: marius.hofert@uwaterloo.ca

Abstract: The new class of matrix-tilted Archimedean copulas is introduced. It combines properties of Archimedean and elliptical copulas by introducing a tilting matrix in the stochastic representation of Archimedean copulas, similar to the Cholesky factor for elliptical copulas. Basic properties of this copula construction are discussed and a further extension outlined.

Keywords: Archimedean copulas; elliptical copulas; stochastic representation; generalization; tilting

1. Introduction

Elliptical distributions are among the most important multivariate distributions and are widely used, for example, in finance, insurance, and risk management to model multivariate risk factor changes. A d -dimensional random vector \mathbf{X} is said to be *elliptically distributed* with location vector $\mathbf{0}$, if it admits the stochastic representation

$$\mathbf{X} = \mathbf{R}\mathbf{A}\mathbf{U}, \quad (1)$$

where the (random) *radial part* $R \geq 0$ is independent of the random vector \mathbf{U} which follows a uniform distribution on the Euclidean unit sphere \mathbb{S}^{k-1} and $\mathbf{A} \in \mathbb{R}^{d \times k}$ is a matrix of rank k (see [Cambanis et al. \(1981\)](#)); in what follows, we assume $R \sim F_R$ with $F_R(0) = 0$ and $k = d$. *Elliptical copulas* are the copulas arising from elliptical distributions via Sklar's Theorem; we assume the reader is familiar with the notion of copulas and related concepts of measuring dependence between the components of a random vector (see [Nelsen 2006](#) for an introduction). Despite their popularity (partly due to the comparably simple simulation approach based on the stochastic representation (1)), elliptical copulas suffer from well-known limitations such as radial symmetry, which, in particular, implies that the lower and upper tail dependence coefficients are equal. This is often an unrealistic assumption from a practical point of view.

Another popular class of copulas are *Archimedean copulas*, which are copulas admitting the representation

$$C(\mathbf{u}) = \psi(\psi^{-1}(u_1) + \dots + \psi^{-1}(u_d)), \quad \mathbf{u} \in [0, 1]^d,$$

for some $\psi : [0, \infty) \rightarrow [0, 1]$ known as (*Archimedean*) *generator*. [McNeil and Nešlehová \(2009\)](#) showed that a stochastic representation similar to (1) can be derived for Archimedean copulas. Archimedean copulas arise as survival copulas of random vectors \mathbf{X} with stochastic representation

$$\mathbf{X} = \mathbf{R}\mathbf{U}, \quad (2)$$

where the radial part R is as in (1), independent of \mathbf{U} , but \mathbf{U} is now uniformly distributed on the standard simplex $\Delta_d = \{\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d \mid x_j \geq 0, j = 1, \dots, d, x_1 + \dots + x_d = 1\}$,



Citation: Hofert, Marius, and Johanna F. Ziegel. 2021. Matrix-Tilted Archimedean Copulas. *Risks* 9: 68. <https://doi.org/10.3390/risks9040068>

Academic Editor: Mogens Steffensen

Received: 5 January 2021

Accepted: 24 March 2021

Published: 6 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

that is, $\mathbf{U} \sim U(\Delta_d)$. The Archimedean generator ψ corresponds to the Williamson d -transform $\mathcal{W}_d[F_R]$ of the radial part distribution F_R of F and is given by

$$\psi(t) = \mathcal{W}_d[F_R](t) = \int_{(t, \infty)} \left(1 - \frac{t}{r}\right)^{d-1} dF_R(r), \quad t \in [0, \infty). \quad (3)$$

In contrast to elliptical copulas, Archimedean copulas are not restricted to radial symmetry. They are often given explicitly and properties can typically be described in terms of the generator ψ . However, one major drawback is that they are exchangeable, which implies, for example, that all pairs of random variables share the same dependence structure. This might be unrealistic from a practical point of view (see [Embrechts and Hofert 2011](#) for a discussion).

Analogously to the stochastic representation given in (1), the main contribution of this paper is to introduce a matrix A in the stochastic representation (2) which allows one to create asymmetric copulas which are limited to neither exchangeability nor radial symmetry. The construction of this new class of copulas, referred to as *matrix-tilted Archimedean copulas*, is given in Section 2. Section 3 discusses properties of matrix-tilted Archimedean copulas. Further extensions are mentioned in Section 4, while Section 5 concludes. All proofs are provided in the Appendix A. Other novel copula constructions not further discussed here include, for example, those in ([Genest et al. 2018](#); [Krupskii and Joe 2013 2015](#); [Quessy and Durocher 2019](#); [Quessy et al. 2016](#)).

2. The Construction

We are interested in the survival copulas of multivariate random vectors of the form

$$\mathbf{X} = R\mathbf{A}\mathbf{U}, \quad (4)$$

where R and $\mathbf{U} \sim U(\Delta_d)$ are as above, and $A = (a_{ij})_{i,j=1,\dots,d}$ is a $(d \times d)$ -matrix with real entries $a_{ij} \in \mathbb{R}$. The intuition behind the resulting dependence structure is the following. Naturally, our construction allows for asymmetric copulas; the degree of asymmetry could be quantified, for example, with the (scaled) supremal distance between the survival copula and copula of (4) (see [Rosco and Joe 2013](#)). For positive entries in A , mass gets squeezed towards the diagonal. For general entries, it is also possible to move mass towards the secondary diagonal. Depending on the different conditions imposed on A , the dependence structure of \mathbf{X} can vary substantially, which can make this construction interesting for scenario generation in risk applications. For example, Figures 1 and 2 show samples of the empirical survival copula of \mathbf{X} in (4) based on the same 1000 realizations of R and \mathbf{U} but tilted with different matrices A ; note that all scatter plots of samples of such type in this paper were generated in this way. The points are colored according to the quadrant the corresponding realization of \mathbf{X} lie in and the boundary curves of the support of the copula density is depicted.




Figure 1. Scatter plots of 1000 samples from matrix-tilted Archimedean copulas with different tilting matrices A but otherwise equal realizations of R and \mathbf{U} . The plot on the left simply corresponds to a Gumbel copula with parameter $\theta = 2$ (Kendall’s tau 0.5). The plot on the right also displays the boundary curves (spanned by \tilde{e}_1, \tilde{e}_2 ; see Proposition 1 below).




Figure 2. Similar to Figure 1 with other tilting matrices A .

In what follows, we discuss a number of interesting cases and properties of matrix-tilted Archimedean copulas.

3. Properties

For the sake of tractability of the calculations and the fact that properties of multivariate copulas are often studied for bivariate margins (measures of association, tail dependence, symmetries, etc.), we focus on the bivariate case, that is, $d = 2$, and assume that

$$a_{11} = a_{22} = 1, \quad a_{12} \leq 1, \quad a_{21} \leq 1, \quad a_{12}a_{21} < 1; \quad (\text{CPD})$$

Section 3.1 addresses why Assumption (CPD) is natural. It often turns out to be useful to consider the random point

$$\tilde{\mathbf{U}} = \mathbf{A}\mathbf{U}, \quad (5)$$

so the image of \mathbf{U} under \mathbf{A} . All such images are denoted by their respective notation using a tilde. If $\mathbf{A} = \mathbf{I}_2$, that is, the identity matrix in the bivariate case, then $\tilde{\mathbf{U}} = \mathbf{A}\mathbf{U} = \mathbf{U}$. The columns of \mathbf{A} are e_1, e_2 in this case (the canonical basis vectors of \mathbb{R}^2) and our construction simply leads to the class of Archimedean copulas. In the general case, \mathbf{A} transforms e_1, e_2 to \tilde{e}_1, \tilde{e}_2 . Figures 3 and 4 depict the space (randomly) spanned by (2) and (4), corresponding to Figures 1 and 2.




Figure 3. Depicting how (5) maps the vectors $e_1 = (1, 0)$ and $e_2 = (0, 1)$ to $\tilde{e}_1 = (1, a_{21} = 1/2)$ and $\tilde{e}_2 = (a_{12} = 1/20, 1)$, respectively, corresponding to Figure 1.




Figure 4. Similar to Figure 3 but with vectors \tilde{e}_1, \tilde{e}_2 corresponding to Figure 2.

3.1. Special Cases

In this section, we briefly address various special cases of our construction and explain why Assumption (CPD) is natural.

First note that copulas are invariant under strictly increasing transformations of the margins. We can thus always multiply each component of \mathbf{X} by positive constants without losing flexibility of our dependence model; the same scaling is commonly used for elliptical copulas where one can assume A to be the Cholesky factor of a correlation (instead of a more general covariance) matrix. Therefore, we can equally well assume that $a_{11} = a_{22} = 1$, the first part of Assumption (CPD). The convex cone between the half-lines spanned by $\tilde{e}_j, j \in \{1, 2\}$, the columns of A , does not depend on the order of the columns in A . We can thus fix the determinant to be positive, the last part of Assumption (CPD). The assumptions $a_{12} \leq 1$ and $a_{21} \leq 1$ are for computational convenience. In the limiting case, when $a_{12} = a_{21} = 1$, we obtain $\det A = 0$ and the corresponding matrix-tilted Archimedean copula is the upper Fréchet–Hoeffding bound M .

3.2. The Multivariate Survival Function

We can now provide the survival function of $\mathbf{X} = RAU$ under Assumption (CPD).

Theorem 1. Under Assumption (CPD), the joint survival function of Model (4) at (x_1, x_2) is given as follows.

(i) If $0 \leq a_{12} < 1$ and $0 \leq a_{21} < 1$, then

$$\begin{cases} 1, & \text{if } a_{21}x_1 \geq x_2 \text{ and } a_{12}x_2 \geq x_1, \\ \frac{1-a_{12}a_{21}}{(1-a_{12})(1-a_{21})} \psi\left(\frac{(1-a_{21})x_1+(1-a_{12})x_2}{1-a_{12}a_{21}}\right) \\ \quad - \frac{a_{12}}{1-a_{12}} \psi\left(\frac{x_1}{a_{12}}\right) - \frac{a_{21}}{1-a_{21}} \psi\left(\frac{x_2}{a_{21}}\right), & \text{if } a_{21}x_1 < x_2 \text{ and } a_{12}x_2 < x_1, \\ \mathbb{1}_{\{x_2 \leq 0\}} + \frac{1}{1-a_{21}} \left(\psi(x_2) - a_{21} \psi\left(\frac{x_2}{a_{21}}\right) \right) \mathbb{1}_{\{x_2 > 0\}}, & \text{if } a_{21}x_1 < x_2 \text{ and } a_{12}x_2 \geq x_1, \\ \mathbb{1}_{\{x_1 \leq 0\}} + \frac{1}{1-a_{12}} \left(\psi(x_1) - a_{12} \psi\left(\frac{x_1}{a_{12}}\right) \right) \mathbb{1}_{\{x_1 > 0\}}, & \text{if } a_{21}x_1 \geq x_2 \text{ and } a_{12}x_2 < x_1. \end{cases}$$

(ii) If $a_{12} < 0$ and $a_{21} < 0$, then

$$\begin{cases} \mathbb{1}_{\{x_2 > 0\}} \left(\frac{a_{12}}{1-a_{12}} \psi(x_1/a_{12}) + \frac{1}{1-a_{21}} \psi(x_2) \right) \\ \quad + \mathbb{1}_{\{x_1 > 0\}} \left(\frac{a_{21}}{1-a_{21}} \psi(x_2/a_{21}) + \frac{1}{1-a_{12}} \psi(x_1) \right) \\ \quad + \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 \leq 0\}} \left(\frac{a_{12}}{1-a_{12}} \psi(x_1/a_{12}) + \frac{a_{21}}{1-a_{21}} \psi(x_2/a_{21}) + 1 \right), & \text{if } a_{21}x_1 \geq x_2 \text{ and } a_{12}x_2 \geq x_1, \\ \frac{1-a_{12}a_{21}}{(1-a_{12})(1-a_{21})} \psi\left(\frac{(1-a_{21})x_1+(1-a_{12})x_2}{1-a_{12}a_{21}}\right), & \text{if } a_{21}x_1 < x_2 \text{ and } a_{12}x_2 < x_1, \\ \frac{1}{1-a_{21}} \psi(x_2) + \frac{a_{12}}{1-a_{12}} \psi(x_1/a_{12}), & \text{if } a_{21}x_1 < x_2 \text{ and } a_{12}x_2 \geq x_1, \\ \frac{1}{1-a_{12}} \psi(x_1) + \frac{a_{21}}{1-a_{21}} \psi(x_2/a_{21}), & \text{if } a_{21}x_1 \geq x_2 \text{ and } a_{12}x_2 < x_1. \end{cases}$$

(iii) If $0 \leq a_{12} < 1$ and $a_{21} < 0$, then

$$\begin{cases} \mathbb{1}_{\{x_2 > 0\}} \frac{1}{1-a_{21}} \psi(x_2) + \mathbb{1}_{\{x_2 \leq 0\}} \left(1 + \frac{a_{21}}{1-a_{21}} \psi(x_2/a_{21}) \right), & \text{if } a_{21}x_1 \geq x_2 \text{ and } a_{12}x_2 \geq x_1, \\ \frac{1-a_{12}a_{21}}{(1-a_{12})(1-a_{21})} \psi\left(\frac{(1-a_{21})x_1+(1-a_{12})x_2}{1-a_{12}a_{21}}\right) - \frac{a_{12}}{(1-a_{12})} \psi(x_1/a_{12}), & \text{if } a_{21}x_1 < x_2 \text{ and } a_{12}x_2 < x_1, \\ \frac{1}{1-a_{21}} \psi(x_2), & \text{if } a_{21}x_1 < x_2 \text{ and } a_{12}x_2 \geq x_1, \\ \frac{a_{21}}{1-a_{21}} \psi(x_2/a_{21}) + \mathbb{1}_{\{x_1 \leq 0\}} \\ \quad + \mathbb{1}_{\{x_1 > 0\}} \frac{1}{1-a_{12}} (\psi(x_1) - a_{12} \psi(x_1/a_{12})), & \text{if } a_{21}x_1 \geq x_2 \text{ and } a_{12}x_2 < x_1. \end{cases}$$

(iv) If $a_{12} = 1$ and $0 \leq a_{21} < 1$, then

$$\begin{cases} 1, & \text{if } a_{21}x_1 \geq x_2 \geq x_1, \\ \frac{1}{1-a_{21}} \left((1 - \frac{x_2}{x_1}) \bar{F}_R(x_1) + \frac{x_2}{x_1} \psi(x_1) \right) - \frac{a_{21}}{1-a_{21}} \psi(x_2/a_{21}), & \text{if } a_{21}x_1 < x_2 < x_1, \\ \frac{1}{1-a_{21}} (\psi(x_2) - a_{21} \psi(x_2/a_{21})), & \text{if } a_{21}x_1 < x_2 \text{ and } x_2 \geq x_1, \\ \bar{F}_R(\max\{0, x_1\}), & \text{if } a_{21}x_1 \geq x_2 \text{ and } x_2 < x_1. \end{cases}$$

(v) If $a_{12} = 1$ and $a_{21} < 0$, then

$$\begin{cases} \mathbb{1}_{\{x_2 \leq 0\}} \left(1 + \frac{a_{21}}{1-a_{21}} \psi(x_2/a_{21}) \right) + \mathbb{1}_{\{x_2 > 0\}} \frac{1}{1-a_{21}} \psi(x_2), & \text{if } a_{21}x_1 \geq x_2 \geq x_1, \\ \frac{1}{1-a_{21}} \left((1 - \frac{x_2}{x_1}) \bar{F}_R(x_1) + \frac{x_2}{x_1} \psi(x_1) \right), & \text{if } a_{21}x_1 < x_2 < x_1, \\ \frac{1}{1-a_{21}} \psi(x_2), & \text{if } a_{21}x_1 < x_2 \text{ and } x_2 \geq x_1, \\ \bar{F}_R(\max\{0, x_1\}) + \frac{a_{21}}{1-a_{21}} \psi(x_2/a_{21}), & \text{if } a_{21}x_1 \geq x_2 \text{ and } x_2 < x_1. \end{cases}$$

The remaining cases can be obtained by exchanging the roles of a_{12}, a_{21} and x_1, x_2 .

The following corollary considers two interesting special cases for the parameters a_{12} and a_{21} .

Corollary 1. Suppose Assumption (CPD) holds. Then, the joint survival function of Model (4) is given as follows.

(i) If $a_{21} = 0$ and $a_{12} < 1$, then

$$\begin{cases} 1, & \text{if } 0 \geq x_2 \text{ and } a_{12}x_2 \geq x_1, \\ \frac{1}{1-a_{12}} \psi(x_1 + (1 - a_{12})x_2) - \frac{a_{12}}{(1-a_{12})} \psi(x_1/a_{12}), & \text{if } 0 < x_2 \text{ and } a_{12}x_2 < x_1, \\ \frac{1}{1-a_{21}} \psi(x_2), & \text{if } 0 < x_2 \text{ and } a_{12}x_2 \geq x_1, \\ \mathbb{1}_{\{x_1 \leq 0\}} + \mathbb{1}_{\{x_1 > 0\}} \frac{1}{1-a_{12}} (\psi(x_1) - a_{12} \psi(x_1/a_{12})), & \text{if } 0 \geq x_2 \text{ and } a_{12}x_2 < x_1. \end{cases}$$

(ii) If $a_{12} = a_{21} = a$, then, for $0 \leq a < 1$, we have

$$\begin{cases} 1, & \text{if } ax_1 \geq x_2 \text{ and } ax_2 \geq x_1, \\ \frac{1}{1-a} \left((1 + a) \psi\left(\frac{x_1+x_2}{1+a}\right) - a \psi\left(\frac{x_1}{a}\right) - a \psi\left(\frac{x_2}{a}\right) \right), & \text{if } ax_1 < x_2 \text{ and } ax_2 < x_1, \\ \mathbb{1}_{\{x_2 \leq 0\}} + \frac{1}{1-a} \left(\psi(x_2) - a \psi\left(\frac{x_2}{a}\right) \right) \mathbb{1}_{\{x_2 > 0\}}, & \text{if } ax_1 < x_2 \text{ and } ax_2 \geq x_1, \\ \mathbb{1}_{\{x_1 \leq 0\}} + \frac{1}{1-a} \left(\psi(x_1) - a \psi\left(\frac{x_1}{a}\right) \right) \mathbb{1}_{\{x_1 > 0\}}, & \text{if } ax_1 \geq x_2 \text{ and } ax_2 < x_1, \end{cases}$$

and, for $a \in (-1, 0)$, we obtain that

$$\begin{cases} \mathbb{1}_{\{x_2 > 0\}} \left(\frac{a}{1-a} \psi(x_1/a) + \frac{1}{1-a} \psi(x_2) \right) \\ \quad + \mathbb{1}_{\{x_1 > 0\}} \left(\frac{a}{1-a} \psi(x_2/a) + \frac{1}{1-a} \psi(x_1) \right) \\ \quad + \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 \leq 0\}} \left(\frac{a}{1-a} \psi(x_1/a) + \frac{a}{1-a} \psi(x_2/a) + 1 \right), & \text{if } ax_1 \geq x_2 \text{ and } ax_2 \geq x_1, \\ \frac{1+a}{1-a} \psi\left(\frac{x_1+x_2}{1+a}\right), & \text{if } ax_1 < x_2 \text{ and } ax_2 < x_1, \\ \frac{1}{1-a} \psi(x_2) + \frac{a}{1-a} \psi(x_1/a), & \text{if } ax_1 < x_2 \text{ and } ax_2 \geq x_1, \\ \frac{1}{1-a} \psi(x_1) + \frac{a}{1-a} \psi(x_2/a), & \text{if } ax_1 \geq x_2 \text{ and } ax_2 < x_1. \end{cases}$$

3.3. The Boundary of the Support of the Copula Density

The following proposition provides almost sure boundaries for X_2 , the second component of X in (4). The statements are directly clear from Figures 3 and 4. The boundaries

carry over to the boundaries for \mathbf{U} in the unit square after the map with the correct marginal survival functions (see also Figures 1 and 2).

Proposition 1. Under Assumption (CPD), the following inequalities hold almost surely for the component X_2 of $\mathbf{X} = (X_1, X_2)$ of Model (4).

- (1) If $a_{12} < 0$, $X_2 \geq x/a_{12}$ for all $x < 0$ and $X_2 \geq a_{21}x$ for all $x \geq 0$.
- (2) If $a_{12} = 0$, $X_2 \geq a_{21}x$ for all $x \geq 0$.
- (3) If $0 < a_{12} \leq 1$, $a_{21}x \leq X_2 \leq x/a_{12}$ for all $x \geq 0$.

3.4. Tail Dependence

In this section, we address the notion of tail dependence for the multivariate model as given in (4). Tail dependence is a copula property, so the coefficients of tail dependence only depend on the copula of (4). For the following result, note that a function $h : (0, \infty) \rightarrow \mathbb{R}$ is called *regularly varying* at $x_0 \in [0, \infty]$ of index $\alpha \in \mathbb{R}$ (denoted by $h \in \text{RV}_\alpha$) if

$$\lim_{x \rightarrow x_0} \frac{h(tx)}{h(x)} = t^\alpha, \quad t > 0.$$

Proposition 2. Let Assumption (CPD) hold and suppose that $a_{12} = a_{21} = a$. Furthermore, let the marginal distribution of X_1 and X_2 be denoted by $F = F_1 = F_2$.

- (1) Let $a \in (0, 1)$. If $\bar{F}^-(0) < \infty$, then $\lambda_L = 0$. If $\bar{F}^-(0) = \infty$ and $\psi' \in \text{RV}_{-\alpha}$ at ∞ , then

$$\lambda_L = 2 \frac{2^{-\alpha}(1+a)^\alpha - a^\alpha}{1 - a^\alpha}.$$

If $\psi' \in \text{RV}_{-\alpha}$ at 0, then

$$\lambda_U = 2 \frac{1 - 2^{-\alpha}(1+a)^\alpha}{1 - a^\alpha}.$$

- (2) Let $a \in (-1, 0)$. If $\bar{F}^-(0) < \infty$, then $\lambda_L = 0$. If $\bar{F}^-(0) = \infty$ and $\psi' \in \text{RV}_{-\alpha}$ at ∞ , then

$$\lambda_L = 2 \left(\frac{1+a}{2} \right)^\alpha.$$

Furthermore, $\lambda_U = 0$.

Figure 5 shows λ_L and λ_U of Part (1) of Proposition 2 as functions of the parameter a for various α .

Assessing λ_L and λ_U in the case $F_1 \neq F_2$ is more difficult. At least, the symmetric case $a_{12} = a_{21}$ can provide bounds. For example, if $\bar{F}_1^-(u) \leq \bar{F}_2^-(u)$ for all u sufficiently small, then (compare with the proof of Proposition 2)

$$\begin{aligned} \lambda_L &= \lim_{u \downarrow 0} \frac{C(u, u)}{u} = \lim_{u \downarrow 0} \frac{\mathbb{P}(X_1 > \bar{F}_1^-(u), X_2 > \bar{F}_2^-(u))}{\mathbb{P}(X_1 > \bar{F}_1^-(u))} \\ &\leq \lim_{u \downarrow 0} \frac{\mathbb{P}(X_1 > \bar{F}_1^-(u), X_2 > \bar{F}_1^-(u))}{\mathbb{P}(X_1 > \bar{F}_1^-(u))} \end{aligned} \tag{6}$$

and one can proceed as in the proof of Proposition 2 to evaluate (6); if $\bar{F}_1^-(u) \geq \bar{F}_2^-(u)$ for all u sufficiently small, then the inequality leads to a lower bound for λ_L .




Figure 5. λ_L and λ_U of Part (1) of Proposition 2 as functions of the parameter a for various α .

4. Randomization and Other Extensions

As shown above, tilting an Archimedean copula through a matrix A makes the model more flexible by introducing different kinds of asymmetry. It is also possible to modify tail dependence properties through A . However, some parameter choices reduce the support of the copula to strict subsets of the unit square. This might be a desired feature for certain applications (a classical example is the copula of (X_1, X_2) with $X_1 \leq X_2$ almost surely), but equally well unnatural in others. A possible remedy may be to randomize the tilting matrix A ; theoretical properties can then be derived by conditioning on A . While such *random-matrix-tilted Archimedean copulas* allow for very flexible models, inference is an open question.

As a first example, we revisit the right-hand side of Figure 1 and randomize the entry $a_{21} = 0.5$ of A with a standard uniform distribution. The result is shown on the left-hand side of Figure 6. Note that, although hardly visible here, the lower boundary as depicted on the right-hand side of Figure 1 still exists but the upper one is “washed out”. If we randomize both a_{12} and a_{21} , then both boundaries disappear. The right-hand side of Figure 6 shows such a case, where a_{12} and a_{21} are (independently) randomized by Beta(1/10, 1) and Beta(2, 1) distributions, respectively. Note that both of these randomizations satisfy Assumption (CPD).

As another example, let us consider the right-hand side of Figure 2 and randomize the entry $a_{21} = -0.05$ of A via a uniform distribution on $[-10, 0]$ (see the left-hand side of Figure 7) or via $a_{21} = -0.1 - \min\{E, 20\}$ with E being standard exponential (see the right-hand side of Figure 7). As above, both randomizations satisfy Assumption (CPD).

Another option not discussed here would be to make the randomized a_{12}, a_{21} dependent by another copula.

A different approach to tilting (2) by a matrix A that we do not explore is the following. Consider a random vector

$$\mathbf{X} = R\mathbf{A}\mathbf{U}, \tag{7}$$

where R is a radial part as in (1), \mathbf{U} is uniformly distributed on the ℓ^1 -sphere $\{\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d \mid |x_1| + \dots + |x_d| = 1\}$ independently of R , and A is a $d \times d$ -matrix. Then, one could consider the copula of the componentwise modulus $|\mathbf{X}|$ of \mathbf{X} . Figures 8 and 9 show the copulas of $|\mathbf{X}|$ for R and A as chosen in Figures 2 and 7, respectively, and \mathbf{X} given by (7). One observes that, for a deterministic tilting matrix A (see Figure 8), the construction does not seem to yield anything visually much different from a Gumbel copula (in distribution). We do not have proof of this fact but we have seen this behavior in all simulation examples.




Figure 6. Scatter plots of 1000 samples from random-matrix-tilted Archimedean copulas with randomized entry $a_{21} \sim U(0,1)$ (left) and independent $a_{12} \sim \text{Beta}(1/10,1)$, $a_{21} \sim \text{Beta}(2,1)$ (right) based on the example displayed on the right-hand side of Figure 1.




Figure 7. Similar to Figure 6 based on the right-hand side of Figure 2, with $a_{21} \sim U(-10,0)$ (left) and $a_{21} = -0.1 - \min\{E, 20\}$ with E being standard exponential (right).




Figure 8. Scatter plots of the copulas of $|X|$ corresponding to Figure 2.




Figure 9. Scatter plots of the copulas of $|X|$ corresponding to Figure 7.

The situation is more promising in the case of a randomized tiling matrix A (see Figure 9, especially the left-hand side).

5. Conclusions

Inspired by the construction of elliptical copulas, we present a new class of matrix-tilted Archimedean copulas, which generalize Archimedean copulas. Matrix-tilted Archimedean copulas are easy to simulate from and allow asymmetries to be modeled. In the bivariate case, the joint survival function and boundaries of the support of the multivariate model which gives rise to matrix-tilted Archimedean copulas can be given, and, in the case of equal margins and symmetric tilting matrix A , the tail dependence coefficients can be derived. Randomized tilting matrices A and the componentwise modulus of the model provide interesting extensions. Although their theoretical treatment is not straightforward, matrix-tilted Archimedean copulas are especially easy to incorporate into simulation-based applications.

The statistical estimation of matrix-tilted Archimedean copulas is an open problem for future research. A natural idea seems to be an iterative estimation procedure that alternates between finding an optimal Archimedean generator for the data given a specific tilting matrix and finding a best-fitting tilting matrix given the Archimedean generator of the previous step. In addition, random-matrix-tilted Archimedean copulas that do not limit the support to a strict subset of the unit hypercube may be of interest, too, as they may allow for likelihood optimization. Another avenue for future research is to introduce matrix tilting to generalizations of Archimedean copulas. One immediate generalization is to replace $\mathbf{U} \sim U(\Delta_d)$ in (4) by a Dirichlet distribution which would then lead to matrix-tilted Liouville copulas.

Author Contributions: Conceptualization, M.H. and J.F.Z.; methodology, M.H. and J.F.Z.; formal analysis, M.H. and J.F.Z.; investigation, M.H. and J.F.Z.; writing—original draft preparation, M.H. and J.F.Z.; writing—review and editing, M.H. and J.F.Z.; visualization, M.H. and J.F.Z. All authors have read and agreed to the published version of the manuscript.

Funding: The first author's research was funded by NSERC under Discovery Grant RGPIN-5010-2015.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

The following proposition provides the survival function of AU , a result which is used in the proof of Theorem 1 below.

Proposition A1. *Let $\mathbf{U} = (U_1, U_2)$ be uniformly distributed on Δ_2 . Under Assumption (CPD), we obtain that*

$$\mathbb{P}(a_{11}U_1 + a_{12}U_2 > y_1, a_{21}U_1 + a_{22}U_2 > y_2) = \begin{cases} \frac{1 - \max\{y_2, a_{21}\}}{1 - a_{21}} \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 < 1\}}, & \text{if } a_{12} = 1, a_{21} < 1, \\ \frac{1 - \max\{y_1, a_{12}\}}{1 - a_{12}} \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 < 1\}}, & \text{if } a_{12} < 1, a_{21} = 1, \\ \frac{1 - a_{12}a_{21} - (1 - a_{21})y_1 - (1 - a_{12})y_2}{(1 - a_{12})(1 - a_{21})} \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 < 1\}} \mathbb{1}_{\{(1 - a_{12})(1 - y_2) - (1 - a_{21})(y_1 - a_{12}) > 0\}} \\ - \frac{a_{12} - y_1}{1 - a_{12}} \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 < 1\}} - \frac{a_{21} - y_2}{1 - a_{21}} \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 \leq a_{21}\}}, & \text{if } a_{12} < 1, a_{21} < 1. \end{cases}$$

Proof. First, note that, for some $U \sim U(0, 1)$,

$$\begin{aligned} \mathbb{P}(U_1 + a_{12}U_2 > y_1, a_{21}U_1 + U_2 > y_2) &= \mathbb{P}(U + a_{12}(1 - U) > y_1, a_{21}U + 1 - U > y_2) \\ &= \mathbb{P}((1 - a_{12})U > y_1 - a_{12}, (a_{21} - 1)U > y_2 - 1). \end{aligned}$$

If $a_{12} = 1$ and $a_{21} < 1$, then

$$\begin{aligned} \mathbb{P}(U_1 + a_{12}U_2 > y_1, a_{21}U_1 + U_2 > y_2) &= \mathbb{1}_{\{y_1 < 1\}} \mathbb{P}\left(U < \frac{1 - y_2}{1 - a_{21}}\right) \\ &= \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 < 1\}} \min\left\{1, \frac{1 - y_2}{1 - a_{21}}\right\} \\ &= \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 < 1\}} \frac{1 - \max\{y_2, a_{21}\}}{1 - a_{21}}. \end{aligned}$$

Similarly, if $a_{12} < 1$ and $a_{21} = 1$, we can interchange the role of y_1, y_2 and a_{12}, a_{21} and obtain

$$\mathbb{P}(U_1 + a_{12}U_2 > y_1, a_{21}U_1 + U_2 > y_2) = \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 < 1\}} \frac{1 - \max\{y_1, a_{12}\}}{1 - a_{12}},$$

which is as stated in the claim. From now on, assume $a_{12} < 1$ and $a_{21} < 1$, and let $t = (1 - a_{12})(1 - y_2) - (1 - a_{21})(y_1 - a_{12})$. Then,

$$\begin{aligned} & \mathbb{P}(U_1 + a_{12}U_2 > y_1, a_{21}U_1 + U_2 > y_2) \\ &= \mathbb{P}\left(U > \frac{y_1 - a_{12}}{1 - a_{12}}, U < \frac{1 - y_2}{1 - a_{21}}\right) \\ &= \mathbb{1}_{\left\{\frac{y_1 - a_{12}}{1 - a_{12}} < 1\right\}} \mathbb{1}_{\left\{\frac{1 - y_2}{1 - a_{21}} > 0\right\}} \mathbb{1}_{\left\{\frac{y_1 - a_{12}}{1 - a_{12}} < \frac{1 - y_2}{1 - a_{21}}\right\}} \\ &\quad \cdot \left(\min\left\{1, \frac{1 - y_2}{1 - a_{21}}\right\} - \max\left\{0, \frac{y_1 - a_{12}}{1 - a_{12}}\right\}\right) \\ &= \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 < 1\}} \mathbb{1}_{\{t > 0\}} \left(\frac{1 - \max\{y_2, a_{21}\}}{1 - a_{21}} - \frac{\max\{y_1, a_{12}\} - a_{12}}{1 - a_{12}}\right). \end{aligned}$$

In the remaining part of the proof, we rewrite this expression to obtain the form as stated. Using $\mathbb{1}_{\{y_1 < 1\}} = \mathbb{1}_{\{y_1 \leq a_{12}\}} + \mathbb{1}_{\{a_{12} < y_1 < 1\}}$ and $\mathbb{1}_{\{y_2 < 1\}} = \mathbb{1}_{\{y_2 \leq a_{21}\}} + \mathbb{1}_{\{a_{21} < y_2 < 1\}}$, multiplying out the terms, and noting that $t > 0$ for all except the last term D (which easily follows from the bounds in the other indicators), we obtain that

$$\mathbb{P}(U_1 + a_{12}U_2 > y_1, a_{21}U_1 + U_2 > y_2) = A + B + C + D,$$

where

$$\begin{aligned} A &= \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 \leq a_{21}\}}, \\ B &= \frac{1 - y_1}{1 - a_{12}} \mathbb{1}_{\{a_{12} < y_1 < 1\}} \mathbb{1}_{\{y_2 \leq a_{21}\}}, \\ C &= \frac{1 - y_2}{1 - a_{21}} \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{a_{21} < y_2 < 1\}}, \\ D &= \mathbb{1}_{\{a_{12} < y_1 < 1\}} \mathbb{1}_{\{a_{21} < y_2 < 1\}} \mathbb{1}_{\{t > 0\}} \frac{1 - a_{12}a_{21} - (1 - a_{21})y_1 - (1 - a_{12})y_2}{(1 - a_{12})(1 - a_{21})}. \end{aligned}$$

Let

$$\begin{aligned} a &= \frac{1 - a_{12}a_{21}}{(1 - a_{12})(1 - a_{21})}, & b &= \frac{(1 - a_{21})y_1 + (1 - a_{12})y_2}{1 - a_{12}a_{21}}, \\ x &= \frac{(1 - a_{21})(a_{12} - y_1)}{1 - a_{12}a_{21}}, & \tilde{x} &= \frac{(1 - a_{12})(a_{21} - y_2)}{1 - a_{12}a_{21}}. \end{aligned}$$

Basic calculations show that

$$\begin{aligned} \frac{1 - y_1}{1 - a_{12}} &= a(1 - b - \tilde{x}) = a(1 - b) - a\tilde{x}, \\ \frac{1 - y_2}{1 - a_{21}} &= a(1 - b - x) = a(1 - b) - ax, \end{aligned}$$

so that $B = B_1 - B_2$ with

$$B_1 = a(1 - b) \mathbb{1}_{\{a_{12} < y_1 < 1\}} \mathbb{1}_{\{y_2 \leq a_{21}\}}, \quad B_2 = a\tilde{x} \mathbb{1}_{\{a_{12} < y_1 < 1\}} \mathbb{1}_{\{y_2 \leq a_{21}\}}.$$

In addition, $\mathbb{1}_{\{a_{21} < y_2 < 1\}} = \mathbb{1}_{\{y_2 < 1\}} - \mathbb{1}_{\{y_2 \leq a_{21}\}}$ implies that $C = C_1 - C_2$, with

$$C_1 = a(1 - b - x) \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 < 1\}}, \quad C_2 = a(1 - b - x) \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 \leq a_{21}\}}.$$

As for B , we can decompose C_1 into $C_{11} - C_{12}$ with

$$C_{11} = a(1 - b) \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 < 1\}}, \quad C_{12} = ax \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 < 1\}}.$$

Furthermore, C_2 can be written as $C_{22} + A$, where

$$C_{22} = \frac{a_{21} - y_2}{1 - a_{21}} \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 \leq a_{21}\}}.$$

We thus obtain that

$$\begin{aligned} & \mathbb{P}(U_1 + a_{12}U_2 > y_1, a_{21}U_1 + U_2 > y_2) \\ &= A + B + C + D = A + B_1 - B_2 + C_{11} - C_{12} - C_{22} - A + D \\ &= (B_1 + C_{11} + D) - C_{12} - (B_2 + C_{22}). \end{aligned}$$

It is easy to see that $B_2 + C_{22} = \frac{a_{21} - y_2}{1 - a_{21}} \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 \leq a_{21}\}}$. Finally, we can multiply with indicators that $t > 0$ and obtain that

$$\begin{aligned} & B_1 + C_{11} + D \\ &= a(1 - b) (\mathbb{1}_{\{a_{12} < y_1 < 1\}} \mathbb{1}_{\{y_2 \leq a_{21}\}} + \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 < 1\}} \\ & \quad + \mathbb{1}_{\{a_{12} < y_1 < 1\}} \mathbb{1}_{\{a_{21} < y_2 < 1\}} \mathbb{1}_{\{t > 0\}}) \\ &= a(1 - b) \mathbb{1}_{\{t > 0\}} (\mathbb{1}_{\{a_{12} < y_1 < 1\}} \mathbb{1}_{\{y_2 \leq a_{21}\}} + \mathbb{1}_{\{y_1 \leq a_{12}\}} \mathbb{1}_{\{y_2 < 1\}} \\ & \quad + \mathbb{1}_{\{a_{12} < y_1 < 1\}} \mathbb{1}_{\{a_{21} < y_2 < 1\}}). \end{aligned}$$

By first taking out $\mathbb{1}_{\{a_{12} < y_1 < 1\}}$ and consecutively pulling together the terms, we obtain that

$$B_1 + C_{11} + D = a(1 - b) \mathbb{1}_{\{y_1 < 1\}} \mathbb{1}_{\{y_2 < 1\}} \mathbb{1}_{\{t > 0\}},$$

which, by a basic calculation, leads to the claim as stated. \square

The following lemma is an auxiliary result, which is also used in the proof of Theorem 1.

Lemma A1. Let $\psi(t) = \mathcal{W}_2[F_R](t) = \int_{(t, \infty)} (1 - t/r) dF_R(r)$ and $-\infty \leq a \leq b \leq \infty$. Then,

$$\begin{aligned} & \int_{(t, \infty)} \mathbb{1}_{\{a < r \leq b\}} \left(1 - \frac{t}{r}\right) dF_R(r) \\ &= \left(1 - \frac{\min\{t, a\}}{a}\right) \bar{F}_R(\max\{0, a\}) + \min\left\{\frac{t}{\max\{0, a\}}, 1\right\} \psi(\max\{a, t\}) \\ & \quad - \left(1 - \frac{\min\{t, b\}}{b}\right) \bar{F}_R(\max\{0, b\}) - \min\left\{\frac{t}{\max\{0, b\}}, 1\right\} \psi(\max\{b, t\}), \end{aligned}$$

with the convention that $\min\{t/\max\{0, a\}, 1\} = 1$ for $a \leq 0$ (and similarly for the term involving b).

Proof. First, assume $0 < a \leq b$. Writing $\mathbb{1}_{\{r \leq b\}} = 1 - \mathbb{1}_{\{r > b\}}$, we obtain that

$$\begin{aligned} & \int_{(t, \infty)} \mathbb{1}_{\{a < r \leq b\}} \left(1 - \frac{t}{r}\right) dF_R(r) \\ &= \int_{(t, \infty)} \mathbb{1}_{\{r > a\}} \left(1 - \frac{t}{r}\right) dF_R(r) - \int_{(t, \infty)} \mathbb{1}_{\{r > \max\{a, b\}\}} \left(1 - \frac{t}{r}\right) dF_R(r), \end{aligned}$$

so it suffices to consider the first integral. For $t \geq a$, this gives $\psi(t)$. If $t < a$, we have

$$\begin{aligned} \int_{(a,\infty)} \left(1 - \frac{t}{r}\right) dF_R(r) &= \int_{(a,\infty)} \left(1 - \frac{t}{a}\right) + \frac{t}{a} \left(1 - \frac{a}{r}\right) dF_R(r) \\ &= \left(1 - \frac{t}{a}\right) \bar{F}_R(a) + \frac{t}{a} \psi(a). \end{aligned}$$

Hence,

$$\int_{(t,\infty)} \mathbb{1}_{\{r>a\}} \left(1 - \frac{t}{r}\right) dF_R(r) = \left(1 - \frac{\min\{t,a\}}{a}\right) \bar{F}_R(a) + \min\left\{\frac{t}{a}, 1\right\} \psi(\max\{a,t\}).$$

Applying the same idea with a replaced by $\max\{a,b\}$ and putting the pieces together, we obtain that

$$\begin{aligned} &\int_{(t,\infty)} \mathbb{1}_{\{a<r\leq b\}} \left(1 - \frac{t}{r}\right) dF_R(r) \\ &= \left(1 - \frac{\min\{t,a\}}{a}\right) \bar{F}_R(a) + \min\left\{\frac{t}{a}, 1\right\} \psi(\max\{a,t\}) \\ &\quad - \left(1 - \frac{\min\{t,b\}}{b}\right) \bar{F}_R(b) - \min\left\{\frac{t}{b}, 1\right\} \psi(\max\{b,t\}). \end{aligned}$$

Now, assume $a \leq 0 \leq b$. Then,

$$\int_{(t,\infty)} \mathbb{1}_{\{a<r\leq b\}} \left(1 - \frac{t}{r}\right) dF_R(r) = \int_{(t,\infty)} \mathbb{1}_{\{0<r\leq b\}} \left(1 - \frac{t}{r}\right) dF_R(r),$$

so that we can apply the formula for the case $0 < a \leq b$ with $a \searrow 0$ here. This implies that

$$\begin{aligned} &\int_{(t,\infty)} \mathbb{1}_{\{a<r\leq b\}} \left(1 - \frac{t}{r}\right) dF_R(r) \\ &= 0 \cdot 1 + 1 \cdot \psi(t) - \left(1 - \frac{\min\{t,b\}}{b}\right) \bar{F}_R(b) - \min\left\{\frac{t}{b}, 1\right\} \psi(\max\{b,t\}), \end{aligned}$$

which is of the form as stated in this case. Finally, if $a \leq b \leq 0$, then the integral is 0. Again, it is easy to see that the result as stated is indeed 0 in this case. \square

We can now address the proof of Theorem 1.

Proof of Theorem 1. We consider several cases.

Case 1: $a_{12} = 1$ and $a_{21} < 1$.

By Proposition A1,

$$\begin{aligned} &\mathbb{P}(RU_1 + a_{12}RU_2 > x_1, a_{21}RU_1 + RU_2 > x_2) \\ &= \int_{(0,\infty)} \mathbb{P}(U_1 + a_{12}U_2 > x_1/r, a_{21}U_1 + U_2 > x_2/r) dF_R(r) \\ &= \int_{(0,\infty)} \frac{1 - \max\left\{\frac{x_2}{r}, a_{21}\right\}}{1 - a_{21}} \mathbb{1}_{\{r>\max\{x_1,x_2\}\}} dF_R(r). \end{aligned}$$

The fraction in the integrand can be written as $\mathbb{1}_{\{ra_{21} \geq x_2\}} + (1 - \frac{x_2}{r}) \mathbb{1}_{\{ra_{21} < x_2\}} / (1 - a_{21})$ and thus the whole integral as $E + F / (1 - a_{21})$, with

$$\begin{aligned} E &= \int_{(0,\infty)} \mathbb{1}_{\{ra_{21} > x_2\}} \mathbb{1}_{\{r>\max\{x_1,x_2\}\}} dF_R(r), \\ F &= \int_{(0,\infty)} \mathbb{1}_{\{ra_{21} \leq x_2\}} \mathbb{1}_{\{r>\max\{x_1,x_2\}\}} \left(1 - \frac{x_2}{r}\right) dF_R(r). \end{aligned}$$

By distinguishing the cases $a_{21} < 0$, $a_{21} = 0$, and $a_{21} > 0$, we obtain that

$$E = \int_{(0,\infty)} \mathbb{1}_{\left\{r \leq \frac{x_2}{a_{21}}\right\}} \mathbb{1}_{\{r > \max\{x_1, x_2\}\}} dF_R(r) \mathbb{1}_{\{x_2 < 0\}} \mathbb{1}_{\{a_{21} < 0\}} + \int_{(0,\infty)} \mathbb{1}_{\{r > x_1\}} dF_R(r) \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} = 0\}} + \int_{(0,\infty)} \mathbb{1}_{\left\{r > \frac{x_2}{a_{21}}\right\}} \mathbb{1}_{\{r > x_1\}} dF_R(r) \mathbb{1}_{\{a_{21} > 0\}}.$$

Note that, if $x_2 \leq 0$, then $\mathbb{1}_{\{r > \max\{x_1, x_2\}\}} = \mathbb{1}_{\{r > \max\{0, x_1, x_2\}\}} = \mathbb{1}_{\{r > \max\{0, x_1\}\}}$. This implies that

$$E = \int_{(0,\infty)} \mathbb{1}_{\left\{\max\{0, x_1\} < r \leq \frac{x_2}{a_{21}}\right\}} dF_R(r) \mathbb{1}_{\{x_2 < 0\}} \mathbb{1}_{\{a_{21} < 0\}} + \bar{F}_R(\max\{0, x_1\}) \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} = 0\}} + \bar{F}_R\left(\max\left\{0, x_1, \frac{x_2}{a_{21}}\right\}\right) \mathbb{1}_{\{a_{21} > 0\}} = \left(\bar{F}_R(\max\{0, x_1\}) - \bar{F}_R\left(\frac{x_2}{a_{21}}\right)\right) \mathbb{1}_{\left\{\frac{x_2}{a_{21}} > \max\{0, x_1\}\right\}} \mathbb{1}_{\{x_2 < 0\}} \mathbb{1}_{\{a_{21} < 0\}} + \bar{F}_R(\max\{0, x_1\}) \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} = 0\}} + \bar{F}_R\left(\max\left\{0, x_1, \frac{x_2}{a_{21}}\right\}\right) \mathbb{1}_{\{a_{21} > 0\}}.$$

By splitting up F in a similar fashion, we obtain that

$$F = \int_{(0,\infty)} \mathbb{1}_{\left\{r > \max\left\{0, x_1, x_2, \frac{x_2}{a_{21}}\right\}\right\}} \left(1 - \frac{x_2}{r}\right) dF_R(r) \mathbb{1}_{\{a_{21} < 0\}} + \int_{(0,\infty)} \mathbb{1}_{\{r > \max\{x_1, x_2\}\}} \left(1 - \frac{x_2}{r}\right) dF_R(r) \mathbb{1}_{\{x_2 > 0\}} \mathbb{1}_{\{a_{21} = 0\}} + \int_{(0,\infty)} \mathbb{1}_{\left\{\max\{x_1, x_2\} < r \leq \frac{x_2}{a_{21}}\right\}} \left(1 - \frac{x_2}{r}\right) dF_R(r) \mathbb{1}_{\{x_2 > 0\}} \mathbb{1}_{\{a_{21} > 0\}}.$$

Distinguishing the cases of different sign of x_2 in the last integral leads to

$$F = \int_{(x_2, \infty)} \mathbb{1}_{\{r > x_1\}} \left(1 - \frac{x_2}{r}\right) dF_R(r) \mathbb{1}_{\{x_2 > 0\}} \mathbb{1}_{\{a_{21} < 0\}} + \int_{(x_2/a_{21}, \infty)} \mathbb{1}_{\{r > x_1\}} \left(1 - \frac{x_2}{r}\right) dF_R(r) \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} < 0\}} + \int_{(x_2, \infty)} \mathbb{1}_{\{r > x_1\}} \left(1 - \frac{x_2}{r}\right) dF_R(r) \mathbb{1}_{\{x_2 > 0\}} \mathbb{1}_{\{a_{21} = 0\}} + \int_{(x_2, \infty)} \mathbb{1}_{\left\{\max\{0, x_1\} < r \leq \frac{x_2}{a_{21}}\right\}} \left(1 - \frac{x_2}{r}\right) dF_R(r) \mathbb{1}_{\{x_2 > 0\}} \mathbb{1}_{\{a_{21} > 0\}} \cdot \mathbb{1}_{\left\{\max\{0, x_1\} < \frac{x_2}{a_{21}}\right\}}.$$

Note that the first and third integrals on the right-hand side of the last equation can be pulled together. In addition, note that the second integral can be written as

$$a_{21} \left(\int_{(x_2/a_{21}, \infty)} \left(\frac{1}{a_{21}} - 1\right) \mathbb{1}_{\{r > x_1\}} dF_R(r) + \int_{(x_2/a_{21}, \infty)} \mathbb{1}_{\{r > x_1\}} \left(1 - \frac{x_2/a_{21}}{r}\right) dF_R(r) \right) \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} < 0\}}.$$

Applying Lemma A1 leads to

$$F = \left(\left(1 - \frac{\min\{x_1, x_2\}}{x_1}\right) \bar{F}_R(\max\{0, x_1\}) + \min\left\{\frac{x_2}{\max\{0, x_1\}}, 1\right\} \right)$$

$$\begin{aligned}
 & \cdot \psi(\max\{x_1, x_2\}) \mathbb{1}_{\{x_2 > 0\}} \mathbb{1}_{\{a_{21} \leq 0\}} \\
 & + \left(\left(1 - \frac{\min\{\max\{0, x_1\}, x_2\}}{\max\{0, x_1\}} \right) \bar{F}_R(\max\{0, x_1\}) \right. \\
 & \quad + \min\left\{ \frac{x_2}{\max\{0, x_1\}}, 1 \right\} \psi(\max\{0, x_1, x_2\}) \\
 & \quad - \left(1 - \frac{\min\{x_2, x_2/a_{21}\}}{x_2/a_{21}} \right) \bar{F}_R(\max\{0, x_2/a_{21}\}) \\
 & \quad \left. - \min\left\{ \frac{x_2}{\max\{0, x_2/a_{21}\}}, 1 \right\} \psi\left(\max\left\{x_2, \frac{x_2}{a_{21}}\right\}\right) \right) \mathbb{1}_{\{x_2 > 0\}} \mathbb{1}_{\{a_{21} > 0\}} \\
 & \cdot \mathbb{1}_{\{\max\{0, x_1\} < \frac{x_2}{a_{21}}\}} \\
 & + a_{21} \left(\left(\frac{1}{a_{21}} - 1 \right) \bar{F}_R\left(\max\left\{x_1, \frac{x_2}{a_{21}}\right\}\right) + \left(1 - \frac{\min\{x_1, x_2/a_{21}\}}{x_1} \right) \right. \\
 & \quad \cdot \bar{F}_R(\max\{0, x_1\}) + \min\left\{ \frac{x_2/a_{21}}{\max\{0, x_1\}}, 1 \right\} \psi\left(\max\left\{x_1, \frac{x_2}{a_{21}}\right\}\right) \left. \right) \\
 & \cdot \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} < 0\}} \\
 = & \left(\left(1 - \frac{\min\{x_1, x_2\}}{x_1} \right) \bar{F}_R(\max\{0, x_1\}) + \min\left\{ \frac{x_2}{\max\{0, x_1\}}, 1 \right\} \right. \\
 & \cdot \psi(\max\{x_1, x_2\}) \left. \right) \mathbb{1}_{\{x_2 > 0\}} \\
 & - \left((1 - a_{21}) \bar{F}_R(x_2/a_{21}) + a_{21} \psi\left(\frac{x_2}{a_{21}}\right) \right) \mathbb{1}_{\{x_2 > 0\}} \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_1 < \frac{x_2}{a_{21}}\}} \\
 & + a_{21} \left(\left(\frac{1}{a_{21}} - 1 \right) \bar{F}_R\left(\max\left\{x_1, \frac{x_2}{a_{21}}\right\}\right) + \left(1 - \frac{\min\{x_1, x_2/a_{21}\}}{x_1} \right) \right. \\
 & \quad \cdot \bar{F}_R(\max\{0, x_1\}) + \min\left\{ \frac{x_2/a_{21}}{\max\{0, x_1\}}, 1 \right\} \psi\left(\max\left\{x_1, \frac{x_2}{a_{21}}\right\}\right) \left. \right) \\
 & \cdot \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} < 0\}}.
 \end{aligned}$$

Recalling that the result is $E + F/(1 - a_{21})$, we obtain the form as claimed.

Case 2: $a_{12} < 1$ and $a_{21} = 1$.

This case directly follows from Case 1 by interchanging a_{12} and a_{21} , as well as x_1 and x_2 .

Case 3: $a_{12} < 1$ and $a_{21} < 1$.

Since $\mathbb{P}(RU_1 + a_{12}RU_2 > x_1, a_{21}RU_1 + RU_2 > x_2) = \int_{(0, \infty)} \mathbb{P}(U_1 + a_{12}U_2 > x_1/r, a_{21}U_1 + U_2 > x_2/r) dF_R(r)$, let us first consider the integrand. By Proposition A1, we have that

$$\mathbb{P}(U_1 + a_{12}U_2 > x_1/r, a_{21}U_1 + U_2 > x_2/r) = A - B - C,$$

where

$$\begin{aligned}
 A &= \frac{1 - a_{12}a_{21} - (1 - a_{21})\frac{x_1}{r} - (1 - a_{12})\frac{x_2}{r}}{(1 - a_{12})(1 - a_{21})} \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \\
 & \cdot \mathbb{1}_{\left\{ \frac{x_1(1 - a_{21}) + x_2(1 - a_{12})}{1 - a_{12}a_{21}} < r \right\}}, \\
 B &= \frac{a_{12} - \frac{x_1}{r}}{1 - a_{12}} \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}}, & C &= \frac{a_{21} - \frac{x_2}{r}}{1 - a_{21}} \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}}.
 \end{aligned}$$

It is straightforward to check that

$$a_{21}x_1 \geq x_2 \iff \frac{x_1(1 - a_{21}) + x_2(1 - a_{12})}{1 - a_{12}a_{21}} \leq x_1, \tag{A1}$$

We now consider four different cases.

Case 3.1: $a_{21}x_1 \geq x_2$ and $a_{12}x_2 \geq x_1$. Condition (A1) and the last equivalence imply that

$$\mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\left\{\frac{x_1(1 - a_{21}) + x_2(1 - a_{12})}{1 - a_{12}a_{21}} < r\right\}} = \mathbb{1}_{\{x_1 < r\}},$$

so that

$$A = \frac{1 - a_{12}a_{21} - (1 - a_{21})\frac{x_1}{r} - (1 - a_{12})\frac{x_2}{r}}{(1 - a_{12})(1 - a_{21})} \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}}.$$

This can be rewritten as

$$\begin{aligned} A &= \left(1 + \frac{1}{1 - a_{12}}\left(a_{12} - \frac{x_1}{r}\right) + \frac{1}{1 - a_{21}}\left(a_{21} - \frac{x_2}{r}\right)\right) \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \\ &= \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \\ &\quad + \frac{1}{1 - a_{12}}\left(a_{12} - \frac{x_1}{r}\right) \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} \geq 0\}} \\ &\quad + \frac{1}{1 - a_{12}}\left(a_{12} - \frac{x_1}{r}\right) \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} < 0\}} \\ &\quad + \frac{1}{1 - a_{21}}\left(a_{21} - \frac{x_2}{r}\right) \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{21} \geq 0\}} \\ &\quad + \frac{1}{1 - a_{21}}\left(a_{21} - \frac{x_2}{r}\right) \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{21} < 0\}}. \end{aligned}$$

Furthermore, $x_1 \leq a_{12}x_2$ implies that

$$\mathbb{1}_{\{a_{12} \geq 0\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{x_1 < ra_{12}\}} = \mathbb{1}_{\{a_{12} \geq 0\}} \mathbb{1}_{\{x_2 < r\}},$$

so that

$$\begin{aligned} B &= \frac{1}{1 - a_{12}}\left(a_{12} - \frac{x_1}{r}\right) (\mathbb{1}_{\{a_{12} \geq 0\}} \mathbb{1}_{\{x_1 \leq ra_{12}\}} + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1 \leq ra_{12}\}}) \mathbb{1}_{\{x_2 < r\}} \\ &= \frac{1}{1 - a_{12}}\left(a_{12} - \frac{x_1}{r}\right) (\mathbb{1}_{\{a_{12} \geq 0\}} \mathbb{1}_{\{x_2 < r\}} + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_2 < r \leq \frac{x_1}{a_{12}}\}}). \end{aligned}$$

Similarly, we obtain that

$$C = \frac{1}{1 - a_{21}}\left(a_{21} - \frac{x_2}{r}\right) (\mathbb{1}_{\{a_{21} \geq 0\}} \mathbb{1}_{\{x_1 < r\}} + \mathbb{1}_{\{a_{21} < 0\}} \mathbb{1}_{\{x_1 < r \leq \frac{x_2}{a_{21}}\}}).$$

Summarizing the terms leads to

$$\begin{aligned} A - B - C &= \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \\ &\quad + \mathbb{1}_{\{a_{12} \geq 0\}} \frac{1}{1 - a_{12}}\left(a_{12} - \frac{x_1}{r}\right) (\mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} - \mathbb{1}_{\{x_2 < r\}}) \\ &\quad + \mathbb{1}_{\{a_{12} < 0\}} \frac{1}{1 - a_{12}}\left(a_{12} - \frac{x_1}{r}\right) (\mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} - \mathbb{1}_{\{x_2 < r \leq \frac{x_1}{a_{12}}\}}) \\ &\quad + \mathbb{1}_{\{a_{21} \geq 0\}} \frac{1}{1 - a_{21}}\left(a_{21} - \frac{x_2}{r}\right) (\mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} - \mathbb{1}_{\{x_1 < r\}}) \\ &\quad + \mathbb{1}_{\{a_{21} < 0\}} \frac{1}{1 - a_{21}}\left(a_{21} - \frac{x_2}{r}\right) (\mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} - \mathbb{1}_{\{x_1 < r \leq \frac{x_2}{a_{21}}\}}). \end{aligned}$$

Since $\mathbb{1}_{\{x_2 < r \leq \frac{x_1}{a_{12}}\}} = \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}}$ and $\mathbb{1}_{\{x_1 < r \leq \frac{x_2}{a_{21}}\}} = \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{r \leq \frac{x_2}{a_{21}}\}}$, we have

$$\begin{aligned} A - B - C &= \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \\ &\quad + \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r} \right) \mathbb{1}_{\{x_2 < r\}} \left(\mathbb{1}_{\{a_{12} < 0\}} \left(\mathbb{1}_{\{x_1 < r\}} \right. \right. \\ &\quad \left. \left. - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) - \mathbb{1}_{\{a_{12} \geq 0\}} \mathbb{1}_{\{x_1 \geq r\}} \right) \\ &\quad + \frac{1}{1 - a_{21}} \left(a_{21} - \frac{x_2}{r} \right) \mathbb{1}_{\{x_1 < r\}} \left(\mathbb{1}_{\{a_{21} < 0\}} \left(\mathbb{1}_{\{x_2 < r\}} \right. \right. \\ &\quad \left. \left. - \mathbb{1}_{\{r \leq \frac{x_2}{a_{21}}\}} \right) - \mathbb{1}_{\{a_{21} \geq 0\}} \mathbb{1}_{\{x_2 \geq r\}} \right). \end{aligned}$$

Consider the case $a_{12} \geq 0$. If $a_{12} = 0$, then $x_1 \leq a_{12}x_2 = 0$, so that $\mathbb{1}_{\{x_1 \geq r\}} = 0$. If $a_{12} > 0$ and $x_1 > 0$, then our above assumptions imply that $x_1 > a_{12}a_{21}x_1 \geq a_{12}x_2 \geq x_1$, a contradiction. Thus, $x_1 \leq 0$ if $a_{12} \geq 0$. Similarly, $x_2 \leq 0$ if $a_{21} \geq 0$. This implies that

$$\begin{aligned} A - B - C &= \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 < r\}} \\ &\quad + \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r} \right) \mathbb{1}_{\{a_{12} < 0\}} \left(\mathbb{1}_{\{x_1 < r\}} - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) \mathbb{1}_{\{x_2 < r\}} \\ &\quad + \frac{1}{1 - a_{21}} \left(a_{21} - \frac{x_2}{r} \right) \mathbb{1}_{\{a_{21} < 0\}} \left(\mathbb{1}_{\{x_2 < r\}} - \mathbb{1}_{\{r \leq \frac{x_2}{a_{21}}\}} \right) \mathbb{1}_{\{x_1 < r\}}. \end{aligned}$$

For $a_{12} < 0$, we have $x_1 \leq a_{12}x_2 \leq 0$ if $x_2 \geq 0$. Thus, for $a_{12} < 0$, we have that

$$\begin{aligned} \left(\mathbb{1}_{\{x_1 < r\}} - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) \mathbb{1}_{\{x_2 < r\}} &= \left(\mathbb{1}_{\{x_1 < r\}} - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{x_1 > 0\}} \mathbb{1}_{\{x_2 \geq 0\}} \\ &\quad + \left(\mathbb{1}_{\{x_1 < r\}} - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 \geq 0\}} \\ &\quad + \left(\mathbb{1}_{\{x_1 < r\}} - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{x_1 > 0\}} \mathbb{1}_{\{x_2 < 0\}} \\ &\quad + \left(\mathbb{1}_{\{x_1 < r\}} - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 < 0\}} \\ &= 0 + \left(1 - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) \cdot 1 \cdot 1 \cdot \mathbb{1}_{\{x_2 \geq 0\}} \\ &\quad + \left(\mathbb{1}_{\{x_1 < r\}} - 0 \right) \cdot 1 \cdot \mathbb{1}_{\{x_1 > 0\}} \mathbb{1}_{\{x_2 < 0\}} \\ &\quad + \left(1 - \mathbb{1}_{\{r \leq \frac{x_1}{a_{12}}\}} \right) \cdot 1 \cdot \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 < 0\}} \\ &= \mathbb{1}_{\{r > \frac{x_1}{a_{12}}\}} \mathbb{1}_{\{x_2 \geq 0\}} + \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_1 > 0\}} \mathbb{1}_{\{x_2 < 0\}} \\ &\quad + \mathbb{1}_{\{r > \frac{x_1}{a_{12}}\}} \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 < 0\}}. \end{aligned}$$

Similarly,

$$\begin{aligned} &\left(\mathbb{1}_{\{x_2 < r\}} - \mathbb{1}_{\{r \leq \frac{x_2}{a_{21}}\}} \right) \mathbb{1}_{\{x_1 < r\}} \\ &= \mathbb{1}_{\{r > \frac{x_2}{a_{21}}\}} \mathbb{1}_{\{x_1 \geq 0\}} + \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{x_1 < 0\}} \mathbb{1}_{\{x_2 > 0\}} + \mathbb{1}_{\{r > \frac{x_2}{a_{21}}\}} \mathbb{1}_{\{x_1 < 0\}} \mathbb{1}_{\{x_2 \leq 0\}}. \end{aligned}$$

Integrating the terms now leads to

$$\begin{aligned} &\int_{(0,\infty)} (A - B - C) dF_R(r) \\ &= \int_{(0,\infty)} \mathbb{1}_{\{r > \max\{x_1, x_2, 0\}\}} dF_R(r) \\ &\quad + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_2 \geq 0\}} \frac{1}{1 - a_{12}} \int_{(0,\infty)} \mathbb{1}_{\{r > \frac{x_1}{a_{12}}\}} \left(a_{12} - \frac{x_1}{r} \right) dF_R(r) \end{aligned}$$

$$\begin{aligned}
& + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1 > 0\}} \mathbb{1}_{\{x_2 < 0\}} \frac{1}{1 - a_{12}} \int_{(0, \infty)} \mathbb{1}_{\{r > x_1\}} \left(a_{12} - \frac{x_1}{r} \right) dF_R(r) \\
& + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 < 0\}} \frac{1}{1 - a_{12}} \int_{(0, \infty)} \mathbb{1}_{\{r > \frac{x_1}{a_{12}}\}} \left(a_{12} - \frac{x_1}{r} \right) dF_R(r) \\
& + \mathbb{1}_{\{a_{21} < 0\}} \mathbb{1}_{\{x_1 \geq 0\}} \frac{1}{1 - a_{21}} \int_{(0, \infty)} \mathbb{1}_{\{r > \frac{x_2}{a_{21}}\}} \left(a_{21} - \frac{x_2}{r} \right) dF_R(r) \\
& + \mathbb{1}_{\{a_{21} < 0\}} \mathbb{1}_{\{x_1 < 0\}} \mathbb{1}_{\{x_2 > 0\}} \frac{1}{1 - a_{21}} \int_{(0, \infty)} \mathbb{1}_{\{r > x_2\}} \left(a_{21} - \frac{x_2}{r} \right) dF_R(r) \\
& + \mathbb{1}_{\{a_{21} < 0\}} \mathbb{1}_{\{x_1 < 0\}} \mathbb{1}_{\{x_2 \leq 0\}} \frac{1}{1 - a_{21}} \int_{(0, \infty)} \mathbb{1}_{\{r > \frac{x_2}{a_{21}}\}} \left(a_{21} - \frac{x_2}{r} \right) dF_R(r) \\
= & \bar{F}_R(\max\{x_1, x_2, 0\}) + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_2 \geq 0\}} \frac{a_{12}}{1 - a_{12}} \psi(x_1/a_{12}) \\
& + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1 > 0\}} \mathbb{1}_{\{x_2 < 0\}} \left(\frac{1}{1 - a_{12}} \psi(x_1) - \bar{F}_R(\max\{x_1, 0\}) \right) \\
& + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 < 0\}} \frac{a_{12}}{1 - a_{12}} \psi(x_1/a_{12}) \\
& + \mathbb{1}_{\{a_{21} < 0\}} \mathbb{1}_{\{x_1 \geq 0\}} \frac{a_{21}}{1 - a_{21}} \psi(x_2/a_{21}) \\
& + \mathbb{1}_{\{a_{21} < 0\}} \mathbb{1}_{\{x_1 < 0\}} \mathbb{1}_{\{x_2 > 0\}} \left(\frac{1}{1 - a_{21}} \psi(x_2) - \bar{F}_R(\max\{x_2, 0\}) \right) \\
& + \mathbb{1}_{\{a_{21} < 0\}} \mathbb{1}_{\{x_1 < 0\}} \mathbb{1}_{\{x_2 \leq 0\}} \frac{a_{21}}{1 - a_{21}} \psi(x_2/a_{21}).
\end{aligned}$$

Case 3.2: $a_{21}x_1 < x_2$ and $a_{12}x_2 < x_1$.

It is straightforward to check that, in this case,

$$\frac{x_1(1 - a_{21}) + x_2(1 - a_{12})}{1 - a_{12}a_{21}} > \max\{x_1, x_2\}$$

so that

$$\begin{aligned}
A &= \frac{1 - a_{12}a_{21} - (1 - a_{21})\frac{x_1}{r} - (1 - a_{12})\frac{x_2}{r}}{(1 - a_{12})(1 - a_{21})} \mathbb{1}_{\left\{ \frac{x_1(1 - a_{21}) + x_2(1 - a_{12})}{1 - a_{12}a_{21}} < r \right\}} \\
&= \frac{1 - a_{12}a_{21}}{(1 - a_{12})(1 - a_{21})} \left(1 - \frac{(1 - a_{21})x_1 + (1 - a_{12})x_2}{r(1 - a_{12}a_{21})} \right) \mathbb{1}_{\left\{ \frac{x_1(1 - a_{21}) + x_2(1 - a_{12})}{1 - a_{12}a_{21}} < r \right\}}.
\end{aligned}$$

For B , note that $a_{12} < 0$ and $x_1 \leq ra_{12}$ imply that $x_2 > \frac{x_1}{a_{12}} \geq r$. For $a_{12} = 0$, $x_1 > a_{12}x_2 = 0$. Furthermore, $a_{12} > 0$, and $x_1 \leq ra_{12}$ imply that $x_2 < \frac{x_1}{a_{12}} \leq r$. Therefore,

$$\begin{aligned}
\mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} < 0\}} &= 0, \\
\mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} = 0\}} &= \mathbb{1}_{\{x_1 \leq 0\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} = 0\}} = 0, \\
\mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} > 0\}} &= \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{a_{12} > 0\}}.
\end{aligned}$$

Thus,

$$\begin{aligned}
B &= \frac{a_{12} - \frac{x_1}{r}}{1 - a_{12}} \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} = \frac{a_{12} - \frac{x_1}{r}}{1 - a_{12}} (\mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} < 0\}} \\
&+ \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} = 0\}} + \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{a_{12} > 0\}}) \\
&= \frac{a_{12} - \frac{x_1}{r}}{1 - a_{12}} \mathbb{1}_{\left\{ \frac{x_1}{a_{12}} \leq r \right\}} \mathbb{1}_{\{a_{12} > 0\}}.
\end{aligned}$$

Similarly,

$$C = \frac{a_{21} - \frac{x_2}{r}}{1 - a_{21}} \mathbb{1}_{\{\frac{x_2}{a_{21}} \leq r\}} \mathbb{1}_{\{a_{21} > 0\}}.$$

Note that, in our case here, $(1 - a_{21})x_1 + (1 - a_{12})x_2 \geq 0$ because

$$0 > (1 - a_{21})x_1 + (1 - a_{12})x_2 \iff x_1 + x_2 < x_1 a_{21} + x_2 a_{12},$$

but the right-hand side is less than or equal to $x_1 + x_2$. In addition, if $a_{12} > 0$, we have that $x_1 \geq 0$, because, otherwise ($x_1 < 0$), we have $x_2 < \frac{x_1}{a_{12}} < 0$, which then leads to $(1 - a_{21})x_1 + (1 - a_{12})x_2 < 0$, a contradiction (note that $a_{12}a_{21} < 1$). This implies that

$$\begin{aligned} \int_{(0,\infty)} (A - B - C) dF_R(r) &= \frac{1 - a_{12}a_{21}}{(1 - a_{12})(1 - a_{21})} \psi\left(\frac{(1 - a_{21})x_1 + (1 - a_{12})x_2}{1 - a_{12}a_{21}}\right) \\ &\quad - \frac{a_{12}}{1 - a_{12}} \mathbb{1}_{\{a_{12} > 0\}} \mathbb{1}_{\{x_1 \geq 0\}} \psi\left(\frac{x_1}{a_{12}}\right) \\ &\quad - \frac{a_{21}}{1 - a_{21}} \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_2 \geq 0\}} \psi\left(\frac{x_2}{a_{21}}\right). \end{aligned}$$

Case 3.3: $a_{21}x_1 < x_2$ and $a_{12}x_2 \geq x_1$.

Let $x_2 < r$. If $a_{12} \geq 0$, then $x_1 \leq a_{12}x_2 < a_{12}r \leq r$, so $\mathbb{1}_{\{x_1 < r\}} = 1$. If $a_{12} < 0$, then $a_{21}x_1 < x_2 \leq x_1/a_{12}$, thus $(a_{21} - 1/a_{12})x_1 \leq 0$ and therefore $(a_{12}a_{21} - 1)x_1 \geq 0$. This implies that $x_1 \leq 0$ and thus that $\mathbb{1}_{\{x_1 < r\}} = 1$, so this indicator drops out of the term A . In addition, note that $a_{12}x_2 \geq x_1$ implies $x_1(1 - a_{21}) + x_2(1 - a_{12}) \leq x_2(1 - a_{12}a_{21}) < r(1 - a_{12}a_{21})$ for $x_2 < r$. Hence, the corresponding indicator in the term A is also always one. We therefore obtain that

$$A = \frac{1 - a_{12}a_{21} - (1 - a_{21})\frac{x_1}{r} - (1 - a_{12})\frac{x_2}{r}}{(1 - a_{12})(1 - a_{21})} \mathbb{1}_{\{x_2 < r\}}.$$

By considering the cases of different signs for a_{12} , we can write the term B as

$$\begin{aligned} B &= \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r}\right) \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} \\ &= \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r}\right) \left(\mathbb{1}_{\{a_{12} \geq 0\}} \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} + \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}}\right). \end{aligned}$$

If $x_2 < r$ and $a_{12} \geq 0$, then $x_1 \leq a_{12}x_2 < a_{12}r$, so $\mathbb{1}_{\{a_{12} \geq 0\}} \mathbb{1}_{\{x_1 \leq ra_{12}\}} \mathbb{1}_{\{x_2 < r\}} = \mathbb{1}_{\{a_{12} \geq 0\}} \mathbb{1}_{\{x_2 < r\}} = (1 - \mathbb{1}_{\{a_{12} < 0\}}) \mathbb{1}_{\{x_2 < r\}}$. By summarizing the indicators, this yields

$$B = \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r}\right) \left(\mathbb{1}_{\{x_2 < r\}} - \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_2 < r\}} \mathbb{1}_{\{x_1 > ra_{12}\}}\right).$$

Since $x_2 \leq x_1/a_{12} < r$ if $a_{12} < 0$ and $ra_{12} < x_1$, it follows that

$$B = \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r}\right) \left(\mathbb{1}_{\{x_2 < r\}} - \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1/a_{12} < r\}}\right).$$

Similarly, for C ,

$$C = \frac{1}{1 - a_{21}} \left(a_{21} - \frac{x_2}{r}\right) \left(\mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}} + \mathbb{1}_{\{a_{21} \leq 0\}} \mathbb{1}_{\{x_1 < r\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}}\right).$$

Note that the indicator function $\mathbb{1}_{\{x_1 < r\}}$ drops out, since, if $a_{21} > 0$ and $x_2 \leq ra_{21}$, then $x_1 < x_2/a_{21} \leq r$. The second summand of indicators completely vanishes, since, if

$a_{21} < 0$ and $x_2 \leq ra_{21}$, then $a_{21}x_1 < x_2 \leq ra_{21}$ implies $x_1 > r$. If $a_{21} = 0$, the contradiction $0 < x_2 \leq 0$ follows. We thus obtain that

$$C = \frac{1}{1 - a_{21}} \left(a_{21} - \frac{x_2}{r} \right) \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}}.$$

Pulling the terms together, we obtain that

$$\begin{aligned} A - B - C &= \frac{1}{1 - a_{21}} \left(1 - \frac{x_2}{r} \right) \mathbb{1}_{\{x_2 < r\}} + \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r} \right) \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1/a_{12} < r\}} \\ &\quad - \frac{1}{1 - a_{21}} \left(a_{21} - \frac{x_2}{r} \right) \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}}. \end{aligned}$$

Distinguishing the cases $x_2 \leq 0$ and $x_2 > 0$, we obtain that

$$\begin{aligned} A - B - C &= \frac{1}{1 - a_{21}} \left(\left(1 - \frac{x_2}{r} \right) \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{x_2 < r\}} - \left(a_{21} - \frac{x_2}{r} \right) \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}} \right. \\ &\quad \left. + \mathbb{1}_{\{x_2 > 0\}} \left(\left(1 - \frac{x_2}{r} \right) \mathbb{1}_{\{x_2 < r\}} - \left(a_{21} - \frac{x_2}{r} \right) \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}} \right) \right) \\ &\quad + \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r} \right) \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\left\{ \frac{x_1}{a_{12}} < r \right\}}. \end{aligned}$$

Note that $\mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{x_2 < r\}} = \mathbb{1}_{\{x_2 \leq 0\}}$ and $\mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}} = \mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} > 0\}}$ $= \mathbb{1}_{\{x_2 \leq 0\}} (1 - \mathbb{1}_{\{a_{21} \leq 0\}})$. We show next that $\mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} \leq 0\}} = 0$. To this end, let $x_2 \leq 0$. If $a_{21} < 0$, then $a_{21}x_1 < x_2 \leq 0$ implies that $x_1 > 0$ and thus $a_{12}x_2 \geq x_1 > 0$. If $x_2 = 0$, then a contradiction follows immediately. Otherwise, we see that $a_{12} < 0$. This yields $a_{12}a_{21} > 1$ in contradiction to the assumption that $1 - a_{12}a_{21} > 0$. If $a_{21} = 0$, then $0 = a_{21}x_1 < x_2 \leq 0$ yields a contradiction immediately. Therefore, $\mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} \leq 0\}} = 0$ and hence $\mathbb{1}_{\{x_2 \leq 0\}} \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}} = \mathbb{1}_{\{x_2 \leq 0\}}$. Similarly, if $a_{12} < 0$ and $x_1/a_{12} < r$, we show that $x_1 < 0$. To this end, let $x_1 \geq 0$. Then, $a_{12}x_2 \geq x_1 \geq 0$, so that $x_2 \leq 0$, and thus $a_{21}x_1 < x_2 \leq 0$. The case $x_1 = 0$ directly leads to a contradiction. If $x_1 > 0$, then $a_{21} < 0$, which implies that $a_{12}x_2 \geq x_1 > x_2/a_{21}$. As above, this leads to a contradiction to the assumption that $1 - a_{12}a_{21} > 0$. Thus, $\mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\left\{ \frac{x_1}{a_{12}} < r \right\}} = \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\left\{ \frac{x_1}{a_{12}} < r \right\}} \mathbb{1}_{\{x_1 < 0\}}$.

Plugging in these relations and summarizing the terms, we obtain that

$$\begin{aligned} A - B - C &= \mathbb{1}_{\{x_2 \leq 0\}} + \frac{\mathbb{1}_{\{x_2 > 0\}}}{1 - a_{21}} \left(\left(1 - \frac{x_2}{r} \right) \mathbb{1}_{\{r > x_2\}} - \left(a_{21} - \frac{x_2}{r} \right) \mathbb{1}_{\{a_{21} > 0\}} \mathbb{1}_{\{x_2 \leq ra_{21}\}} \right) \\ &\quad + \frac{1}{1 - a_{12}} \left(a_{12} - \frac{x_1}{r} \right) \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\left\{ \frac{x_1}{a_{12}} < r \right\}} \mathbb{1}_{\{x_1 < 0\}}. \end{aligned}$$

Integrating the terms now leads to

$$\begin{aligned} &\int_{(0, \infty)} (A - B - C) dF_R(r) \\ &= \mathbb{1}_{\{x_2 \leq 0\}} + \left(\frac{1}{1 - a_{21}} \psi(x_2) - \frac{a_{21}}{1 - a_{21}} \psi\left(\frac{x_2}{a_{21}}\right) \mathbb{1}_{\{a_{21} > 0\}} \right) \mathbb{1}_{\{x_2 > 0\}} \\ &\quad + \frac{a_{12}}{1 - a_{12}} \psi\left(\frac{x_1}{a_{12}}\right) \mathbb{1}_{\{a_{12} < 0\}} \mathbb{1}_{\{x_1 < 0\}}. \end{aligned}$$

Case 3.4: $a_{21}x_1 \geq x_2$ and $a_{12}x_2 < x_1$.

Interchanging the roles of a_{21}, a_{12} and x_1, x_2 , we obtain that

$$\begin{aligned} & \int_{(0,\infty)} (A - B - C) dF_R(r) \\ &= \mathbb{1}_{\{x_1 \leq 0\}} + \left(\frac{1}{1 - a_{12}} \psi(x_1) - \frac{a_{12}}{1 - a_{12}} \psi\left(\frac{x_1}{a_{12}}\right) \mathbb{1}_{\{a_{12} > 0\}} \right) \mathbb{1}_{\{x_1 > 0\}} \\ & \quad + \frac{a_{21}}{1 - a_{21}} \psi\left(\frac{x_2}{a_{21}}\right) \mathbb{1}_{\{a_{21} < 0\}} \mathbb{1}_{\{x_2 < 0\}}. \end{aligned}$$

□

Finally, here is the proof of Proposition 2.

Proof of Proposition 2. (1) Recall that $\mathbf{U} \stackrel{d}{=} (\bar{F}_1(X_1), \bar{F}_2(X_2))$ for $\mathbf{X} = (X_1, X_2)$ as in (4). Then,

$$\frac{C(u, u)}{u} = \frac{\mathbb{P}(\bar{F}_1(X_1) \leq u, \bar{F}_2(X_2) \leq u)}{\mathbb{P}(\bar{F}_1(X_1) \leq u)} = \frac{\mathbb{P}(X_1 > \bar{F}_1^-(u), X_2 > \bar{F}_2^-(u))}{\mathbb{P}(X_1 > \bar{F}_1^-(u))}.$$

Under the given assumptions, note that the distribution of (X_1, X_2) is supported in the first quadrant, the marginal survival functions are equal (denoted by \bar{F}), and, for $x \geq 0$, we have by Corollary 1 that

$$\mathbb{P}(X_1 > x, X_2 > x) = \frac{2}{1 - a} \left(\frac{1 + a}{2} \psi\left(\frac{2}{1 + a}x\right) - a\psi\left(\frac{x}{a}\right) \right), \tag{A2}$$

$$\bar{F}(x) = \mathbb{P}(X_1 > x) = \mathbb{P}(X_2 > x) = \frac{1}{1 - a} \left(\psi(x) - a\psi\left(\frac{x}{a}\right) \right). \tag{A3}$$

Therefore, we obtain that

$$\lambda_L = \lim_{u \downarrow 0} \frac{C(u, u)}{u} = 2 \lim_{x \uparrow \bar{F}^-(0)} \frac{\frac{1 + a}{2} \psi\left(\frac{2}{1 + a}x\right) - a\psi\left(\frac{x}{a}\right)}{\psi(x) - a\psi\left(\frac{x}{a}\right)}. \tag{A4}$$

Let $\bar{F}^-(0) < \infty$.

Since $(1 + a)/2 \in (1/2, 1)$ and ψ is non-increasing, $\frac{1+a}{2}\psi\left(\frac{2}{1+a}x\right) < \psi(x)$ for all $x \in [0, \bar{F}^-(0))$. Furthermore, there exists an $\tilde{x} \in [0, \bar{F}^-(0))$ such that the numerator of (A4) is zero but the denominator is greater than zero for all $x \in [\tilde{x}, \bar{F}^-(0))$. Hence $\lambda_L = 0$.

If $\bar{F}^-(0) = \infty$, applying l'Hôpital's Rule leads to

$$\lambda_L = 2 \lim_{x \uparrow \infty} \frac{\psi'\left(\frac{2x}{1 + a}\right) - \psi'\left(\frac{x}{a}\right)}{\psi'(x) - \psi'\left(\frac{x}{a}\right)}.$$

Dividing by $\psi'(x)$ and using regular variation, we obtain that

$$\lambda_L = 2 \lim_{x \uparrow \infty} \frac{\frac{\psi'(x \cdot 2/(1 + a))}{\psi'(x)} - \frac{\psi'(x/a)}{\psi'(x)}}{1 - \frac{\psi'(x/a)}{\psi'(x)}} = 2 \frac{\left(\frac{1 + a}{2}\right)^\alpha - a^\alpha}{1 - a^\alpha}.$$

Now, consider λ_U and note that

$$\begin{aligned}\lambda_U &= \lim_{u \uparrow 1} \frac{1 - 2u + C(u, u)}{1 - u} = 2 - \lim_{u \uparrow 1} \frac{1 - C(u, u)}{1 - u} \\ &= 2 - \lim_{u \uparrow 1} \frac{1 - \mathbb{P}(X_1 > \bar{F}^-(u), X_2 > \bar{F}^-(u))}{1 - \mathbb{P}(X_1 > \bar{F}^-(u))} \\ &= 2 - \lim_{x \downarrow 0} \frac{1 - \mathbb{P}(X_1 > x, X_2 > x)}{1 - \mathbb{P}(X_1 > x)}.\end{aligned}\quad (\text{A5})$$

Using (A2) and (A3), and proceeding similarly as above (with regular variation at 0), one obtains λ_U as stated.

(2) Under the given assumptions, using Corollary 1, we obtain that

$$\begin{aligned}\mathbb{P}(X_1 > x, X_2 > x) &= \begin{cases} \frac{2a}{1-a} \psi(x/a) + 1, & \text{if } x \leq 0, \\ \frac{1+a}{1-a} \psi(2x/(1+a)), & \text{if } x > 0, \end{cases} \\ \bar{F}(x) = \mathbb{P}(X_1 > x) = \mathbb{P}(X_2 > x) &= \begin{cases} \frac{a}{1-a} \psi(x/a) + 1, & \text{if } x \leq 0, \\ \frac{1}{1-a} \psi(x), & \text{if } x > 0. \end{cases}\end{aligned}$$

This yields

$$\lambda_L = \lim_{u \downarrow 0} \frac{C(u, u)}{u} = \lim_{x \uparrow \bar{F}^-(0)} \frac{(1+a)\psi(2x/(1+a))}{\psi(x)}.$$

If $\bar{F}^-(0) < \infty$, then $\lambda_L = 0$. If $\bar{F}^-(0) = \infty$, then l'Hôpital's Rule and regular variation imply that

$$\lambda_L = 2 \lim_{x \uparrow \infty} \frac{\psi'(\frac{2}{1+a}x)}{\psi'(x)} = 2 \left(\frac{2}{1+a}\right)^{-\alpha}.$$

Now, consider λ_U and note that the margins live on \mathbb{R} . Similar to above, we obtain that

$$\lambda_U = 2 - \lim_{x \downarrow -\infty} \frac{1 - \mathbb{P}(X_1 > x, X_2 > x)}{1 - \mathbb{P}(X_1 > x)}$$

which is easily seen to be 0.

□

References

- Cambanis, Stamatis, Steel Huang, and Gordon Simons. 1981. On the theory of elliptically contoured distributions. *Journal of Multivariate Analysis* 11: 368–85. [\[CrossRef\]](#)
- Embrechts, Paul, and Marius Hofert. 2011. Comments on: Inference in multivariate Archimedean copula models. *TEST* 20: 263–70. [\[CrossRef\]](#)
- Genest, Christian, Johanna Nešlehová, and Louis-Paul Rivest. 2018. The class of multivariate max-id copulas with ℓ_1 -norm symmetric exponent measure. *Bernoulli* 24: 3751–90. [\[CrossRef\]](#)
- Krupskii, Pavel, and Harry Joe. 2013. Factor copula models for multivariate data. *Journal of Multivariate Analysis* 120: 85–101. [\[CrossRef\]](#)
- Krupskii, Pavel, and Harry Joe. 2015. Structured factor copula models: Theory, inference and computation. *Journal of Multivariate Analysis* 138: 53–73. [\[CrossRef\]](#)
- McNeil, Alexander J., and Johanna Nešlehová. 2009. Multivariate Archimedean copulas, d -monotone functions and l_1 -norm symmetric distributions. *The Annals of Statistics* 37: 3059–97. [\[CrossRef\]](#)
- Nelsen, Roger B. 2006. *An Introduction to Copulas*. Berlin/Heidelberg: Springer.
- Quessy, Jean-François, and Martin Durocher. 2019. The class of copulas arising from squared distributions: Properties and inference. *Econometrics and Statistics* 12: 148–66. [\[CrossRef\]](#)

-
- Quessy, Jean-François, Louis-Paul Rivest, and Marie-Hélène Toupin. 2016. On the family of multivariate chi-square copulas. *Journal of Multivariate Analysis* 152: 40–60. [[CrossRef](#)]
- Rosco, J. F., and Harry Joe. 2013. Measures of tail asymmetry for bivariate copulas. *Statistical Papers* 54: 709–26. [[CrossRef](#)]