

+ Model

ARTICLE IN PRESS



ELSEVIER

Molecular and Cellular Endocrinology xxx (2006) xxx–xxx



Molecular and Cellular Endocrinology

www.elsevier.com/locate/mce

Regulation of junction dynamics in the testis—Transcriptional and post-translational regulations of cell junction proteins

Wing-Yee Lui, Will M. Lee*

Department of Zoology, The University of Hong Kong, Pokfulam Road, Hong Kong, PR China

Abstract

Cell junctions are the sites at which cells attach to the neighboring cells. They do not only maintain tissue integrity, their turnover also plays a crucial role in cell development and morphogenesis. In the testis, tight junctions and adherens junctions are dynamically remodeled to allow the movement of post-meiotic germ cells across the seminiferous epithelium and the timely release of spermatids into the tubular lumen. There is growing evidence that this dynamic remodeling of cell junctions is mediated by several mechanisms at the transcriptional and post-translational levels. This review summarizes what is known about the transcriptional regulation, ubiquitination and endocytosis that are involved in modulating junction dynamics in epithelial cells. It also highlights the recent findings on the regulation of junction dynamics in the testis and the specific areas that require further research for a thorough understanding of the role of junction remodeling in spermatogenesis. Understanding the junction dynamics in the seminiferous epithelium may unfold new targets for non-hormonal male contraceptive development.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Sertoli cells; Cell junctions; Transcription; Post-translation; Ubiquitination

1. Introduction

In rodent testes, a type A1 spermatogonium (diploid, 2n) gives rise to 256 mature spermatids (haploid, n) in the seminiferous epithelium during spermatogenesis (for reviews, see Cheng and Mruk, 2002; de Kretser and Kerr, 1988). Differentiating germ cells must traverse from the basal to the adluminal compartment of the seminiferous epithelium to complete spermatogenesis. Tight junctions (TJs) between Sertoli cells at the basal compartment form the blood–testis barrier (BTB), which segregates most part of spermatogenesis from the systemic circulation (for reviews, see Dym and Cavicchia, 1977; Dym and Fawcett, 1970; Russell and Peterson, 1985). At the late stage VIII through early stage IX, preleptotene and leptotene spermatocytes must traverse the BTB, where junctions are rapidly broken and subsequently reassemble to allow the spermatocytes to pass through (for reviews, see Cheng and Mruk, 2002; Lui et al., 2003d; Russell, 1977).

The translocation of germ cells across the seminiferous epithelium involves not only the dynamic modulation of TJ at the BTB, but also the extensive restructuring of actin-based adherens junctions (AJs) between Sertoli cells as well as between Sertoli

and germ cells thereafter, so that differentiating germ cells could move towards the adluminal compartment for further development (for reviews, see Cheng and Mruk, 2002; Lui et al., 2003d; Mruk and Cheng, 2004).

Previous studies in our laboratory have shown that the restructuring of cell junctions in the testis involves an array of biomolecules such as cytokines, proteases, protease inhibitors and extracellular matrix proteins, and signaling molecules including cAMP and cGMP (Lee and Cheng, 2004; Lui et al., 2001, 2003b, 2003f; Siu and Cheng, 2004; Siu et al., 2003a, 2003b; Wong et al., 2004).

Cytokines such as transforming growth factor- β 3 (TGF- β 3) and tumor necrosis factor- α (TNF- α) have been shown to be involved in regulating junction proteins such as claudin-11, occludin and ZO-1 at their transcriptional levels in Sertoli cells (Lui et al., 2001). Cytokine-mediated transcriptional regulation of those proteins results in altering the permeability of the TJ barrier both in vitro and in vivo (Lui et al., 2001, 2003b, 2003f; Siu and Cheng, 2004; Wong et al., 2004). These results clearly demonstrate that the transcriptional regulation of the junction proteins is one of the major regulatory mechanisms to modulate the junction dynamics in the testis. Apart from transcriptional regulation of junction proteins, our recent studies have demonstrated that junction restructuring is also mediated via post-translational modification of the junction protein. Ubiquitination regulates the bioavailability of targeted junction proteins

* Corresponding author. Tel.: +852 22990842; fax: +852 25599114.
E-mail address: hrszlw@hku.hk (W.M. Lee).

at the site of cell–cell contact, resulting in the opening and closing of cell junctions (Lui and Lee, 2005).

The structure and function of cell junctions in the testis have been reviewed extensively (for reviews, see Cheng and Mruk, 2002; Lui et al., 2003d; Mruk and Cheng, 2004). Here we attempt to highlight specific research areas that deserve attention in future studies. In this review, we shall focus on the (i) transcriptional regulation and (ii) post-translational modification of cell junction proteins at the site of cell–cell contact, which in turn regulate cell junction dynamics.

2. Structure and molecular composition of cell junctions in the testis

Morphologically, TJs form a continuous circumferential seal above the basal lamina of the seminiferous tubules in the testis. The molecular architecture of TJs has been unraveled rapidly in recent years (for reviews, see Fanning et al., 1999; Martin-Padura et al., 1998; Mitic et al., 2000). Three classes of integral membrane proteins have been positively identified in the testis including occludin, claudins and junctional adhesion molecules (JAM) (Martin-Padura et al., 1998; Moroi et al., 1998; Tsukita et al., 2001). Several peripheral membrane proteins, which are linked to the integral membrane proteins have also been identified at the site of TJs in the testis. They include zonula occludens-1 (ZO-1), ZO-2, cingulin and several others (for reviews, see Byers et al., 1991, 1992; Cheng and Mruk, 2002; Jesaitis and Goodenough, 1994; Lui et al., 2003d; Mruk and Cheng, 2004). Some of the peripheral proteins function to connect the integral membrane protein with the actin filament inside the cell.

Adherens junctions link cytoskeletal elements from one cell (e.g. Sertoli cell) to the same type or another type of cell (e.g. germ cell) or to the extracellular matrix, creating a network that maintains the architecture of the testis. More importantly, the dynamic nature of AJs in the testis permits the translocation of developing germ cells across the seminiferous epithelium during spermatogenesis. There are three classes of interlocking protein complexes identified at the AJs. They are the cadherin–catenin complex, nectin–afadin complex and integrin–laminin complex (Bouchard et al., 2000; Chapin et al., 2001; Chung et al., 1999; Lee et al., 2003; Ozaki-Kuroda et al., 2002; Siu et al., 2003b; Wine and Chapin, 1999). The regulatory mechanisms of the formation and disruption of TJs and AJs have been examined extensively (for reviews, see Cheng and Mruk, 2002; Lui et al., 2003d, 2003e; Mruk and Cheng, 2004), and the promoter analyses of occludin, claudins and cadherin have also begun (Batlle et al., 2000; Bolos et al., 2003; Chen et al., 2000; Comijn et al., 2001; Li and Mrsny, 2000; Mankertz et al., 2000).

3. Regulation of TJ dynamics by transcriptional regulation of tight junction proteins

Previous studies from our laboratory and others have revealed that cytokines down-regulate the transcription of the junction proteins in primary Sertoli cells and the human intestinal cell line, HT-29/B6 (Hellani et al., 2000; Lui et al., 2001; Mankertz et al., 2000; Siu et al., 2003a). This cytokine-mediated transcrip-

tional repression resulted in the reduced expression of junction proteins, concomitant with an increase in barrier permeability.

Although major advances have been made in the past 20 years in identifying the components of TJs as well as the signaling pathways involved in the regulation of TJ dynamics (for reviews, see Cheng and Mruk, 2002; Lui et al., 2003d, 2003e; Mruk and Cheng, 2004), the transcriptional regulation of TJ proteins involved in modulating TJ junction dynamics is limited to the transmembrane proteins such as occludin and claudins (Ikenouchi et al., 2003; Luk et al., 2004; Mankertz et al., 2000; Sakaguchi et al., 2002; Wachtel et al., 2001). It is obvious that studies of transcriptional regulation of other TJ protein members, including the peripheral membrane proteins, should be expanded to further our understanding of the precise regulatory mechanism(s) by which TJ dynamics are regulated at the transcriptional level.

3.1. Occludin

Occludin, a 64 kDa protein, was the first TJ integral membrane protein identified in many epithelia including the rat seminiferous epithelium (Furuse et al., 1993). Numerous studies have implicated that cytokines such as TNF α and interferon γ (IFN γ) exert negative regulatory effects on the expression of occludin in epithelial cells along with the disruption of TJ barrier monitored by transepithelial resistance (TER) measurement (Hellani et al., 2000; Lui et al., 2001; Mankertz et al., 2000; Siu et al., 2003a; Wachtel et al., 2001). Studies from Mankertz et al. (2000) have identified the promoter sequence essential for the regulation of occludin expression and TJ formation in HT-29/B6 cells. The 208 bp DNA fragment upstream from the putative transcription start site of occludin gene was shown to be necessary and sufficient in mediating the basal promoter activity. It was also shown that TNF α impaired TJ barrier function by lowering the expression of occludin in HT-29/B6 cells through the suppression of the promoter activity (Mankertz et al., 2000). Although a number of potential *cis*-acting motifs pertinent to TNF α -mediated gene transcription, such as NF-IL6 and NF- κ B (GGGAGGAGGC, at position 1753), were identified within the human occludin promoter sequences, the detailed intracellular pathway that mediates TNF α -dependent occludin gene repression remains to be elucidated.

Other studies by Wachtel et al. (2001) have shown a similar negative effect of TNF α on occludin gene transcription in astrocytes, but not in brain endothelial cells and Madin–Darby canine kidney cells (MDCK). It was found that TNF α suppressed occludin mRNA level, but not ZO-1 expression in astrocytes and the removal of TNF α from astrocytes could restore basal expression of occludin. The effect of TNF α on occludin expression in astrocytes is mediated through TNF α type-1 receptor and NF- κ B. It was suggested that NF- κ B might either function as a negative regulator through a direct interaction with the *cis*-acting motif located on occludin promoter or exert an indirect effect by activating a repressor that acted on the occludin promoter (Wachtel et al., 2001).

TNF α could decrease occludin expression via the transcriptional repression in HT-29/B6 cells and astrocytes, but it showed

no effect on brain endothelial cells and MDCK cells. These studies have demonstrated that TNF α represses the occludin promoter in a cell-specific manner. Recent findings in our laboratory have demonstrated that TNF α and TGF- β 3 play a crucial role in regulating TJ dynamics in the testis. They have been shown to exert a negative effect on occludin transcription, which in turn reduces the occludin protein level thus leading to an increase in the testicular barrier permeability (Lui et al., 2001, 2003c; Siu et al., 2003a). The effect of TGF- β 3 on occludin expression has been confirmed recently in in vivo studies (Lui et al., 2003f); however, the molecular mechanism on how TNF α and TGF- β 3 down-regulate the occludin promoter activity in Sertoli cells is not clearly known. Work remains to be done to elucidate the transcriptional repression machinery and the significance of the transcriptional repression of occludin in modulating TJ dynamics in the testis.

Recent studies have demonstrated that the Snail superfamily of repressors encoding transcription factors of the zinc-finger type are involved in transcriptional regulation of occludin (Ikenouchi et al., 2003). The two members of this superfamily, Snail and Slug, were found to repress the occludin transcription in many epithelial cells including the human breast cancer cell, MCF7 and the mammary epithelial cell, Eph4 (Ikenouchi et al., 2003; Ohkubo and Ozawa, 2004). Most of the studies of Snail and Slug have been focused on their role in the epithelium-mesenchyme transition (for reviews, see Nieto, 2002; Thiery, 2002). The functional importance of Snail superfamily in spermatogenesis has not been investigated. In this regard, it is noteworthy to examine whether Snail or Slug could elicit a stage-specific expression of junction proteins in the seminiferous epithelial cycle. Would Snail and Slug be involved in the complete shutdown of occludin expression at the stages VIII-IX of the epithelial cycle, when the BTB is disassembled to allow the translocation of preleptotene and leptotene spermatocytes towards the adluminal compartment of the seminiferous epithelium? The molecular function of Snail superfamily in the testis has yet to be defined.

3.2. Claudins

Similar to occludin, claudins are also integral membrane proteins (~23 kDa) found at the sites of TJs in both epithelia and endothelia (for reviews, see Heiskala et al., 2001; Morita et al., 1999a; Tsukita and Furuse, 1999). They are from large gene family with more than 20 members (for reviews, see Morita et al., 1999a; Tepass, 2003). Different mammalian tissues have their own combination of claudin members (Morita et al., 1999a). For instance, several claudin members have been found in the testis and they are claudin-1, -3, -4, -5, -7, -8, and -11 (Tsukita et al., 2001). Among them, claudin-11 is highly expressed in the testis and is an important TJ building block that constitutes TJ strands between Sertoli cells in the testis (Gow et al., 1999; Morita et al., 1999b).

Studies from Ikenouchi et al. and Ohkubo et al. have unraveled the importance of Snail in the transcriptional regulation of claudins (Ikenouchi et al., 2003; Ohkubo and Ozawa, 2004). The promoters of mouse claudin-3, -4 and -7 were character-

ized and each promoter has at least six E-boxes (Ikenouchi et al., 2003). The E-box motif, with sequence [CA(G/C)(G/C)TG], is identical to the binding sites for Snail (Ikenouchi et al., 2003). When Snail expression vector and claudin reporter constructs were co-transfected into Eph4 cells, the promoters of claudin-3, -4 and -7 were repressed remarkably. While co-transfection of mutant Snail lacking the N-terminal SNAG domain, which is essential for the repressor activity, the repression effect of Snail on claudin-7 promoter was impaired (Ikenouchi et al., 2003). These studies clearly illustrated that the transcription of claudins was directly regulated by Snail through modulating their promoter activities. However, studies from Ohkubo and Ozawa (2004) have shown that Snail is involved in direct transcriptional repression of occludin, but not claudin-1. It was found that overexpression of Snail in MDCK cells could only decrease the protein synthesis of claudin, suggesting that Snail down-regulated claudin-1 through the control of post-transcriptional events (Ohkubo and Ozawa, 2004). Such discrepancy in the role of Snail on transcriptional regulation of claudin genes as revealed by the two studies might be related to the use of different cell lines as well as the members of claudin gene chosen for study.

In view of the physiological significance of Snail superfamily in transcriptional regulation of TJ components, it is important that Snail and Slug should be studied more vigorously in the testis for the purpose of elucidating their role in the migration of preleptotene/leptotene spermatocytes across the BTB during the late stage VIII and early stage IX of the epithelial cycle.

Apart from Snail, several transcription factors such as the β -catenin/Tcf complex, Cdx homeodomain proteins/hepatocyte nuclear factor-1 α and Sp1 were reported to bind directly to claudin-1, -2 and -19 promoters, respectively (Luk et al., 2004; Miwa et al., 2000; Sakaguchi et al., 2002). For instance, β -catenin/Tcf complex binds the two putative Tcf4 binding elements in the 5' flanking region of claudin-1 to activate transcription (Miwa et al., 2000). HNF-1 α enhances Cdx2-mediated activation of claudin-2 promoter in Caco-2 cells and HNF-1 α is an organ-specific regulator of claudin-2 expression in the liver (Sakaguchi et al., 2002). These findings indicate that expressions of different claudin members in different tissues are under unique and sophisticated regulatory control.

Previous studies from our laboratory have shown that TGF- β 3 can down-regulate the expression of claudin-11 in cultured Sertoli cells at the time of TJ assembly, which in turn perturbs the TJ permeability barrier (Lui et al., 2001). These results thus suggest claudin-11 plays a vital role in the formation and maintenance of TJ barrier in the testis. Recent studies have also revealed that follicle stimulating hormone (FSH) and TNF α exert a negative effect on claudin-11 transcription in mouse Sertoli cells (Hellani et al., 2000). It was also found that the FSH-driven transcriptional repression of claudin-11 gene is mediated through the cAMP/protein kinase A pathway (Hellani et al., 2000). It is believed that a unique transcriptional mechanism might exist to confine the tissue-specific expression of claudin-11 in the testis and brain. Such a postulate is supported by the observation that organ-specific transcription factor such as HNF-1 α is involved

287 in regulating the expression of claudin-2 in the liver, but not in
288 other tissues (Sakaguchi et al., 2002).

289 Several *cis*-acting motifs including NY-F and GATA motifs
290 have been identified to be involved in the activation of
291 mouse claudin-11 transcription. Gel-shift analysis and co-
292 immunoprecipitation studies have shown that NY-F, GATA-1
293 and CREB form a transcriptional complex and bind to the same
294 GATA/NY-F overlapping motif on mouse claudin-11 promoter,
295 and overexpression of these transcription factors significantly
296 increased the claudin-11 promoter activity (Lui et al., unpub-
297 lished observation). These findings clearly suggest that the tran-
298 scriptional machinery of claudin-11 gene is different from the
299 other claudin members. In the proximal claudin-11 promoter,
300 some testis-specific *cis*-acting motifs such as SRY were iden-
301 tified by sequence analysis. It is noteworthy to determine how
302 SRY plays a role in regulating the expression of claudin-11 gene
303 in the adult testis as it has been suggested that during the testic-
304 ular development in the fetus, the expression of claudin-11 gene
305 is probably under the control of SRY (Hellani et al., 2000).

306 4. Regulation of AJ dynamics by transcriptional 307 regulation of adherens junction proteins

308 Ectoplasmic specializations (ES) are specialized actin-
309 based cell–cell AJs unique to the testis. Cadherin-catenin,
310 nectin/afadin and integrin/laminin complexes are the interlock-
311 ing protein complexes that can be found at the ES in the testis
312 (for reviews, see Cheng and Mruk, 2002; Lui et al., 2003e; Mruk
313 and Cheng, 2004). They can be found between Sertoli cells at the
314 basal region of the seminiferous epithelium (basal ES) as well as
315 at the apical region of the seminiferous epithelium where devel-
316 oping and mature spermatids attach onto Sertoli cells (apical
317 ES). The turnover of ES permits the movement of spermato-
318 cytes across the epithelium and allows the release of mature
319 spermatids from the seminiferous epithelium (for reviews, see
320 Cheng and Mruk, 2002; Mruk and Cheng, 2004). Previous stud-
321 ies in our laboratory have identified biomolecules such as Rho
322 GTPases and signaling pathways such as integrin-linked kinase
323 that are involved in AJ disassembly and reassembly in the testis
324 (Lui et al., 2003a; Siu et al., 2003b).

325 Studies from this and other laboratories have demonstrated
326 that the transcriptional regulation of AJ components also plays
327 a critical role in modulating the expression of AJ proteins on
328 epithelial cells, which results in the destruction of cell junctions
329 (Batlle et al., 2000; Cano et al., 2000). Understanding the tran-
330 scriptional mechanism by which transcription factors act on the
331 promoters of AJ components in other epithelial cells will pro-
332 vide a useful guideline on similar studies in the testis and the
333 modulation of AJ dynamics in the testis.

334 4.1. Cadherin

335 Previous studies have shown that the loss of cadherin expres-
336 sion is responsible for the breakdown of intercellular adhesion,
337 suggesting that the regulation of cadherin gene transcription is
338 one of the predominant mechanisms to control the AJ dynamics
339 in epithelial cells (Perl et al., 1998).

340 The transcriptional regulation of cadherin was extensively
341 studied in the field of cancer biology since the down-regulation
342 of E-cadherin expression is highly pertinent to the development
343 of tumors and their progression (Birchmeier and Behrens, 1994;
344 Takeichi, 1993). Recent studies have identified several transcrip-
345 tional factors including bHLH factor E12/E47, the two-handed
346 zinc factors ZEB-1 (δ EF1) and ZEB-2 (SIP-1) that are involved
347 in transcriptional repression of the cadherin gene (Comijn et al.,
348 2001; Grootclaes and Frisch, 2000; Perez-Moreno et al., 2001).
349 Interestingly, all these transcriptional repressors act through the
350 interaction with specific E-boxes on the proximal promoter of
351 cadherin, resulting in down-regulation of cadherin expression
352 (Comijn et al., 2001; Grootclaes and Frisch, 2000; Perez-
353 Moreno et al., 2001).

354 There is emerging evidence showing that the Snail super-
355 family of transcriptional factors are involved in the regulation of
356 cadherin gene transcription (for reviews, see Nieto, 2002; Thiery,
357 2002). Studies from several laboratories have demonstrated that
358 Snail and Slug zinc-finger proteins repressed the endogenous
359 E-cadherin expression in a panel of epithelial tumor cell lines of
360 different origins, ranging from bladder carcinoma to breast carci-
361 noma (Batlle et al., 2000; Hajra et al., 2002; Kurrey and Bapat,
362 2005). The Snail- and Slug-mediated repression pathways act
363 through the putative E-box motifs on the E-cadherin proximal
364 promoter (Batlle et al., 2000; Bolos et al., 2003; Ikenouchi et al.,
365 2003). This specific E-cadherin repression mechanism has been
366 unraveled by Peinado et al. It was found that Snail mediated
367 E-cadherin repression by the recruitment of the Sin3A/histone
368 deacetylase 1 (HDAC1)/HDAC2 complex at the E-cadherin pro-
369 moter (Peinado et al., 2004). At there, HDAC1/HDAC2 deacety-
370 lated the histone H3 and H4 proteins. This suggests that Snail
371 mediates chromatin remodeling and histone modifications to
372 repress the cadherin expression (Peinado et al., 2004).

373 4.2. Nectin

374 Nectin is a newly identified AJ integral membrane protein.
375 Much of the works performed previously focused on the identifi-
376 cation of its interacting partners and localization. Until recently,
377 the gene knockout studies have illustrated that nectin-2 is a major
378 component of the ES in the testis and plays a crucial role in sper-
379 matogenesis (Bouchard et al., 2000). Loss of nectin-2 in male
380 mice results in infertility. Proper formation and destruction of
381 nectin-2-based AJs between Sertoli cells and germ cells allow
382 the movement of developing germ cells (Bouchard et al., 2000;
383 Ozaki-Kuroda et al., 2002).

384 In our laboratory, we have recently isolated the mouse nectin-
385 2 promoter for detailed characterization. It was found that Sp1
386 and cAMP response element (CRE) motifs at the proximal pro-
387 moter played a crucial role in regulating gene transcription.
388 Interestingly, gel-shift assays, overexpression analysis and chro-
389 matin immunoprecipitation assays have unequivocally shown
390 that not only CREB protein interacts with the CRE *cis*-acting
391 motif, c-Jun, but not c-fos, also acts through the CRE motif
392 to up-regulate the nectin-2 transcription in Sertoli cells (Lui
393 et al, unpublished observation). This transcriptional regulation
394 is functionally significant to the testicular physiology as cyclic




Fig. 1. A schematic diagram illustrating the location of the cell junctions between Sertoli cells and different types of germ cells in the seminiferous epithelium. The blood–testis barrier, which is constituted by tight junctions, divides the epithelium into the basal and adluminal compartment. Anchoring junctions that are present in the seminiferous epithelium including (i) cell–cell actin-based adherens junctions such as ectoplasmic specialization and the tubulobulbar complex, (ii) cell–cell intermediate filament-based desmosome-like junctions. The dynamics rearrangement of cell junctions is essential to allow the movement of germ cells from the basal compartment to the adluminal compartment to complete spermatogenesis.

395 expression of CREB in the seminiferous epithelium concomi- 418
 396 tant with the expression of nectin-2 gene was observed when 419
 397 isolated staged seminiferous tubules were analysed (Don and 420
 398 Stelzer, 2002; Waeber et al., 1991). 421

399 5. Regulation of junction dynamics via 422 400 post-translational modification of junction protein at the 423 401 site of cell–cell contact 424

402 Cell junctions between Sertoli–Sertoli and Sertoli–germ cells 425
 403 are dynamically modulated and such changes can result in the 426
 404 translocation of differentiating germ cells from the basal to the 427
 405 adluminal compartment of the seminiferous epithelium for sper- 428
 406 miation. Undoubtedly, transcriptional regulation of the junction 429
 407 components is an important mechanism to control the expres- 430
 408 sion of the junction proteins (Ivanov et al., 2005; Nieto, 2002; 431
 409 Thiery, 2002). Still, the fate of the existing junction proteins 432
 410 at the site of cell–cell contact is a major determinant factor on 433
 411 junction dynamics. For instance, remodeling of junction pro- 434
 412 teins during epithelial morphogenesis occurs in 1 h (Schock 435
 413 and Perrimon, 2002), whereas the half-lives of the junction 436
 414 proteins are much longer, up to 12 h for occludin (Wong and 437
 415 Gumbiner, 1997). The disparity between the stability of junc- 438
 416 tion proteins and rapid junction remodeling apparently suggests 439
 417 that the transcriptional regulation of the junction protein is not 440

the sole mechanism to achieve rapid remodeling of the cell 418
 junctions. 419

Ubiquitination and endocytosis are recognized as the two 420
 essential mechanisms of targeted protein degradation whereby 421
 the targeted proteins are removed by the proteasome and the 422
 lysosomal system, respectively (for reviews, see Le Roy and 423
 Wrana, 2005; Takei et al., 2005; Wilkinson, 2000). There is 424
 growing evidence that ubiquitination and endocytosis of junc- 425
 tion proteins at the site of cell–cell contact are effective mech- 426
 anisms for the remodeling of cell junctions in dynamic situa- 427
 tions, where junctions must be rapidly broken and reassembled 428
 (Fujita et al., 2002; Kamei et al., 1999; Le et al., 1999; Lui and Lee, 2005; 429
 Paterson et al., 2003; Taya et al., 1998; Traweger et al., 2002). 430

Herein, we attempt to review (i) the mechanism of protein 431
 ubiquitination and deubiquitination, (ii) the covalent attach- 432
 ment of ubiquitin to the targeted junction proteins or removing ubi- 433
 quitin to rescue degradation, and (iii) endocytosis of cell junc- 434
 tion proteins (Fig. 1). 435

436 6. Ubiquitination 437

The ubiquitin conjugation system is composed of ubiquitin 437
 and three enzymes namely ubiquitin-activating enzyme (E1), 438
 ubiquitin-conjugating enzyme (E2) and ubiquitin-ligase (E3) 439
 (for reviews, see Hershko and Ciechanover, 1998; Hicke, 2001). 440




Fig. 2. A schematic drawing illustrating the dynamic regulation of junction proteins through ubiquitination and deubiquitination. At steady-state, junction proteins are organized at the site of cell–cell contacts. Under certain conditions such as cAMP-mediated junction disassembly, E3 ligase recognizes and binds to the specific junction protein substrate. Meanwhile, E1 activates ubiquitin and the activated ubiquitin is then conjugated with E2. With the aid of E2, E3 ligase functions to mediate the transfer of ubiquitin chains to the junction protein. The ubiquitinated protein may either be degraded into short peptides by proteasome with the release of ubiquitin chain, or undergo deubiquitination by deubiquitinating enzymes (DUBs). DUBs act on the ubiquitin chain and cleave the ubiquitin chain from that protein, resulting in rescuing the protein from degradation. The junction protein can be recycled back to the cell surface, which in turn facilitates the reassembly of cell junctions.

Ubiquitin, an evolutionary conserved protein with 76 amino acid residues, is first activated through the glycine residue in its C-terminus by E1. An E1-ubiquitin intermediate is formed with a high-energy thioester bond. The activated ubiquitin then passes a thiol group onto one of the E2. The E2 when bound to an E3 ligase transfers activated ubiquitin to the target protein (Fig. 2). Most organisms contain one or two E1s since E1 is able to carry out the activation of ubiquitin for all modifications, whereas E2 and E3 have their substrate specificities. Each E2 transfers ubiquitin specifically to a single or several E3 ligases, whilst each E3 ligase can associate with unique protein substrates through recognizing similar motifs (for review, see Pickart, 2001).

The protein substrate tagged with a polyubiquitin chain is then recognized and degraded into short peptides by the 26S proteasome complex. The ubiquitin is then released and recycled (for reviews, see Hicke, 2001; Pickart, 2001) (Fig. 2).

7. Deubiquitination

Ubiquitination can be reversed by members of a large family of enzymes known as isopeptidases or deubiquitinating enzymes (for reviews, see Kim et al., 2003; Wilkinson, 2000; Wing, 2003). Generally, the deubiquitinating enzymes can be divided into two main types (for review, see Wing, 2003). The first group functions to regenerate ubiquitin from proteolytic remnants produced by the proteasomes, thereby to speed up the proteasome-dependent proteolysis. The second group of the deubiquitinating enzymes is responsible for the reverse process of ubiquitin conjugation. In the process of deubiquitination, the

enzymes cleave the isopeptide bond of poly-ubiquitin chain proximal to the target proteins or hydrolyze the poly-ubiquitin chains into ubiquitin monomers, and thereby prevent the degradation of protein by the proteasome (Hochstrasser, 1995). By the action of the deubiquitinating enzymes, the ubiquitinated protein which is originally destined to the degradation process can then be rescued (Fig. 2).

8. Attachment of ubiquitin to the targeted proteins for degradation by proteasome

Using yeast two-hybrid screening, an E3 ubiquitin ligase Itch was identified to bind specifically to the NH₂-terminal portion of occludin (Traweger et al., 2002). This novel interaction between Itch and occludin is involved in the ubiquitination of occludin in MDCK cells, and the degradation of occludin is sensitive to proteasome inhibition (Traweger et al., 2002). Such interaction can also be found in the Sertoli cells by which TJs are dynamically rearranged to allow the movement of germ cells in the seminiferous epithelium (Lui and Lee, 2005). In addition, a novel interaction between Itch and UBC4 (an ubiquitin-conjugating enzyme) on occludin was detected by co-immunoprecipitation. Using the cAMP-mediated TJ disruption model, we have shown that an increase in protein levels of Itch and UBC4 along with a significant reduction in endogenous occludin was detected when TJs were disrupted by dibutyl-cAMP (db-cAMP). Addition of MG-132 (a proteasome inhibitor) could prevent db-cAMP-induced TJ disruption by altering the rate of occludin degradation (Lui and Lee, 2005). These studies support the notion that

the ubiquitination of TJ protein is one of the regulatory mechanisms in modulating the junction dynamics in the testis.

Recently, a specific E3 ubiquitin ligase targeted to E-cadherin was identified by yeast two-hybrid system (Fujita et al., 2002). This E3, known as Hakai, contains SH2, RING, zinc-finger and proline-rich domains. Hakai interacts with E-cadherin in a tyrosine phosphorylation-dependent manner and induces ubiquitination of the E-cadherin complex (Fujita et al., 2002). The activation by tyrosine kinases induces the tyrosine phosphorylation of the E-cadherin, which in turn attracts Hakai. Such interaction triggers the ubiquitination and degradation of the E-cadherin complex (Fujita et al., 2002; Pece and Gutkind, 2002).

Apart from the identification of E3 ubiquitin ligases specific to the integral protein of cell junctions, it has been reported that F-box/WD40-repeat protein functions as an E3 ubiquitin ligase to regulate the ubiquitination of β -catenin, a peripheral protein associated with cadherin at the AJ (Hart et al., 1999; Jiang and Struhl, 1998; Latres et al., 1999; Marikawa and Elinson, 1998). The ubiquitinated β -catenin is recognized by the proteasome and rapidly degraded. Several studies have illustrated the function of F-box/WD40-repeat protein on the degradation of β -catenin, however, all studies focused on its effect on growth and development via the Wntless-Wnt signaling pathway (Hart et al., 1999; Jiang and Struhl, 1998; Latres et al., 1999; Marikawa and Elinson, 1998). Therefore, the effect of E3-ligase-mediated β -catenin degradation on cell adhesion has not been fully characterized.

9. Deconjugating ubiquitin from the multiubiquitinated substrate to avoid protein degradation

Protein ubiquitination is a reversible process, and it has become increasingly obvious that ubiquitin deconjugation plays an important role in rescuing the ubiquitinated proteins from proteolysis (for reviews, see Kim et al., 2003; Wilkinson, 2000; Wing, 2003).

In the ubiquitin-proteasome pathway, the ubiquitinated substrates either undergo degradation by the proteasome or stabilization through the action of the deubiquitinating enzyme, suggesting that the bioavailability of proteins is tightly regulated by the action of the ubiquitinating and deubiquitinating enzymes (for review, see Wilkinson, 2000). As a result, the disassembly and reassembly of cell junctions could be manipulated by the precise regulation on the junction protein levels at the site of cell–cell contacts through the specific pathways of ubiquitination and deubiquitination (Fig. 2).

Studies by Taya et al. (1998) have shown that one of the deubiquitinating enzymes, Fam, is capable of regulating the degradation of peripheral component of nectin-based cell–cell adhesion. Immunofluorescence staining has shown that Fam is specifically localized at the sites of cell–cell contact in confluent MDCKII cells. Fam not only co-localizes with AF-6 (afadin), but also interacts with AF-6 in vitro and in vivo. More importantly, Fam can exert its deubiquitinating activity in vivo to release ubiquitin from the ubiquitinated AF-6, suggesting that the degradation of AF-6 at the site of cell–cell adhesion is regulated by Fam (Taya et al., 1998).

Apart from deubiquitinating the ubiquitinated AF-6, other studies have demonstrated that Fam also interacts with another AJ peripheral components, β -catenin. It was found that Fam could stabilize β -catenin by inhibiting its degradation, thereby prolonging the half-life of β -catenin (Taya et al., 1999). Although the detailed mechanism underlying the stabilization of β -catenin by Fam has not been fully elucidated, it is presumably through the deubiquitination of β -catenin.

It is obvious that the degradation of peripheral membrane proteins AF-6 and β -catenin are regulated through the FAM-mediated deubiquitination. However, virtually no deubiquitinating enzyme specific to the integral membrane protein of AJs and TJs has been identified. Work should be expanded to identify the candidates involved in deubiquitination of the junction proteins as well as to understand the precise regulatory mechanism(s) by which junction dynamics are regulated utilizing these deubiquitinating enzymes.

Although the exact mechanisms of ubiquitination and deubiquitination are not fully understood, the junction proteins at the site of cell–cell contacts are in an ubiquitination-deubiquitination equilibrium. Thus, the activity and the expression of the E3 ligases and the deubiquitinating enzymes at the site of cell–cell contacts play a crucial role in affecting this equilibrium, which in turn modulates the junction dynamics (Fujita et al., 2002).

10. Endocytosis of junction proteins

The level of junction proteins at the site of cell–cell contacts could be modulated by transcriptional regulation and/or protein degradation through the ubiquitin-proteasome pathway (Thiery, 2002). The endocytosis and recycling of junction proteins have recently emerged as an alternative mechanism allowing cells to undergo rapid changes in morphology in response to extracellular stimuli (Kamei et al., 1999; Le et al., 1999; Paterson et al., 2003) (Fig. 3).

Studies from Ivanov et al. have demonstrated that internalization of TJ proteins such as occludin, JAM-1, claudins-1 and -4 were observed in T84 epithelial cells by proteinase protection assay and immunocytochemistry (Ivanov et al., 2004). Addition of pharmacological inhibitors of clathrin-mediated endocytosis blocked the process, suggesting the endocytosis is clathrin-dependent. However, those TJ proteins are targeted neither for recycling nor for degradation in lysosomes. It was proposed that the endocytosed TJ proteins were shuffled into a unique storage compartment, hence providing a new mechanism to disrupt the TJ barrier (Ivanov et al., 2004).

Matsuda et al. have shown that the dynamic remodeling of TJs involves the elongation and shortening of individual TJs between two adjacent cells during intercellular motility (Matsuda et al., 2004). During the shortening of the individual TJs, vesicular structures containing claudin-3 were found in the cytoplasm. Interestingly, occludin, ZO-1 and JAM, which are the major building blocks of TJs, were not detected in the claudin-containing vesicles, suggesting that claudin-3, but not other TJ components were selectively segregated during TJ internalization (Matsuda et al., 2004). These results suggest that distinctive




Fig. 3. Disassembly and reassembly of cell junctions are mediated by endocytosis and recycling of cell junction proteins. Junction proteins are internalized via different endocytic structures including caveolin-coated vesicle (a), clathrin-coated vesicle (b) and actin-coated vacuolae (phagosome) (c). Internalized proteins are delivered into early endosomes. Subsequently they are delivered to either late endosome which targets for degradation in lysosome (d) or recycling endosome for channeling back to the cell surface (e).

605 internalization of claudins plays a crucial role in the remodeling of TJs and the selective regulation of claudin endocytosis is important in intercellular motility (Matsuda et al., 2004).

606
607
608 Endocytosis not only plays a role in TJ dynamics, its effect on the regulation of AJ dynamics is well-documented (Kamei et al., 1999; Le et al., 1999; Palacios et al., 2001, 2002). For instance, using surface biotinylation and recycling assays, Le et al. have shown that some of the E-cadherin at the cell surface are actively internalized and then recycled back to the plasma membrane through the classic clathrin-mediated endocytosis pathway (Le et al., 1999). When cells were cultured in Ca^{2+} -depleted medium, a significant increase in the endocytosis and recycling cell surface E-cadherin was observed. The reassembly of AJs by Ca^{2+} -repletion was inhibited by bafilomycin-mediated disruption of the endocytosed E-cadherin recycling (Le et al., 1999). These results support the notion that endocytosis and the recycling of the junction proteins are involved in the regulation of junction dynamics.

623 Subsequent studies have also identified the signaling cascades and biomolecules involved in the internalization of AJ proteins (Kamei et al., 1999; Palacios et al., 2001, 2002; Paterson et al., 2003). The diversity of the internalization mechanisms for junction protein has already been covered in an excellent recent review (Ivanov et al., 2005). Readers are strongly encouraged to seek additional information on these subject areas from this article.

631 Previous studies in our laboratory have demonstrated that RhoB GTPase are also involved in regulating Sertoli–germ AJ dynamics (Lui et al., 2003a). Several small GTPases such as Rho and Rab family members are involved directly and indirectly in the endocytosis and recycling of E-cadherin in several epithelial cells (Kamei et al., 1999; Palacios et al., 2001, 2002;

637 Paterson et al., 2003). For instance, the activation of ARF6 in MDCK cells promotes the clathrin-dependent internalization of E-cadherin, resulting in the disassembly of AJs without the remodeling of actin filament (Palacios et al., 2001). Recently, the molecular mechanism of ARF6 on AJ disassembly has been identified. It was found that ARF6-GTP interacted with and recruited Nm23-H1, a nucleoside diphosphate kinase, to facilitate dynamin-mediated endocytosis during AJ disassembly (Palacios et al., 2002). All these studies clearly suggest small GTPases play an important role in regulating the endocytosis of junction proteins, resulting in junction remodeling.

11. Concluding remarks

649 In this review, we have summarized some of the recent findings in the study of junction dynamics in epithelial cells, some of the potentially important regulators such as E3 ligase, and regulatory pathways of junction dynamics recently identified in the testis. It is obvious that many questions remain to be addressed. For instance, the precise transcriptional regulation in controlling the testis-specific expression of junction proteins and transcriptional repression of TJ proteins at the stages VIII and IX, by which preleptotene and leptotene spermatocytes traverse the BTB, remain unknown.

659 Studies of the effect of ubiquitination on junction dynamics in the seminiferous epithelium are very limited, apart from a recent study assessing the role of Itch (E3 ligase) on the Sertoli TJ barrier in vitro. This apparently is a priority area that needs to be further investigated in the near future. As such, the identification of specific E3 ligases targeted to different junction proteins and the elucidation of the precise regulatory mechanisms are needed to be addressed.

Another interesting topic that deserves further investigation is whether endocytosis plays a role in modulating the junction dynamics in the testis. Many studies have demonstrated that endocytosis of junction protein is a rapid and effective way in reorganizing cell junctions. This information from the studies of other epithelial cells provides a blueprint for us to examine the role of endocytosis on junction dynamics in the seminiferous epithelium.

A thorough understanding of germ cell migration during spermatogenesis, in particular, the cell junction dynamics in the seminiferous epithelium, would allow the identification of new targets for non-hormonal male contraceptive development. For instance, selective repression of the junction protein expression at the apical ES, such as nectin-2, might induce premature release of spermatids into the tubular lumen via the transcriptional regulation. Alternatively, if prolonged expression of junction proteins could be procured, germ cells might be trapped in the seminiferous epithelium for an extended period leading to apoptosis. This post-meiotic approach of male contraception can be achieved by the identification of the testis-specific transcription factors that are involved in regulating junction protein expression in the seminiferous epithelium.

The precise control of junction protein turnover in the seminiferous epithelium provides another line of potential for male contraceptive development. For example, alteration of ubiquitination or endocytosis of junction proteins will interfere the dynamic control of junction disassembly and reassembly in the seminiferous epithelium and thus may cause the loss of fertility in the male.

Acknowledgements

This work was supported by grants from the Hong Kong Research Grant Council (HKU 7413/04M and HKU 7536/05M) and The University of Hong Kong (Seed Fund for the Strategic Theme on Healthy Aging to WML and CRCG Small Project Funding to WYL).

References

Battle, E., Sancho, E., Franci, C., Dominguez, D., Monfar, M., Baulida, J., Garcia De Herreros, A., 2000. The transcription factor snail is a repressor of E-cadherin gene expression in epithelial tumour cells. *Nat. Cell Biol.* 2, 84–89.

Birchmeier, W., Behrens, J., 1994. Cadherin expression in carcinomas: role in the formation of cell junctions and the prevention of invasiveness. *Biochim. Biophys. Acta* 1198, 11–26.

Bolos, V., Peinado, H., Perez-Moreno, M.A., Fraga, M.F., Esteller, M., Cano, A., 2003. The transcription factor Slug represses E-cadherin expression and induces epithelial to mesenchymal transitions: a comparison with Snail and E47 repressors. *J. Cell Sci.* 116, 499–511.

Bouchard, M.J., Dong, Y., McDermott Jr., B.M., Lam, D.H., Brown, K.R., Shelanski, M., Bellve, A.R., Racaniello, V.R., 2000. Defects in nuclear and cytoskeletal morphology and mitochondrial localization in spermatozoa of mice lacking nectin-2, a component of cell–cell adherens junctions. *Mol. Cell Biol.* 20, 2865–2873.

Byers, S., Graham, R., Dai, H.N., Hoxter, B., 1991. Development of Sertoli cell junctional specializations and the distribution of the tight-junction-associated protein ZO-1 in the mouse testis. *Am. J. Anat.* 191, 35–47.

Byers, S.W., Citi, S., Anderson, J.M., Hoxter, B., 1992. Polarized functions and permeability properties of rat epididymal epithelial cells in vitro. *J. Reprod. Fertil.* 95, 385–396.

Cano, A., Perez-Moreno, M.A., Rodrigo, I., Locascio, A., Blanco, M.J., del Barrio, M.G., Portillo, F., Nieto, M.A., 2000. The transcription factor snail controls epithelial-mesenchymal transitions by repressing E-cadherin expression. *Nat. Cell Biol.* 2, 76–83.

Chapin, R.E., Wine, R.N., Harris, M.W., Borchers, C.H., Haseman, J.K., 2001. Structure and control of a cell–cell adhesion complex associated with spermiation in rat seminiferous epithelium. *J. Androl.* 22, 1030–1052.

Chen, Y., Lu, Q., Schneeberger, E.E., Goodenough, D.A., 2000. Restoration of tight junction structure and barrier function by down-regulation of the mitogen-activated protein kinase pathway in ras-transformed Madin–Darby canine kidney cells. *Mol. Biol. Cell* 11, 849–862.

Cheng, C.Y., Mruk, D.D., 2002. Cell junction dynamics in the testis: Sertoli–germ cell interactions and male contraceptive development. *Physiol. Rev.* 82, 825–874.

Chung, S.S., Lee, W.M., Cheng, C.Y., 1999. Study on the formation of specialized inter-Sertoli cell junctions in vitro. *J. Cell Physiol.* 181, 258–272.

Comijn, J., Berx, G., Vermassen, P., Verschuere, K., van Grunsven, L., Bruyneel, E., Mareel, M., Huylebroeck, D., van Roy, F., 2001. The two-handed E box binding zinc finger protein SIP1 downregulates E-cadherin and induces invasion. *Mol. Cell* 7, 1267–1278.

de Kretser, D.M., Kerr, J.B., 1988. The cytology of the testis. In: Knobil, E., Neill, J. (Eds.), *The Physiology of Reproduction*, vol. 1. Raven, New York, pp. 837–932.

Don, J., Stelzer, G., 2002. The expanding family of CREB/CREM transcription factors that are involved with spermatogenesis. *Mol. Cell Endocrinol.* 187, 115–124.

Dym, M., Cavicchia, J.C., 1977. Further observations on the blood–testis barrier in monkeys. *Biol. Reprod.* 17, 390–403.

Dym, M., Fawcett, D.W., 1970. The blood–testis barrier in the rat and the physiological compartmentation of the seminiferous epithelium. *Biol. Reprod.* 3, 308–326.

Fanning, A.S., Mitic, L.L., Anderson, J.M., 1999. Transmembrane proteins in the tight junction barrier. *J. Am. Soc. Nephrol.* 10, 1337–1345.

Fujita, Y., Krause, G., Scheffner, M., Zechner, D., Leddy, H.E., Behrens, J., Sommer, T., Birchmeier, W., 2002. Hakai, a c-Cbl-like protein, ubiquitinates and induces endocytosis of the E-cadherin complex. *Nat. Cell Biol.* 4, 222–231.

Furuse, M., Hirase, T., Itoh, M., Nagafuchi, A., Yonemura, S., Tsukita, S., 1993. Occludin: a novel integral membrane protein localizing at tight junctions. *J. Cell Biol.* 123, 1777–1788.

Gow, A., Southwood, C.M., Li, J.S., Pariali, M., Riordan, G.P., Brodie, S.E., Dianas, J., Bronstein, J.M., Kachar, B., Lazzarini, R.A., 1999. CNS myelin and sertoli cell tight junction strands are absent in *Osp/claudin-11* null mice. *Cell* 99, 649–659.

Grooteclaes, M.L., Frisch, S.M., 2000. Evidence for a function of CtBP in epithelial gene regulation and anoikis. *Oncogene* 19, 3823–3828.

Hajra, K.M., Chen, D.Y., Fearon, E.R., 2002. The SLUG zinc-finger protein represses E-cadherin in breast cancer. *Cancer Res.* 62, 1613–1618.

Hart, M., Concordet, J.P., Lassot, I., Albert, I., del los Santos, R., Durand, H., Perret, C., Rubinfeld, B., Margottin, F., Benarous, R., Polakis, P., 1999. The F-box protein β -TrCP associates with phosphorylated β -catenin and regulates its activity in the cell. *Curr. Biol.* 9, 207–210.

Heiskala, M., Peterson, P.A., Yang, Y., 2001. The roles of claudin superfamily proteins in paracellular transport. *Traffic* 2, 93–98.

Hellani, A., Ji, J., Mauduit, C., Deschildre, C., Tabone, E., Benahmed, M., 2000. Developmental and hormonal regulation of the expression of oligodendrocyte-specific protein/claudin 11 in mouse testis. *Endocrinology* 141, 3012–3019.

Hershko, A., Ciechanover, A., 1998. The ubiquitin system. *Annu. Rev. Biochem.* 67, 425–479.

Hicke, L., 2001. A new ticket for entry into budding vesicles-ubiquitin. *Cell* 106, 527–530.

Hochstrasser, M., 1995. Ubiquitin, proteasomes, and the regulation of intracellular protein degradation. *Curr. Opin. Cell Biol.* 7, 215–223.

- Ikenouchi, J., Matsuda, M., Furuse, M., Tsukita, S., 2003. Regulation of tight junctions during the epithelium-mesenchyme transition: direct repression of the gene expression of claudins/occludin by Snail. *J. Cell Sci.* 116, 1959–1967.
- Ivanov, A.I., Nusrat, A., Parkos, C.A., 2004. Endocytosis of epithelial apical junctional proteins by a clathrin-mediated pathway into a unique storage compartment. *Mol. Biol. Cell* 15, 176–188.
- Ivanov, A.I., Nusrat, A., Parkos, C.A., 2005. Endocytosis of the apical junctional complex: mechanisms and possible roles in regulation of epithelial barriers. *Bioessays* 27, 356–365.
- Jesaitis, L.A., Goodenough, D.A., 1994. Molecular characterization and tissue distribution of ZO-2, a tight junction protein homologous to ZO-1 and the Drosophila discs-large tumor suppressor protein. *J. Cell Biol.* 124, 949–961.
- Jiang, J., Struhl, G., 1998. Regulation of the Hedgehog and Wingless signalling pathways by the F-box/WD40-repeat protein Slimb. *Nature* 391, 493–496.
- Kamei, T., Matozaki, T., Sakisaka, T., Kodama, A., Yokoyama, S., Peng, Y.F., Nakano, K., Takaishi, K., Takai, Y., 1999. Coendocytosis of cadherin and c-Met coupled to disruption of cell–cell adhesion in MDCK cells—regulation by Rho, Rac and Rab small G proteins. *Oncogene* 18, 6776–6784.
- Kim, J.H., Park, K.C., Chung, S.S., Bang, O., Chung, C.H., 2003. Deubiquitinating enzymes as cellular regulators. *J. Biochem. (Tokyo)* 134, 9–18.
- Kurrey, N.K.K.A., Bapat, S.A., 2005. Snail and Slug are major determinants of ovarian cancer invasiveness at the transcription level. *Gynecol. Oncol.* 97, 155–165.
- Latres, E., Chiaur, D.S., Pagano, M., 1999. The human F box protein β -Trcp associates with the Cul1/Skp1 complex and regulates the stability of β -catenin. *Oncogene* 18, 849–854.
- Le Roy, C., Wrana, J.L., 2005. Clathrin- and non-clathrin-mediated endocytic regulation of cell signalling. *Nat. Rev. Mol. Cell Biol.* 6, 112–126.
- Le, T.L., Yap, A.S., Stow, J.L., 1999. Recycling of E-cadherin: a potential mechanism for regulating cadherin dynamics. *J. Cell Biol.* 146, 219–232.
- Lee, N.P., Cheng, C.Y., 2004. Nitric oxide/nitric oxide synthase, spermatogenesis, and tight junction dynamics. *Biol. Reprod.* 70, 267–276.
- Lee, N.P., Mruk, D., Lee, W.M., Cheng, C.Y., 2003. Is the cadherin/catenin complex a functional unit of cell–cell actin-based adherens junctions in the rat testis? *Biol. Reprod.* 68, 489–508.
- Li, D., Mrsny, R.J., 2000. Oncogenic Raf-1 disrupts epithelial tight junctions via downregulation of occludin. *J. Cell Biol.* 148, 791–800.
- Lui, W.Y., Lee, W.M., 2005. cAMP perturbs inter-Sertoli tight junction permeability barrier in vitro via its effect on proteasome-sensitive ubiquitination of occludin. *J. Cell Physiol.* 203, 564–572.
- Lui, W.Y., Lee, W.M., Cheng, C.Y., 2001. Transforming growth factor- β 3 perturbs the inter-Sertoli tight junction permeability barrier in vitro possibly mediated via its effects on occludin, zonula occludens-1, and claudin-11. *Endocrinology* 142, 1865–1877.
- Lui, W.Y., Lee, W.M., Cheng, C.Y., 2003a. Sertoli–germ cell adherens junction dynamics in the testis are regulated by RhoB GTPase via the ROCK/LIMK signaling pathway. *Biol. Reprod.* 68, 2189–2206.
- Lui, W.Y., Lee, W.M., Cheng, C.Y., 2003b. TGF- β s: their role in testicular function and Sertoli cell tight junction dynamics. *Int. J. Androl.* 26, 147–160.
- Lui, W.Y., Lee, W.M., Cheng, C.Y., 2003c. Transforming growth factor β 3 regulates the dynamics of Sertoli cell tight junctions via the p38 mitogen-activated protein kinase pathway. *Biol. Reprod.* 68, 1597–1612.
- Lui, W.Y., Mruk, D., Lee, W.M., Cheng, C.Y., 2003d. Sertoli cell tight junction dynamics: their regulation during spermatogenesis. *Biol. Reprod.* 68, 1087–1097.
- Lui, W.Y., Mruk, D.D., Lee, W.M., Cheng, C.Y., 2003e. Adherens junction dynamics in the testis and spermatogenesis. *J. Androl.* 24, 1–14.
- Lui, W.Y., Wong, C.H., Mruk, D.D., Cheng, C.Y., 2003f. TGF- β 3 regulates the blood–testis barrier dynamics via the p38 mitogen activated protein (MAP) kinase pathway: an in vivo study. *Endocrinology* 144, 1139–1142.
- Luk, J.M., Tong, M.K., Mok, B.W., Tam, P.C., Yeung, W.S., Lee, K.F., 2004. Sp1 site is crucial for the mouse claudin-19 gene expression in the kidney cells. *FEBS Lett.* 578, 251–256.
- Mankertz, J., Tavalali, S., Schmitz, H., Mankertz, A., Riecken, E.O., Fromm, M., Schulzke, J.D., 2000. Expression from the human occludin promoter is affected by tumor necrosis factor α and interferon γ . *J. Cell Sci.* 113, 2085–2090.
- Marikawa, Y., Elinson, R.P., 1998. β -TrCP is a negative regulator of Wnt/ β -catenin signaling pathway and dorsal axis formation in *Xenopus* embryos. *Mech. Dev.* 77, 75–80.
- Martin-Padura, I., Lostaglio, S., Schneemann, M., Williams, L., Romano, M., Fruscella, P., Panzeri, C., Stoppacciaro, A., Ruco, L., Villa, A., Simmons, D., Dejana, E., 1998. Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *J. Cell Biol.* 142, 117–127.
- Matsuda, M., Kubo, A., Furuse, M., Tsukita, S., 2004. A peculiar internalization of claudins, tight junction-specific adhesion molecules, during the intercellular movement of epithelial cells. *J. Cell Sci.* 117, 1247–1257.
- Mitic, L.L., Van Itallie, C.M., Anderson, J.M., 2000. Molecular physiology and pathophysiology of tight junctions. I. Tight junction structure and function: lessons from mutant animals and proteins. *Am. J. Physiol. Gastrointest. Liver Physiol.* 279, G250–G254.
- Miwa, N., Furuse, M., Tsukita, S., Niikawa, N., Nakamura, Y., Furukawa, Y., 2000. Involvement of claudin-1 in the beta-catenin/Tcf signaling pathway and its frequent upregulation in human colorectal cancers. *Oncol. Res.* 12, 469–476.
- Morita, K., Furuse, M., Fujimoto, K., Tsukita, S., 1999a. Claudin multigene family encoding four-transmembrane domain protein components of tight junction strands. *Proc. Natl. Acad. Sci. U.S.A.* 96, 511–516.
- Morita, K., Sasaki, H., Fujimoto, K., Furuse, M., Tsukita, S., 1999b. Claudin-11/OSP-based tight junctions of myelin sheaths in brain and Sertoli cells in testis. *J. Cell Biol.* 145, 579–588.
- Moroi, S., Saitou, M., Fujimoto, K., Sakakibara, A., Furuse, M., Yoshida, O., Tsukita, S., 1998. Occludin is concentrated at tight junctions of mouse/rat but not human/guinea pig Sertoli cells in testes. *Am. J. Physiol.* 274, C1708–C1717.
- Mruk, D.D., Cheng, C.Y., 2004. Sertoli–Sertoli and Sertoli–germ cell interactions and their significance in germ cell movement in the seminiferous epithelium during spermatogenesis. *Endocr. Rev.* 25, 747–806.
- Nieto, M.A., 2002. The snail superfamily of zinc-finger transcription factors. *Nat. Rev. Mol. Cell Biol.* 3, 155–166.
- Ohkubo, T., Ozawa, M., 2004. The transcription factor Snail downregulates the tight junction components independently of E-cadherin downregulation. *J. Cell Sci.* 117, 1675–1685.
- Ozaki-Kuroda, K., Nakanishi, H., Ohta, H., Tanaka, H., Kurihara, H., Mueller, S., Irie, K., Ikeda, W., Sakai, T., Wimmer, E., Nishimune, Y., Takai, Y., 2002. Nectin couples cell–cell adhesion and the actin scaffold at heterotypic testicular junctions. *Curr. Biol.* 12, 1145–1150.
- Palacios, F., Price, L., Schweitzer, J., Collard, J.G., D'Souza-Schorey, C., 2001. An essential role for ARF6-regulated membrane traffic in adherens junction turnover and epithelial cell migration. *EMBO J.* 20, 4973–4986.
- Palacios, F., Schweitzer, J.K., Boshans, R.L., D'Souza-Schorey, C., 2002. ARF6-GTP recruits Nm23-H1 to facilitate dynamin-mediated endocytosis during adherens junctions disassembly. *Nat. Cell Biol.* 4, 929–936.
- Paterson, A.D., Parton, R.G., Ferguson, C., Stow, J.L., Yap, A.S., 2003. Characterization of E-cadherin endocytosis in isolated MCF-7 and Chinese hamster ovary cells: the initial fate of unbound E-cadherin. *J. Biol. Chem.* 278, 21050–21057.
- Pece, S., Gutkind, J.S., 2002. E-cadherin and Hakai: signalling, remodeling or destruction? *Nat. Cell Biol.* 4, E72–E74.
- Peinado, H., Ballestar, E., Esteller, M., Cano, A., 2004. Snail mediates E-cadherin repression by the recruitment of the Sin3A/histone deacetylase 1 (HDAC1)/HDAC2 complex. *Mol. Cell Biol.* 24, 306–319.
- Perez-Moreno, M.A., Locascio, A., Rodrigo, I., Dhondt, G., Portillo, F., Nieto, M.A., Cano, A., 2001. A new role for E12/E47 in the repression of E-cadherin expression and epithelial-mesenchymal transitions. *J. Biol. Chem.* 276, 17431–17424.
- Perl, A.K., Wilgenbus, P., Dahl, U., Semb, H., Christofori, G., 1998. A causal role for E-cadherin in the transition from adenoma to carcinoma. *Nature* 392, 190–193.

- 927 Pickart, C.M., 2001. Mechanisms underlying ubiquitination. *Annu. Rev. Biochem.* 70, 503–533. 962
- 928 Russell, L., 1977. Movement of spermatocytes from the basal to the adluminal compartment of the rat testis. *Am. J. Anat.* 148, 313–328. 963
- 929 Russell, L.D., Peterson, R.N., 1985. Sertoli cell junctions: morphological and functional correlates. *Int. Rev. Cytol.* 94, 177–211. 964
- 930 Russell, L.D., Peterson, R.N., 1985. Sertoli cell junctions: morphological and functional correlates. *Int. Rev. Cytol.* 94, 177–211. 965
- 931 Sakaguchi, T., Gu, X., Golden, H.M., Suh, E., Rhoads, D.B., Reinecker, H.C., 2002. Cloning of the human claudin-2 5'-flanking region revealed a TATA-less promoter with conserved binding sites in mouse and human for caudal-related homeodomain proteins and hepatocyte nuclear factor-1 α . *J. Biol. Chem.* 277, 21361–21370. 966
- 932 Schock, F., Perrimon, N., 2002. Molecular mechanisms of epithelial morphogenesis. *Annu. Rev. Cell. Dev. Biol.* 18, 463–493. 967
- 933 Siu, M.K., Cheng, C.Y., 2004. Interactions of proteases, protease inhibitors, and the beta1 integrin/laminin γ 3 protein complex in the regulation of ectoplasmic specialization dynamics in the rat testis. *Biol. Reprod.* 70, 945–964. 968
- 934 Siu, M.K., Lee, W.M., Cheng, C.Y., 2003a. The interplay of collagen IV, tumor necrosis factor- α , gelatinase B (matrix metalloprotease-9), and tissue inhibitor of metalloproteases-1 in the basal lamina regulates Sertoli cell-tight junction dynamics in the rat testis. *Endocrinology* 144, 371–387. 969
- 935 Siu, M.K., Mruk, D.D., Lee, W.M., Cheng, C.Y., 2003b. Adhering junction dynamics in the testis are regulated by an interplay of β 1-integrin and focal adhesion complex-associated proteins. *Endocrinology* 144, 2141–2163. 970
- 936 Takei, K., Yoshida, Y., Yamada, H., 2005. Regulatory mechanisms of dynamin-dependent endocytosis. *J. Biochem. (Tokyo)* 137, 243–247. 971
- 937 Takeichi, M., 1993. Cadherins in cancer: implications for invasion and metastasis. *Curr. Opin. Cell Biol.* 5, 806–811. 972
- 938 Taya, S., Yamamoto, T., Kanai-Azuma, M., Wood, S.A., Kaibuchi, K., 1999. The deubiquitinating enzyme Fam interacts with and stabilizes β -catenin. *Genes Cells* 4, 757–767. 973
- 939 Taya, S., Yamamoto, T., Kano, K., Kawano, Y., Iwamatsu, A., Tsuchiya, T., Tanaka, K., Kanai-Azuma, M., Wood, S.A., Mattick, J.S., Wood, S.A., Mattick, J.S., Kaibuchi, K., 1998. The Ras target AF-6 is a substrate of the fam deubiquitinating enzyme. *J. Cell Biol.* 142, 1053–1062. 974
- 940 Tepass, U., 2003. Claudin complexities at the apical junctional complex. *Nat. Cell Biol.* 5, 595–597. 975
- 941 Thiery, J.P., 2002. Epithelial-mesenchymal transitions in tumour progression. *Nat. Rev. Cancer* 2, 442–454. 976
- 942 Traweger, A., Fang, D., Liu, Y.C., Stelzhammer, W., Krizbai, I.A., Fresser, F., Bauer, H.C., Bauer, H., 2002. The tight junction-specific protein occludin is a functional target of the E3 ubiquitin-protein ligase itch. *J. Biol. Chem.* 277, 10201–10208. 977
- 943 Tsukita, S., Furuse, M., 1999. Occludin and claudins in tight-junction strands: leading or supporting players? *Trends Cell Biol.* 9, 268–273. 978
- 944 Tsukita, S., Furuse, M., Itoh, M., 2001. Multifunctional strands in tight junctions. *Nat. Rev. Mol. Cell Biol.* 2, 285–293. 979
- 945 Wachtel, M., Bolliger, M.F., Ishihara, H., Frei, K., Bluethmann, H., Gloor, S.M., 2001. Down-regulation of occludin expression in astrocytes by tumour necrosis factor (TNF) is mediated via TNF type-1 receptor and nuclear factor-kappa B activation. *J. Neurochem.* 78, 155–162. 980
- 946 Waeber, G., Meyer, T.E., LeSieur, M., Hermann, H.L., Gerard, N., Habener, J.F., 1991. Developmental stage-specific expression of cyclic adenosine 3',5'-monophosphate response element-binding protein CREB during spermatogenesis involves alternative exon splicing. *Mol. Endocrinol.* 5, 1418–1430. 981
- 947 Wilkinson, K.D., 2000. Ubiquitination and deubiquitination: targeting of proteins for degradation by the proteasome. *Semin. Cell Dev. Biol.* 11, 141–148. 982
- 948 Wine, R.N., Chapin, R.E., 1999. Adhesion and signaling proteins spatiotemporally associated with spermatiation in the rat. *J. Androl.* 20, 198–213. 983
- 949 Wing, S.S., 2003. Deubiquitinating enzymes—the importance of driving in reverse along the ubiquitin-proteasome pathway. *Int. J. Biochem. Cell Biol.* 35, 590–605. 984
- 950 Wong, C.H., Mruk, D.D., Lui, W.Y., Cheng, C.Y., 2004. Regulation of blood-testis barrier dynamics: an in vivo study. *J. Cell Sci.* 117, 783–798. 985
- 951 Wong, V., Gumbiner, B.M., 1997. A synthetic peptide corresponding to the extracellular domain of occludin perturbs the tight junction permeability barrier. *J. Cell Biol.* 136, 399–409. 986
- 952 987 988 989 990 991 992 993 994 995 996 997