

## **3261 Effects of Children Toothpastes on Enamel Caries Using Two Models**

**[A. ITTHAGARUN](#)**<sup>1</sup>, P. THAVEESANGPANICH<sup>1</sup>, N.M. KING<sup>1</sup>, and J.S. WEFEL<sup>2</sup>,  
<sup>1</sup>University of Hong Kong, Hong Kong SAR, China, <sup>2</sup>The University of Iowa,  
Iowa City, USA

**Aims/objectives:** To compare, using two pH-cycling models, the de/remineralization effects of children's toothpastes on primary teeth. **Design:** In vitro single-section and pH-cycling models. **Methods:** Primary teeth were placed in demineralizing solution for 96 hours to produce artificial carious lesions 60-100  $\mu\text{m}$  deep. They were cut into 100  $\mu\text{m}$  thick sections and assigned to 6 groups. Sections in Groups A and D were exposed to a non-fluoridated toothpaste, those in Groups B and E to half-pea-sized (0.16 grams) and those in Groups C and F to pea-sized portions (0.32 grams) of a 500ppm F toothpaste. pH-cycling Model I (Groups A, B, C), without added fluoride, ran for 7 days, while Model II (Groups D, E, F), with 0.25ppm F, ran for 10 days. **Results:** Lesions were evaluated using polarized light microscopy and microradiography. The lesions increased in depth by 64% in Group A, 70% in Group B, and 20% in Group C. For Model II specimens, those in Group D increased by 88% and those in Groups E and F by 61% and 23% respectively. ANOVA and Bonferroni comparison tests revealed that Groups A and B were significantly different from Group C ( $p < 0.001$ ). However, no significant difference existed between Groups A and B ( $p > 0.05$ ). Similarly, the results obtained from Model II showed no significant difference between Groups D and E ( $p > 0.05$ ). However, there was a statistically significant difference when Groups D and E were compared with Group F ( $p < 0.001$ ). **Conclusions:** Both 10-day and 7-day pH-cycling models were suitable for studying carious lesion progression in primary teeth (the demineralizing and remineralizing solutions of the 10-day cycling model contained 0.25ppm F). A pea-sized portion (0.32 grams) of 500ppm F toothpaste could slow down the demineralization progression better than a half-pea-sized portion.

[Seq #348 - Dentifrice](#)

2:00 PM-4:00 PM, Saturday, 12 March 2005 Baltimore Convention Center  
Exhibit Hall E-F

[Back to the Cariology Research Program](#)

[Back to the IADR/AADR/CADR 83rd General Session \(March 9-12, 2005\)](#)