



UNIVERSITY OF HONG KONG
LIBRARY

CLINICAL REPORT
OF THE
TSAN YUK HOSPITAL
FOR THE YEAR
1956



ISSUED BY THE
DEPARTMENT OF OBSTETRICS AND GYNAECOLOGY
UNIVERSITY OF HONG KONG

CLINICAL REPORT
OF THE
TSAN YUK HOSPITAL
FOR THE YEAR
1956



ISSUED BY THE
DEPARTMENT OF OBSTETRICS AND GYNAECOLOGY
OF THE
UNIVERSITY OF HONG KONG

PRINTED BY
THE SOUTH CHINA MORNING POST, LIMITED.
HONG KONG.

CONTENTS

	<i>Pages</i>
1. MEDICAL STAFF	1
2. ABBREVIATIONS	3
3. PREFACE	5
4. COMPARATIVE TABLE	8

SECTION I—THE OBSTETRIC REPORT, 1956

5. STATISTICAL SUMMARY	9
6. DETAILED OBSTETRICAL REPORT	

ARRANGEMENT OF TABLES:—

Table 1. Associated Diseases and Conditions not Detailed Elsewhere	10
2. Abortion	11
3. Cardiac Disease	13
4. Pre-eclampsia	16
5. Hypertension—Essential	26
6. Chronic Nephritis	31
7. Eclampsia	32
8. Accidental Antepartum Haemorrhage ..	33
9. Placenta Praevia	35
10. Hydramnios	37
11. Tumours Associated with Pregnancy, Labour or Puerperium	38
12. Prolonged Labour	39
13. Persistent posterior position of the Occiput and Transverse Arrest	41
14. Uncomplicated Breech Deliveries	45
15. Complicated Breech Deliveries	49
16. Face and Brow Presentations	53
17. Transverse and Oblique Lie	54
18. Compound Presentation	56
19. Multiple Pregnancy	57
20. Labour following Previous Caesarean Section	60
21. Contracted Pelvis	61
22. Failed Forceps	71
23. Prolapse and Presentation of Cord	72
24. Post-partum Haemorrhage	74
25. Manual Removal of the Placenta	78
26. External Version before Labour	81
27. Induction of Labour	83
28. Caesarean Section	87
29. Forceps Deliveries	96
30. Ruptured Uterus	103
31. Version in Labour	104
32. Embryotomy and Craniotomy	105
33. Puerperal Pyrexia: Genital Infection	106
34. Puerperal Pyrexia: Extra-genital Infection	109

CONTENTS – (Cont'd)

	<i>Pages</i>
7. MATERNAL MORTALITY	119
8. DETAILS OF MATERNAL DEATHS	120
9. SPECIAL CASE	123

SECTION II — THE PAEDIATRIC REPORT, 1956

10. PAEDIATRIC REPORT	124
11. STATISTICAL SUMMARY	128

ARRANGEMENT OF TABLES:—

Table 35. Stillbirths	126
36. Neonatal Deaths	134
37. Premature Live Born Infants	139
38. Congenital Malformations	152
39. Birth Trauma	154
40. Neonatal Infections	156
41. Other Neonatal Diseases	158

12. X-RAY DEPARTMENT STATISTICS	162
---	-----

MEDICAL STAFF

Professor of Obstetrics & Gynaecology (Oct. 1938—Nov. 1956)
Gordon King, O.B.E., F.R.C.S. (Eng.), F.R.C.O.G.

Visiting Professor of Obstetrics & Gynaecology
(Dec. 1955—May, 1956):

Nicholson J. Eastman, M.D.
Professor of Obstetrics, Johns Hopkins University
and
Obstetrician-in-Chief to Johns Hopkins Hospital, U.S.A.

Senior Lecturer

Daphne Chun, M.B., B.S., F.R.C.S. (Ed.), M.R.C.O.G.

Lecturers:

Ada S. H. Wong, M.B., B.S., M.R.C.O.G.
Y. K. Tsao, M.B., B.S., M.R.C.O.G.
Christina Chow, M.B., B.S., M.R.C.O.G.

Assistant Lecturers:

K. K. Chow, M.B., B.S.
P. S. Kan, M.B., B.S.

Medical Officer-in-Charge:

H. Y. Cheng, M.B., B.S.

Clinical Assistants:

Samuel Yen, M.B., B.S.
W. H. Ko, M.B., B.S.
David Tse, M.B., B.S. (July-Dec., 1956).
Yu Chi Chiu, M.B., B.S. (July-Dec., 1956).

Senior House-Officers:

Mary Wong, M.B., B.S.
Esther Tam, M.B., B.S. (July-Dec., 1956).

House-Officers:

Kok Sau Hor, M.B., B.S.
Esther Tam, M.B., B.S. (Jan.-June, 1956).
Grace Lim, M.B., B.S.
David Tse, M.B., B.S. (Jan.-June, 1956).
Ronald Perry, M.B., B.S.
Yu Chi Chiu, M.B., B.S. (Jan.-June, 1956).
Ue Shuk Yuen, M.B., B.S.
Sylvia Loo Siew Wah, M.B., B.S.
Therese Lu Ti Li, M.B., B.S.
Tso Hin Yip, M.B., B.S.
Ng Chuan Wai, M.B., B.S.
Cheung Pak Chue, M.B., B.S.

ABBREVIATIONS

Ab.	Abortion
A.C. Head	After coming head
A.P.H.	Ante-partum haemorrhage
A.R.M.	Artificial rupture of membranes
B.	Booked case
B.B.A.	Born before arrival
C.	Child
C.P.	Contracted pelvis
C.S.	Caesarean Section
C.C.S.	Classical Caesarean Section
D.	Died
D.A.A.	Discharged against advice
D-S Catheter	Drew-Smythe's catheter
D.T.A.	Deep transverse arrest
E.C.S.	Extraperitoneal Caesarean Section
F.	Female
F.D.	Fully dilated
I.D.I.	Induction delivery interval
I.P. Eclampsia	Intra-partum eclampsia
I.S.	Interspinous diameter
I.U.D.	Intra-uterine death
I.V.	Intravenous
L.	Live
L.B.	Live-born
L.F.	Low forceps
L.S.C.S.	Lower segment Caesarean Section

M.	Mother
M.	Male
Mac.	Macerated
M.F.	Mid forceps
M.R.P.	Manual removal of placenta
M.S.V.	Mauriceau-Smellie-Viet
M.R. + M.F.	Manual rotation & mid forceps
M.R. + L.F.	Manual rotation & low forceps
N.A.D.	Nothing abnormal detected
N.B.	Non-booked case
N.D.	Neo-natal death
O.B.E.	Oil, bath and enema (Medical induction of labour)
O.C.	Obstetrical conjugate
O.P.	Occipito-posterior
P.E.	Pre-eclampsia
P.P. Eclampsia	Post-partum eclampsia
P.P.H.	Post-partum haemorrhage
P.P.S.	Post-partum sterilization
Q.M.H.	Queen Mary Hospital
T.D.	Transverse diameter of inlet
S.B.	Stillbirth
Spont.	Spontaneous delivery
T.B.	Tuberculosis
wt.	Weight

PREFACE

The Report for the year 1956 shows that the figures for admission and delivery surpass those of any previous year in the history of the Tsan Yuk Hospital. There were no fewer than 8,391 admissions and 7,552 deliveries during the year, thus surpassing by a considerable margin the previous record figures of 1953, when there were 7,221 admissions and 6,817 deliveries. A very considerable strain was therefore imposed upon the new 200-bed hospital thus early in its existence.

A perusal of the statistics presented in this year's report shows that the pattern of work during 1956 followed very closely that of previous years. Some 97% of the patients received antenatal care, and the average number of visits paid by each patient was 5.3. In spite of this, however, pregnancy toxæmia remained one of the big problems in the care of the patients, there being a 5.2% incidence of this condition (though admittedly a number of these cases came from the ranks of the unbooked cases). Altogether 373 operative deliveries were carried out during the year, representing a 2.6% incidence of forceps delivery and a 2.3% incidence of Caesarean Section. There were 4 maternal deaths (an incidence of 0.52 per thousand live and still births), the causes of death being antepartum haemorrhage (1), post-partum haemorrhage (1), acute hepatic necrosis (1) and perforation of the aorta into the oesophagus (1).

An outstanding event of the year was the visit of Professor Nicholson J. Eastman, who came to Hong Kong as Visiting Professor of Obstetrics and Gynaecology from January to June, 1956. Dr. Eastman's visit was a source of great inspiration to all members of the staff, and will long be remembered by all who came into contact with him during this period. The sincere thanks of the Department and of the University are due to the China Medical Board, through whose generosity the visit was made possible.

I feel that this Report must end on a personal note, since it marks the close of my association with the University of Hong Kong, where I have held the Chair of Obstetrics and Gynaecology from October 18th, 1938 to November 22nd, 1956—a period of over 18 years. During this time many changes affecting the Hospital have taken place in Hong Kong. Towards the end of 1938, as a result of the Sino-Japanese hostilities, the first big influx of population began, with the consequence that the number of deliveries in the Hospital increased abruptly from an average of between 2,000 and 2,500 per annum during the years 1933 to 1938 to figures in excess of 3,000 during the years 1939 to 1941. At the end of 1941, when the Second World War spread to the Far East, the work of the Hospital was seriously affected and its doors were actually closed to patients during a portion of the period of the Japanese occupation. In September, 1945, within three weeks of the re-occupation of Hong Kong, the hospital was again functioning and the number of patients admitted has progressed from year to year until the present astronomical figures have been reached. The long cherished dream of a new hospital first began to take form early in 1952, when the Hong Kong Jockey Club made its generous decision to devote the sum of three-and-a-half million dollars to the building of a new Tsan Yuk Hospital. In October, 1952 the Foundation Stone of the new

hospital was laid by Her Royal Highness the Duchess of Kent, and the dream of a new Hospital became translated into reality in June 1955, when the new building was officially opened by His Excellency Sir Alexander Grantham.

Throughout the years, since Professor Totterham first established the relationship between the Hospital and the University in 1926, the Tsan Yuk Hospital has offered unrivalled facilities for the training of medical students and midwives, and it is probably true today to say that, on account of its unique position, the hospital is capable of giving as extensive a training in the modern art of obstetrics to both undergraduate and postgraduate students as any other obstetrical hospital in the world.

During the years since its inception in 1922, the Tsan Yuk Hospital has been fortunate in the long succession of devoted workers who, from the time of Dr. Alice D. Hickling and Miss S. C. Leung onwards, have given of their best in order to enhance the welfare of the hospital. Throughout the 35 years of its history, the Hospital has provided an excellent example of what can be accomplished by a relatively small staff of doctors, nurses and students imbued with the team spirit. I should like to record here my immense gratitude to Matron S. C. Leung, M.B.E., who, throughout the first 30 years of the life of the hospital, until her retirement in 1952, was the moving spirit in everything that affected the well-being of the Hospital. In her retirement she still lives for the Hospital, venerated by successive generations of pupil midwives, medical students, doctors and nurses alike. Miss Leung was ably succeeded by Miss B. Thom, the present matron, who shouldered the immense task of transferring and organising the greatly enlarged nursing and domestic staff required by the new Hospital. To Miss Thom and her whole staff I acknowledge my very great indebtedness. I think with gratitude, also, of my pre-war Assistants, Dr. H. C. Ku, Dr. C. K. Quek and Dr. K. Yang and also of a whole host of house officers, who are now scattered over a very wide area of the Far East and many other parts of the world. To the present members of the Department I wish to tender my warm and sincere thanks for their loyal and devoted service, particularly in the difficult days of the post-war reconstruction, and in this I wish to pay special tribute to the sterling service rendered by Dr. H. Y. Cheng, who has been the Medical Officer-in-Charge of the Hospital since the post-war reorganisation in September, 1945. It has been a matter of particular gratification to me to note the increasing number of former students and house officers who have decided to specialise in Obstetrics and Gynaecology, and who have been successful in obtaining the Membership of the Royal College of Obstetricians and Gynaecologists. It is indeed encouraging to realise that the Department is now almost entirely staffed by former graduates of the University of Hong Kong, who have received the basic portion of their own training within the Department.

In conclusion, I should like to say how delighted I am that Dr. Daphne Chun has been appointed by the University as my successor in the Chair of Obstetrics and Gynaecology. Dr. Chun first entered my Department in the year 1940 and since that date has been a most loyal and energetic member of the staff, carrying a very heavy load of clinical responsibility during the post-war years, at a time when the Department

was grossly understaffed. I look forward with confidence and with pleasure to the future progress of the Department under the guidance of Professor Daphne Chun.

A parting of the ways is inevitably tinged with sorrow, and it is with very sincere regret that I say farewell to my friends of many years in Hong Kong. But I shall always follow with the keenest interest the progress of the Department and of the Hospital under its new and younger leadership.

GORDON KING.

October, 1957.

COMPARATIVE TABLE

	1945	1946	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956
<i>Maternal Mortality (per 1,000 deliveries)</i>												
<i>Booked</i>	—	—	—	0.68	1.61	0.22	1.29	0.31	0.45	0.77	0.16	0.24
<i>Non-booked</i>	5.10	—	2.45	—	7.84	0.18	—	11.45	5.99	13.07	—	10.20
Toaemia (No. of maternal deaths)	—	—	—	—	3	—	—	3	—	2	—	—
Eclampsia (No. of maternal deaths)	—	—	1	—	5	1	3	2	1	1	—	1
Placenta Praevia (No. of maternal deaths)	—	—	1	—	—	—	—	—	—	1	—	—
Morbidity	10.89%	3.74%	2.86%	2.26%	1.51%	1.59%	0.81%	1.20%	0.84%	0.98%	1.97%	5.73%
Forceps	3.56%	2.83%	1.91%	2.11%	2.91%	1.67%	1.56%	1.60%	1.95%	1.73%	1.73%	2.66%
C.S. Rate	—	—	0.08%	0.64%	0.58%	1.06%	0.88%	1.29%	1.03%	1.63%	2.53%	2.30%
No. of Deliveries	337	2,757	3,925	4,549	4,301	5,085	5,819	6,737	6,817	6,606	6,197	7,552

STATISTICAL SUMMARY

	Booked	Non-Booked	Total
Adult patients admitted	8,165	226	8,391
Patients delivered in hospital :			
Primigravidae	1,800	67	1,878
Multiparae	5,523	123	5,646
Patients admitted after delivery :			
Primiparae	1	2	3
Multiparae	22	3	25
Total patients delivered	7,357	195	7,552
Abortions	30	2	32
Patients transferred	37	3	40
Patients discharged undelivered	248	11	259
Maternal deaths	2	2	4
Maternal death rate (per 1,000 live and still-births)			0.523
Infants born in hospital	7,412	195	7,607
Infants born before admission (B.B.A.)	23	5	28
Total infants born	7,435	200	7,635
Infants transferred	16	—	16
Stillbirths	99	20	119
Stillbirth rate (per 1,000 live and stillbirths)			15.59
Neo-natal deaths	130	23	153
Neo-natal death rate (per 1,000 live births)			20.04
Ante-natal Clinic :—			
New patients		9,452	
Re-attendances		29,727	
Total		39,179	
Post-natal Clinic :—			
Total attendances		3,962	

TABLE 1

**ASSOCIATED DISEASES & CONDITIONS NOT DETAILED
ELSEWHERE**

	<i>Booked</i>	<i>Non-Booked</i>	<i>Total</i>
Abdominal pregnancy	1	-	1
Abscess, breast	2	-	2
Abscess, buttock	2	-	2
Amoebiasis	5	-	5
Anaemia, Iron deficiency due to ankylostomiasis	63	8	71
Anaemia, Megaloblastic of pregnancy	2	-	2
Asthma	5	1	6
Avitaminosis	143	2	145
Bacillary dysentery	2	-	2
Bartholin cyst	1	-	1
Bronchiectasis	3	1	4
Congenital recto-vaginal fistula	1	-	1
Dextrocardia	1	-	1
Diabetes mellitus	1	-	1
Empyema	1	-	1
Filariasis	1	-	1
Gonococcal infection	1	-	1
Haematoma of vagina	2	-	2
Hernia, diaphragmatic	3	-	3
Hernia, ventral	1	-	1
Homologous serum jaundice	1	-	1
Hypothyroidism	2	-	2
Hydronephrosis & ureteric stone	2	1	3
Infective hepatitis	3	-	3
Leprosy	2	-	2
Mumps	1	-	1
Naegele's pelvis	5	1	6
Nasopharyngeal Carcinoma	1	-	1
Nontoxic nodular goitre	3	-	3
Paratyphoid C.	1	-	1
Pleural effusion	2	-	2
Postpartum psychosis	3	-	3
Previous pelvic floor repair	1	-	1
Prolapsed uterus	5	-	5
Renal calculi	2	-	2
Septate uterus	1	-	1
Syphilis	273	9	282
Thrombocytopenic purpura	1	-	1
Thrombophlebitis	1	-	1
Thyrototoxicosis	-	1	1
Tuberculous meningitis	1	-	1
Tuberculous pneumonia	1	-	1
Typhoid fever	1	-	1
Urinary infection	39	1	40
Uterus bicornis unicollis	1	-	1
Vaginal septum	6	-	6
Varicose veins	197	1	198

TABLE 2

ABORTION

There were 30 Booked and 2 Non-Booked cases. No maternal death. One case had morbidity of one day. All, except one*, terminated spontaneously.

I). Parity— (Booked and Non-Booked)

Primigravidae	5	(15.62% of all abortions)
Parous	27	(84.38% of all abortions)

II). Types of Abortion:— (Booked and Non-Booked)

Complete	22
Missed	10
Therapeutic	0
Threatened	0

III).

	Spontaneous delivery	Threatened abortion Discharged undelivered	Missed abortion	Therapeutic vaginal delivery	Therapeutic abdominal delivery	Habitual abortion	Total abortion
Booked	16	---	10	---	---	1	---
Non-Booked	2	---	---	---	---	---	---

* Hosp. No. 2389/56N.B.

ABORTION (Continued)

Reg. No.	Age	Pregnancies Before 28 wks	Pregnancies After 28 wks	Maturity	If induced Indication of Method	If Spontaneous cause (if known) and treatment	Result to Mother	Morbidity	Remarks
BOOKED									
311	26	—	3	?	—	Spont.	L	—	Missed abortion
1130	37	2	2	?	—	"	L	—	Missed Abortion, B.B.A. Twins
1222	29	—	1	27	—	"	L	—	Advanced maceration Blood loss 900 c.c.
2080	28	—	3	23-2	—	"	L	—	Blood transfusion 500 c.c.
2399	30	1	7	24	—	"	L	—	Complete abortion
2523	34	1	2	25	—	"	L	—	Missed abortion
2600	35	—	2	18-6	—	"	L	—	Missed abortion, Essential Hypertension
3090	29	1	6	17	—	"	L	—	Complete abortion Placental tissue removed by ovum forceps
3187	24	—	2	20	—	"	L	—	Missed abortion
3203	33	1	4	25-2	—	"	L	—	Complete abortion
3326	32	—	4	14	—	"	L	—	"
3376	38	1	4	10-6	—	"	L	—	"
3647	38	—	5	22	—	"	L	—	"
3677	28	—	2	25-6	—	"	L	—	"
3786	32	—	3	30-4	—	"	L	—	"
3855	22	—	—	?	—	"	L	—	Missed abortion
4067	37	11	1	24-5	—	"	L	—	"
4887	26	—	—	25-6	—	"	L	—	Complete abortion, habitual Missed abortion
4900	32	—	3	20	—	"	L	—	"
5538	41	2	10	19-4	—	"	L	I	Complete abortion
5559	37	—	3	24	—	"	L	—	Mitral Stenosis P.P.S.
5862	29	—	1	26-1	—	"	L	—	Uniovular twins
6110	41	1	1	25-5	—	"	L	—	Complete abortion
6112	23	—	—	26-3	—	"	L	—	"
6202	37	—	4	22	—	"	L	—	"
6753	27	—	1	20-2	—	"	L	—	Complete abortion
6904	29	—	1	?	—	"	L	—	Hydramnios 2,700 c.c.
7089	40	—	6	?	—	"	L	—	Aneurysmal Missed abortion
7719	26	—	2	20	—	"	L	—	"
8137	30	—	5	?	—	"	L	—	"
NON-BOOKED									
2389	19	—	—	?	—	Spont.	L	—	Complete abortion
7456	45	1	3	27-1	—	Manual Removal Spont.	L	—	Complete abortion

TABLE 3

CARDIAC DISEASE

There were 58 cases, an incidence 0.69%, 56 Booked and 2 Non-booked.

There were no maternal deaths.

There were 2 neo-natal deaths and one stillbirth.

There was one spontaneous abortion.

There were 8 patients in failure on admission.

Three patients were sterilised.

13 cases were admitted during pregnancy. One was admitted at 17th wk., 2 at 26th wk., 1 at 28th wk., 1 at 32nd wk., 2 at 36th wk., 1 at 37th wk., 2 at 38th wk., 1 at 39th wk., and 2 at 40th wk., respectively.

1). Nature of heart lesion:—

	Booked	Non-Booked
Mitral Stenosis	45	1
Mitral Stenosis + Aortic Regurgitation	11	1

2). Method of Delivery:—

Method of Delivery	Booked	Non-Booked
Spontaneous	49	2
Bipolar Version	1	
Mid Forceps	3	
Manual Rotation & Mid Forceps	1	
Abortion	1	
Undelivered	1	
TOTAL	56	2

CARDIAC DISEASE (Continued)

Reg. No.	Age	Pregnancies Before 28 wks	Pregnancies After 28 wks	Maturity	Lesion	Degree of failure of compensation (Class 1, 2, 3 or 4 N.Y. Heart Assoc. At Admission Delivery)	Method of Delivery	M.	Result	Remarks
BOOKED										
6811/55	25	-	-	40-6	Mitral Stenosis	I	I	L	L	D.A.A.
6827/55	27	-	1	27-4	"	I	"	L	L	
113/56	30	-	4	39-6	Mitral Stenosis	IV	IV	L	N.D.	Congestive Heart Failure
243	36	-	2	39	Aortic Regurgitation	I	I	L	L	
518	27	-	2	39	Mitral Stenosis	I	I	L	L	
520	36	-	3	38-3	"	I	"	L	L	
524	38	-	2	38	"	IV	IV	L	L	
558	36	-	5	37-6	Mitral Stenosis	I	I	L	L	
660	39	1	6	33-3	Aortic Regurgitation	IV	IV	L	S.B.	Prolapsed Cord, Manual Removal of Placenta
852	23	-	-	40-3	Mitral Stenosis	I	I	L	L	
967	34	-	3	35	"	IV	IV	L	L	
994	37	-	6	37-1	"	I	"	L	L	
1620	29	1	3	38-4	"	I	I	L	L	
1655	34	-	3	39-5	"	II	II	L	L	
2224	36	1	3	41	Mitral Stenosis	II	II	L	L	
2258	30	2	3	3	"	I	"	L	L	
2333	20	-	-	43	"	I	"	L	L	
2383	29	-	2	41-5	"	IV	IV	L	L	
2522	29	-	1	?	"	III	III	L	L	
2942	38	-	2	37-3	"	IV	IV	L	L	
2980	29	1	4	39-3	"	IV	IV	L	L	
2991	41	3	4	39	"	IV	IV	L	L	
3106	36	-	5	38-4	"	I	"	L	L	
3162	33	1	1	41-1	"	I	"	L	L	
3548	38	-	7	38	"	I	"	L	L	
3585	34	1	5	38-2	Mitral Stenosis	II	II	L	L	
3767	26	1	1	3	Aortic Regurgitation	I	I	L	L	Well compensated. D.A.A.
3796	29	-	1	36-6	Mitral Stenosis	II	II	L	L	
4011	26	-	2	33-5	"	I	"	L	L	
4041	31	3	1	38-2	"	II	II	L	L	
4056	42	-	7	40-2	Mitral Stenosis	II	II	L	L	
4979	34	1	1	46-4	Aortic Regurgitation	II	II	L	L	Developed Congestive Heart Failure after Delivery
5010	29	-	2	41	"	II	"	L	L	

CARDIAC DISEASE (Continued)

Reg. No.	Age	Pregnancies 28 wks Before	Maturity	Lesion	Degree of failure of compensation (Class 1, 2, 3 or 4 N.Y.) (Heart Assoc. 1939)	Method of Delivery	Result C.	Remarks
5081	39	1	37-3	Mitral Stenosis	I	"	L	Compensated. P.P.S.
5245	33	1	30-3	Mitral Stenosis	I	Spont.	L	
5330	30	1	39-4	"	IV	M.R. + M.F.	L	
5558	36	1	33	Mitral Stenosis	III	Spont.	L	
5559	37	1	24	Aortic Regurgitation	I	"	L	Abortion
5737	28	1	39-1	Mitral Stenosis	I	"	L	
5803	25	1	39-3	Mitral Stenosis	II	"	L	
5874	29	1	39	Aortic Regurgitation	I	"	L	
5999	26	2	38-2	Mitral Stenosis	II	"	L	
6052	21	1	35-1	"	I	"	L	
6164	31	5	41-4	"	II	"	L	
6324	33	2	42	"	I	"	L	
6498	28	1	40-1	"	III	"	L	
6900	23	1	38-2	"	II	"	L	
7054	32	1	41-1	Mitral Stenosis	III	"	L	
7154	30	1	22-2	Aortic Regurgitation	IV	Undelivered	L	
7209	23	2	37-1	Mitral Stenosis	II	Spont.	L	
7607	29	1	38	Aortic Regurgitation	III	"	L	
7613	36	8	38-1	Mitral Stenosis	I	"	L	
7678	29	3	41-5	"	III	"	L	
7880	35	5	37-5	"	I	"	L	
7965	29	2	38-2	"	II	"	L	
8338	32	3	41-3	"	I	"	L	
NON-BOOKED								
1400	32	5	40	Mitral Stenosis	IV	Spont.	L	
1560	32	1	41-6	Aortic Regurgitation	IV	"	L	

P.P.S. compensated
P.P.S.
Compensated
Congestive Heart Failure. P.E.
D.A.A.
Congestive Heart Failure. Acute
Allergic Dermatitis. Infectious
Mononucleosis. Transferred to
Q.M.H.
Congestive Heart Failure
P.P.S.
Compensated

TABLE 4

PRE-ECLAMPSIA

There were 418 cases of pre-eclampsia, an incidence of 4.98%, 382 Booked and 36 Non-booked. There were 12 stillbirths (4 macerated), an incidence of 2.91%, 8 in Booked and 4 in Non-booked cases; 14 neo-natal deaths (4 twins), an incidence of 3.39%, 11 Booked and 3 Non-booked.

There were 24 sets of twins, an incidence of 5.81%, 21 in Booked and 3 in Non-booked cases. Four cases were discharged undelivered.

Reg. No.	Age	Pregnancies Before 28 wks	Maturity On Ad-At De-mission livery	Albuminuria On Maximum Discharge	Oedema	Height Blood Pressure	No. of Days in hospital before labour or discharge	If labour indicated in State method	Method of Delivery	Result M. C.	Weight of Child gm.	Remarks
BOOKED												
6583/55	35	—	6 36-3	40 None	Slight	160/110	25 Days	—	Spont.	L	3120	Twin Pregnancy
6783/55	38	—	3 35-6	38-5 Trace	Severe	150/110	20 "	—	"	L	2950	
6787/55	24	—	— 42-3	44-1 None	Slight	160/90	12 "	O.B.E.	M.R. + M.F.	L	3300	Indication of Induction: Post-maturity
3/56	35	—	4 42-2	42-2 Cloud	"	160/105	—	—	Spont.	L	4320	
56	28	—	2 41-3	41-3 Trace	"	170/110	—	—	"	L	2920	
155	40	—	9 43-6	43-6 None	"	165/100	—	—	"	L	3130	
182	41	—	9 38-3	38-3 "	"	165/95	—	—	"	L	2920	
192	33	1	6 23-5	undelivered	"	140/90	5 Days	—	Undelivered	L		Undelivered Admitted for Investigation and Treatment
212	38	—	1 38-4	39-1 Trace	"	150/100	4 "	—	Mid Forceps	L	2650	
215	35	—	6 36-1	37-1 None	"	160/110	7 "	—	Spont.	L N.D.	2200	
223	29	—	1 37-3	37-3 Trace	"	160/90	—	—	"	L	3100	
229	34	1	— 38-5	38-5 Trace	"	160/105	—	—	"	L	2650	
237	18	—	5 38-4	38-4 None	"	160/100	—	—	L.S.C.S.	L L	2948	
{ 6841/55	42	1	5 38-6	38-6 "	"	150/105	—	—	Spont.	L L	3390	Admitted at 35-4/7 wks. for Investigation and Treatment
291	32	—	1 39-4	39-5 "	"	140/105	—	—	"	L N.D.	2200	
326	22	—	1 37-2	37-6 "	Moderate	160/90	4 Days	—	Spont.	L	2730	Twin Pregnancy
329	41	—	4 40-4	41 Trace	Severe	155/100	3 "	—	"	L	2500	
350	27	2	1 37-1	37-3 "	Moderate	140/90	2 "	—	"	L	2890	Manual Removal of Placenta
391	33	—	1 34-4	36 "	"	150/100	10 "	—	"	L N.D.	1820	
393	44	—	6 34-6	38 Solid	Slight	105/110	22 "	—	"	L S.B.	2500	Pulmonary Tuberculosis
403	36	—	2 30-1	31-1 None	"	150/100	—	—	"	L	2400	
407	39	—	3 30	30-1 Cloud	"	140/95	1 Day	A.R.M.	Re-polar Version	L	2610	Indication of Induction: Accidental Haemorrhage
											1015	Craniotomy Before Extraction of After-Coming Head: Mixed Accidental Haemorrhage (900 c.c.)

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Maturity On Ad- mission	Albuminuria On Discharge	Oedema	Highest Blood Pressure	No. of Days in hospital before labour or discharge	If labour induced State method	Method of Delivery	Result	Weight of Child gm.	Remarks
425	27	—	3 40-2	None	Slight	160/95	—	—	Spont.	L	3280	
428	32	—	1 36-2	Trace	Slight	160/95	—	—	"	L	3250	
{ 113	22	—	1 39-4	None	"	160/100	—	—	"	L	3270	
{ 436	—	—	—	Trace	—	145/90	—	—	"	L	2977	
{ 6957/55	29	—	1 32-3	Trace	—	165/105	—	—	"	L	2977	Hydrops Foetalis Admitted at 36-5/7 wks. for Investigation and Treatment
{ 507/56	28	—	4 37-1	None	Slight	165/105	—	—	Spont.	L	2370	Admitted at 27-5/7 wks. for Investigation and Treatment
543	34	—	4 35-6	Trace	"	180/120	22 Days	—	L.S.C.S.	L	2387	Elderly Primigravida Contracted Pelvis
555	22	—	—	"	"	140/100	—	—	Spont.	L	3580	
557	28	1	2 42	"	"	185/105	—	—	"	L	3050	
559	34	—	4 38-1	"	Slight	140/100	7 Days	—	"	L	3487	Pulmonary Tuberculosis Admitted at 37-6/7 wks. for Investigation and Treatment
{ 6956/55	43	—	4 39-2	"	"	140/100	—	—	"	L	3440	Admitted at 33 and 37 wks. for Investigation and Treatment
{ 620/56	—	—	—	"	"	160/100	—	—	"	L	2800	
{ 6851/55	27	—	2 39-6	"	"	150/100	—	—	(1) Spont.	L	2980	Admitted at 37-3/7 wks. for Investigation and Treatment
{ 642/56	38	1	2 41	"	Severe	150/100	—	—	(2) Spont.	L	2100	Admitted at 37-1/7 wks. for Investigation and Treatment
649/56	44	—	10 39	"	Slight	160/100	2 Days	—	Spont.	L	3030	Manual Removal of Placenta
{ 663	—	—	—	"	"	160/120	—	—	"	L	3650	Admitted at 26-1/7 wks. for Investigation and Treatment
692	26	—	1 39-1	Trace	"	150/90	—	—	M.R. + M.F.	L	3232	Admitted at 26-1/7 wks. for Investigation and Treatment
{ 6577/55	26	—	1 36	"	"	145/100	—	—	Spont.	L	3280	Admitted at 38 wks. for Investigation and Treatment
{ 700	22	—	—	"	"	170/120	—	—	Transverse Lie	L	2665	Admitted at 36 wks. for Investigation and Treatment
{ 526	—	—	—	"	"	145/90	—	—	Breech Extraction	L	3020	Admitted at 34-3/7 and 36-4/7 wks. for Investigation and Treatment
{ 714	35	—	2 38-5	Cloud	"	145/105	—	—	Spont.	L	3232	
{ 734	24	—	40-4	Trace	"	165/90	—	—	"	L	2390	
737	24	—	39-6	Trace	"	200/115	3 Days	—	"	L	2990	
429	25	—	39-6	Trace	"	150/110	—	—	"	L	3100	
742	30	—	4 40	"	"	200/120	—	—	"	L	3120	
771	29	1	7 36-5	"	"	160/110	—	—	"	L	2985	Admitted at 36-1/7 wks. for Investigation and Treatment
793	17	—	40-6	Cloud	"	150/100	—	—	"	L	3420	Admitted at 35-2/7 wks. for Investigation and Treatment
{ 364	—	—	—	Trace	"	165/120	—	—	M.R. + M.F.	L	2400	Manual Removal of Placenta
{ 820	32	—	4 37-6	"	"	145/90	—	—	Low Forceps	L	2870	
823	28	—	1 40	"	"	145/90	—	—	Spont.	L	2400	
{ 338	—	—	—	Trace	"	145/90	—	—	"	L	2870	
{ 825	—	—	—	Trace	Moderate	145/90	—	—	"	L	2870	
835	23	—	41-3	Trace	"	145/90	—	—	"	L	2870	
836	32	—	2 39-6	Trace	"	145/90	—	—	"	L	2870	
846	31	—	10 39-6	"	"	145/90	—	—	"	L	2870	

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks	Maturity On Ad. - At. De-mission	Albuminuria On Maximum Discharge	Oedema	Highest Blood Pressure	No. of Days in hospital before labour or discharge	If labour induced State method	Method of Delivery	Result	Weight of Child gm.	Remarks
{ 103 } { 848 } { 702 } { 618 }	22 35 30	— 1 —	39 6 40 40 2	Trace None Trace	Slight Moderate Slight	150/105 170/100 140/100	— 2 Days 5 "	— — —	Spont. " Spont.	L L L	3380 3230 2870	Admitted at 33 3/7 wks. for Investigation and Treatment Admitted at 33-2/7 wks. for Investigation and Treatment
939 994	30 37	— 1	40 6 37-1	Trace	"	150/90 150/95	3 Days —	— —	Spont. Spont.	L L	3115 2700	Mitral Stenosis—Class I. Admitted at 29 wks. for Investigation and Treatment Manual Removal of Placenta Cervical Dystocia, Contracted Pelvis
995 1005	22 18	— —	43 36 4	— Cloud	" "	145/105 140/90	— 2 "	— —	M R. + M.F. L.S.C.S.	L L	3160 2980	Admitted at 33 wks. for Investigation and Treatment Hypotonic Uterine Inertia
1060 1089	30 32	— —	39 3 40 3	— Cloud	Moderate	175/100 200/120	— —	— —	Spont. L.S.C.S.	L L	2190 2800	
1137 1156 1176	25 27	— —	40 5 34-1	— —	" Slight	140/100 145/95	— —	— —	Spont. "	L L	3780 3070	
1206 1208 { 1012 } { 1227 }	37 35 35	— 1 — —	39 3 38 6 39 2 41 2	Trace Trace —	Moderate Slight Moderate "	160/90 150/100 150/100	1 Day 2 Days 1 Day —	— — — —	Spont. " " "	L L L L	3350 2720 2840 4140	Admitted at 38 6/7 wks. for Investigation and Treatment
1261	33	1	36 3	Trace	Severe	210/120	2 Days	Low Rup-ture of Membranes	"	L	2150	
1265 1226 1297	29 41 36	— 1 3	41 2 37 4 39 5	Trace Trace	Slight Slight Moderate	150/95 155/90 150/100	— — 1 1/2 Days	— — —	Spont. Spont.	L L L	3450 3460 2700	P.P.S.
1300 1366 1407	35 24 20	— — —	38 3 41 2 37	" "	" Slight	160/100 165/105 145/95	— — —	— — —	" " "	L L L	3000 3120 3162	
1421 1438	30 29	— —	38 1 37 5	Cloud Cloud	Severe Slight	190/100 180/120	5 "	— —	L.S.C.S. Spont	L L	2807 2812	
1475 1484	30 40	— —	38 6 40 5	Trace	Severe Slight	170/110 140/90	6 " 2 "	— —	" "	L L	2480 2500	
1523 { 1216 } 1352	40 32 26	— — —	38 2 40 4 40 4	Trace Trace	" Moderate	140/100 150/100 150/100	— 4 " —	— — —	" " "	L L L	3700 3700 3700	Admitted at 37 4/7 wks. for Investigation and Treatment Pulmonary T.B. Pulmonary T.B.
1568 1602	34 40	— —	38 2 39 6	Trace Cloud	Slight "	160/100 160/110	— 5 "	— —	" "	L L	3700 3500	

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks	Maturity On Ad- At De- mission livery	Albuminuria On Maximum Discharge	Oedema	Highest Blood Pressure	No. of Days in hospital before labour or discharge	If labour induced State method	Method of Delivery	Result	Weight of Child gmt.	Remarks
2448	32	—	38-1	Trace	Slight	155/110	—	—	Spont.	L	3080	
2451	31	—	38-1	Solid	Severe	210/120	7 Days	Stripping of Membranes	"	L	2665	
2452	34	1	34-6	Trace	"	140/100	18 "	—	(1) Vx-Spont. (2) Vx-Spont.	L	2480	Twin Pregnancy
2472	33	—	34-3	Trace	Moderate	140/100	—	—	"	L	3295	
2508	28	1	42	"	"	160/115	1 Day	—	"	L	3070	
2511	37	6	42-1	"	Slight	190/105	—	—	"	L	3372	Pulmonary T.B.
2572	33	—	40-5	"	Moderate	160/90	—	—	"	L	3070	
2614	41	4	38-6	—	"	195/90	—	—	"	L	3370	
2615	39	6	39	Trace	"	165/105	—	—	"	L	2880	
2617	24	—	39-4	Cloud	Moderate	170/105	—	—	(1) Vx-Spont. (2) Br-Spont.	L	2840	Admitted at 35-5/7 wks. and 38-1/7 wks. for Investigation and Treatment. Twin Preg-nancy
2037	38	—	43	—	"	180/90	—	—	"	L	2520	
2663	32	—	39-6	Trace	"	160/90	—	—	Spont.	L	2890	
2688	25	—	36-6	—	Severe	160/115	15 Days	—	"	L	2530	
2281	43	—	39-5	—	"	160/90	2 "	—	"	L	2840	Admitted 36-2/7 wks. for In-vestigation and Treatment.
2754	31	—	35-4	Trace	"	160/106	—	—	"	L	3530	
2775	33	—	40-4	Trace	Slight	155/100	—	—	"	L	3600	
2776	33	—	39-5	Trace	Moderate	170/110	—	—	"	L	3440	
2785	33	—	38-6	Trace	"	140/90	—	—	"	L	3280	
2803	25	—	36-4	—	Slight	160/100	—	—	"	L	3420	Admitted at 34-1/7 wks. for In-vestigation and Treatment.
2478	32	1	36-4	—	"	160/100	—	—	"	L	3890	
2828	42	3	38-5	None	Moderate	150/100	15 Days	—	"	L	2700	Admitted at 27-6/7 wks. for Investigation and Treatment.
2841	19	—	30-6	Trace	Slight	160/110	2 "	—	"	L	2730	Admitted at 34-5/7 wks. for Investigation and Treatment.
2548	18	—	37	—	"	170/130	—	—	"	L	2470	Mitral Stenosis
2867	30	—	37-6	Trace	Slight	176/110	17 Days	—	Spont.	L	3280	
2942	38	—	37-2	None	Moderate	210/100	1 Day	—	Mild Forceps	L	3130	
2974	33	1	38-4	—	Slight	140/90	—	—	Spont.	L	3000	
2985	34	—	37-6	Trace	Slight	160/105	5 Days	—	"	L	3480	
2033	39	—	42-5	"	Slight	150/100	—	—	"	L	3400	Admitted at 36th & 40th week for Investigation and Treatment
2670	36	—	39	None	Moderate	150/90	—	—	"	L	3470	
3016	24	—	39-5	"	"	180/115	—	—	"	L	3180	
3037	28	—	38	Trace	Moderate	140/90	—	—	"	L	3800	
3095	38	—	35-3	"	Slight	220/125	3 Days	—	"	L	1900	
3053	28	—	40-6	"	Moderate	115/90	14 Days	—	"	L	2950	Manual Removal of Placenta
3062	28	—	40-6	None	Moderate	140/95	—	—	"	L	3710	
3114	25	—	40-6	"	"	180/100	—	—	"	L	3050	
3119	25	—	37-3	"	Slight	—	—	—	"	L	—	

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks	Maturity On Admission	Albuminuria On Discharge	Oedema	Highest Blood Pressure	No. of Days in hospital before labour or discharge	If labour induced	Method of Delivery	Result M. C.	Weight of Child gm.	Remarks
3137	20	1	38-5	None	Slight	160/100	—	—	Spont.	L	3620	
3139	25	—	39-4	Trace	Moderate	160/100	5 Days	—	Mid Forceps.	L	3050	Twin Pregnancy
3148	26	1	35-2	Trace	Moderate	150/110	5 "	—	Spont.	L	2000	
3155	28	—	36	None	"	140/90	5 "	—	Mid Forceps.	L	2180	
3166	28	—	38-2	"	Slight	150/100	—	—	Spont.	L	2466	
3169	36	—	?	"	"	190/115	—	—	"	L	2910	
{2551	37	—	39-2	"	"	170/100	—	—	"	L	2810	
{3184		3	39-2	"	"	170/100	—	—	"	L	2870	Admitted at 30-3/7 wks. for Treatment
2104	43	10	41-5	"	"	170/110	—	—	"	L	3700	Admitted at 33-6/7 wks. and 38-5/7 wks. for Investigation and Treatment
2780		—	?	"	"	170/110	—	—	"	L	2100	
3199	28	3	37-6	Trace	"	140/90	—	—	"	L	3550	
3207	30	1	37-2	Trace	"	180/100	—	—	"	L	3120	
3252	40	1	41-2	None	"	160/105	4 Days	—	"	L	3120	
3332	40	6	35-3	Trace	"	165/90	2 Days	—	"	L	2850	
3333	35	—	40-3	Cloud	"	170/110	1 Day	—	"	L	3100	
3339	21	—	40-4	None	"	175/100	1 "	—	L.S.C.S.	L	3060	
3368	32	1	37-6	Cloud	"	130/80	1 Day	—	Spont.	L	3020	
3382	26	—	33-6	Trace	"	175/105	2 Days	—	"	L	3160	
3427	24	—	39-5	None	Moderate	155/100	2 Days	—	"	L	2980	
3452	28	—	37-6	"	"	150/110	—	—	"	L	2850	
3472	45	1	32	"	"	150/110	5 Days	—	"	L	2990	
3504	21	1	39	None	Slight	150/100	—	—	Spont.	L	3100	D.A.A.
3506	26	—	36-4	Trace	"	140/100	—	—	"	L	3440	
3537	42	—	36-5	Trace	Moderate	190/120	2 Days	—	"	L	2840	
3554	28	—	38-3	None	"	150/190	—	—	"	L	2960	
3557	29	3	39	Trace	Slight	180/90	—	—	"	L	3300	
{2759	39	—	38-2	Cloud	"	190/125	3 Days	—	"	L	3170	Admitted at 33-3/7 and 36th wks. for Investigation and Treatment
{3181		3	38-2	Cloud	"	190/125	3 Days	—	"	L	3300	
{3572		5	38-5	Trace	"	190/120	2 "	—	"	L	3110	
3600	31	—	39-6	"	"	170/100	—	—	"	L	2560	Twin Pregnancy
3614	37	—	39-6	"	"	170/100	—	—	"	L	2750	
3699	32	—	36	Trace	Moderate	190/100	—	—	Spont.	L	3120	
3794	36	—	38-4	None	Slight	140/95	—	—	"	L	3750	
3833	43	4	39-3	"	Moderate	140/90	2 Days	—	Low Forceps	L	3570	
{3363	24	—	39-5	Trace	"	165/120	—	—	Spont.	L	3170	Admitted at 37 wks. for Investigation and Treatment
{3882		1	39-5	"	"	165/120	—	—	"	L	3170	
3901	31	—	36-3	None	Slight	180/110	17 "	—	"	L	3570	
3903	19	—	34-4	Cloud	"	200/130	—	—	"	L	2030	
3905	32	—	36-5	Trace	Severe	150/110	—	—	"	L	2450	Twin Pregnancy

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks	Maturity On Ad- mission	Albuminuria On Discharge	Oedema	Highest Blood Pressure	No. of Days in Hospital before labour or discharge	H labour induced by method	Method of Delivery	Result M. C.	Weight of Child gm.	Remarks
3943	28	2	42-2	None	Moderate	135/105	—	—	Spont.	L	3300	
3950	25	—	40	"	Slight	165/115	—	—	—	L	2665	
3975	33	—	38-6	"	"	165/105	—	—	"	L	3670	
3979	37	—	40-6	"	"	145/105	—	—	"	L	2640	
3990	31	—	38-4	"	"	155/110	—	—	"	L	3290	
4021	42	1	40-3	Trace	"	200/120	2 Days	—	"	L	3200	
4038	30	1	42-4	None	Moderate	160/100	—	—	"	L	2770	
4082	24	1	38-2	"	Slight	150/100	—	—	"	L	2650	
4083	24	—	36-5	Trace	Moderate	155/100	—	—	"	L	2610	
4082	23	—	34	"	"	180/120	44 Days	Stripping of Membranes	"	L S.B.	2050	
4084	25	—	37	"	"	140/110	14 Days	—	"	L	2800	
4096	25	—	40	None	Slight	140/100	—	—	"	L	3270	
4104	26	—	39	Trace	Moderate	155/105	—	—	"	L	2540	
4163	32	—	34-1	Solid	"	170/125	13 Days	—	L.S.C.S.	L	3150	
4181	29	—	38-2	Trace	Slight	150/100	5 "	—	Spont.	L	3180	
4183	24	—	38-1	None	Moderate	150/110	—	—	"	L	3020	
4198	34	—	39-2	Trace	Slight	160/90	—	—	L.S.C.S.	L	3150	
4202	24	—	?	None	Moderate	140/90	—	—	Spont.	L	2900	
4213	24	—	41	Trace	"	140/110	4 Days	Stripping of Membranes	"	L	3580	
{ 3928	31	—	37-6	"	Slight	160/105	—	—	"	L	3090	Admitted at 36-3/7 wks. for Investigation and Treatment
{ 4258	39	—	39-4	"	Moderate	145/95	—	—	"	L	3320	
4283	29	—	?	None	"	155/105	3 Days	—	"	L	3570	
4319	29	—	38-6	Trace	"	145/90	1 Day	—	"	L	3856	
4336	27	—	41-2	"	"	165/95	22 Days	O.B.E.	L.S.C.S.	L	3200	D.A.A.
4389	31	—	40-3	None	Slight	165/105	2 "	—	Spont.	L	2850	B.B.A.
4403	32	—	35-5	Trace	"	175/106	—	—	"	L	3100	
4417	26	—	39	Trace	"	170/100	—	—	Spont.	L	2400	
4495	24	1	35-3	Cloud	Severe	170/105	9 Days	—	"	L	3050	
4506	21	—	40	Trace	Moderate	170/100	5 "	—	"	L	2675	
4522	29	—	35-2	Trace	Slight	150/90	3 "	—	"	L	2050	
4529	39	—	37-2	Trace	Moderate	185/115	—	—	"	L	3080	Admitted at 39-4/7 wks. for Investigation and Treatment
3223	29	—	41-3	—	Slight	150/90	—	—	"	L	3700	
4564	28	—	36-2	Trace	"	160/100	—	—	"	L	3700	
4576	20	—	40-6	Trace	"	180/100	—	—	"	L	2608	Readmitted at Term with Normal B.P., No Oedema and No Albuminuria
4590	33	—	37-6	—	Severe	150/100	9 Days	—	Undelivered	L		
{ 4594	26	—	?	Trace	Slight	190/120	1 Day	—	Spont.	L	3100	
4598	27	—	46	"	"	160/100	—	—	"	L	3130	
4602	32	—	39-2	"	"	150/100	—	—	"	L	3090	
4607	27	—	41-6	"	"	160/90	—	—	"	L	3270	

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks	Maternity On Ad. At De-mission	Albuminuria On Maximum Discharge	Oedema	Highest Blood Pressure	No. of Days in hospital before labour or discharge	If labour induced State method	Method of Delivery	Result	Weight of Child gm.	Remarks
2570	36	—	40.3	—	None	140/90	—	—	Spont.	L	2680	Admitted at 38 5/7 wks. for Investigation and Treatment. Twin Pregnancy
4273										L	2710	
4610										L		
4623	24	1	?	Trace	"	160/110	5 Days	—	"	L	2620	
4661	26	—	40.2	—	"	150/105	—	—	"	L	2890	
4674	36	4	36.3	"	"	155/95	15 Days	—	"	L	3420	
4680	25	—	40.6	—	"	150/90	—	—	"	L	3600	
4693	25	—	42.3	Trace	"	150/100	—	—	"	L	3600	
4743	36	1	?	—	Moderate	140/100	5 Days	—	"	L	3820	
4745	31	1	40.4	—	"	160/115	1 Day	—	"	L	3700	
4769	29	—	38.2	Trace	Slight	130/90	3 Days	—	L.S.C.S.	L	3175	
4773	35	—	40.3	—	"	140/100	—	—	"	L	2700	
4782	24	—	40.4	"	"	160/110	—	—	"	L	2950	
4788	37	—	40.2	"	"	160/115	—	—	"	L	5560	Admitted at 38 wks. for Investigation and Treatment
4797	27	—	37.4	—	"	160/105	1 Day	—	(1) Vertex— M.F.	L	3180	Twin Pregnancy
4845	27	—	36.2	Trace	"	140/90	"	—	(2) Vertex— Breech Internal Version	L	1900	
4891	27	—	39.4	"	"	170/110	"	—	L.F.	L	3280	
4896	36	2	37.3	Trace	Moderate	180/120	—	—	Spont.	L	1900	
4908	25	—	39.5	—	"	150/100	—	—	"	L	2880	
4925	43	—	39.6	—	"	150/90	6 Days	—	"	L	3170	
4929	24	—	38.5	Cloud	"	150/110	8	—	M.F.	L	4200	
5013	36	—	39.3	—	"	160/100	1 Day	—	Spont.	L	3050	
5052	28	—	39.6	Trace	Slight	160/100	—	—	Spont.	L	3220	
5071	30	—	38.5	—	"	140/100	—	—	"	L	3750	
4597	39	—	41.1	—	"	150/90	—	—	"	L	2600	
5095						140/100	—	—	"	L	2975	Admitted at 39 wks. for Investigation and Treatment
5165	44	—	38	Trace	"	150/100	—	—	"	L	3250	
5202	29	—	36.3	"	Moderate	160/100	18 Days	—	"	L	2600	
5219	38	—	38.4	"	Slight	140/100	—	—	"	L	2600	
5251	35	—	39.3	Cloud	"	135/70	—	—	"	L	2975	
5268	37	1	41.3	Trace	"	150/110	1 Day	—	L.S.C.S.	L	2495	
5286	27	—	41.6	—	Moderate	160/110	—	—	Spont.	L	2780	
5318	38	—	41	—	Slight	140/90	—	—	"	L	2800	
5336	37	—	34.3	None	Severe	170/110	20 Days	—	"	L	3120	
5338	52	—	22	Cloud	Slight	180/115	49	—	"	L	930	
5339	36	—	38	Trace	"	150/100	—	—	"	L	2850	
5360	25	—	38	Cloud	Severe	180/100	7 Days	—	L.S.C.S.	L	2500	
5393	21	—	36	Trace	Slight	160/120	"	A.R.M.	L.F.	L	3170	

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks	Maturity On Ad. At Dismission	Albuminuria On Maximum Discharge	Oedema	Highest Blood Pressure	No. of Days in Hospital before labour or discharge	If labour induced in State	Method of Delivery	Result M. C.	Weight of Child gm.	Remarks
5399	34	—	36-3	Trace	Slight	160/90	—	—	Spont.	L	3000	
5407	30	—	41-6	None	Moderate	160/120	—	—	"	L	3200	
5432	31	—	39-3	Trace	Slight	160/100	—	—	"	L	3450	
5488	31	—	40-3	None	"	145/100	—	—	"	L	3350	
5540	30	—	37-6	Trace	"	150/90	—	—	"	L	3000	
5560	27	1	40-6	"	"	160/90	—	—	"	L	3175	D.A.A. Admitted at 39-1/7 wks. for investigation and Treatment
5599	38	—	41-2	"	Moderate	145/90	—	—	"	L	3250	
5621	24	—	35-4	Solid	Severe	155/115	3 Days	—	L.S.C.S.	L	2495	D.A.A.
5626	41	3	40-2	None	Slight	140/105	—	—	Spont.	L	2750	
5653	28	—	41-5	Trace	Moderate	180/110	7 Days	Stripping	"	L	3730	Twin Pregnancy
5650	28	—	33-3	None	Severe	165/105	7 Days	Stripping of Membranes	"	L	2400	
5692	38	1	38-4	"	Slight	160/100	3 "	"	"	L	2100	
5701	24	—	37-2	Trace	"	170/100	9 "	"	"	L	3317	D.A.A.
5771	18	—	37-5	"	Severe	150/100	6 "	"	"	L	3940	
5779	24	—	44-2	None	Slight	140/110	9 "	"	"	L	2780	
5856	26	—	39-1	Cloud	"	180/120	1 Day	—	L.S.C.S.	L	3280	
5913	28	—	36-4	None	"	140/100	2 Days	—	Spont.	L	2800	Twin Pregnancy
5993	24	—	38-6	Trace	"	140/100	—	—	"	L	1800	
5994	36	—	39-3	None	"	150/100	—	—	"	L	3280	
5996	37	—	39-2	"	"	150/90	—	—	"	L	3110	
6022	20	—	39-2	Cloud	Moderate	160/110	2 Days	Stripping/ of Membranes	"	L	3000	Trial Labour Successful—4 1/2 hrs.
6032	37	—	39-4	None	Slight	150/90	—	—	"	L	3050	
6052	21	—	35	Trace	Moderate	180/110	1 Day	Stripping of Membranes	"	L	2900	
6058	27	—	42	None	"	140/110	—	—	"	L	2750	
6080	32	2	40-3	"	"	160/100	1 Day	Stripping of Membranes	"	L	3330	
6108	34	—	38-5	None	Slight	150/90	8 Days	A.R.M.	Spont.	L	3370	Indication of Induction: Post-maturity
6109	24	—	42-5	"	Moderate	155/100	4 "	Stripping of Membranes	"	L	2500	Admitted at 32-6/7 wks. for investigation and Treatment
6117	32	—	34-6	"	Severe	140/100	—	—	"	L	1700	
6118	35	—	37-6	"	Slight	150/100	—	—	"	N.D.	2100	
6131	25	—	38	Trace	"	140/90	—	—	"	L	2560	
6133	24	—	35	Solid	Severe	180/120	—	—	L.S.C.S.	L	3150	
6161	30	—	38-1	Trace	Slight	150/110	—	—	"	L	1900	Twin Pregnancy
6197	32	—	40-2	"	"	150/100	7 Days	—	Spont.	L	1600	B.B.A.
6202	30	—	37	Cloud	"	165/120	—	—	L.S.C.S.	L	2200	
6216	24	—	37-6	Trace	Slight	140/100	—	Stripping of Membranes	Spont.	L	2680	
									"	L	2520	
									"	L	2610	

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks	Metamity On Adm. Delivery	Albuminuria On Discharge	Oedema	Highest Blood Pressure	No. of Days in hospital before labour or discharge	If labour induced State method	Method of Delivery	Result M. C.	Weight of Child gm.	Remarks					
6907	37	2	4	38-4	38-5	None	None	Slight	170/100	—	Stripping of Membranes	Spont.	L	L	3800		
6955	21	—	—	40-2	42-3	Trace	Trace	Moderate	160/110	15 Days	—	M.R. + M.F.	L	L	3000		
6963	24	—	1	34-2	37	None	None	Slight	150/105	19 "	—	Mid Forceps	L	N.D.	3370		
6975	26	—	—	39-1	39-1	None	None	Slight	140/100	—	—	Mid Forceps	L	L	2722		
{ 6980	31	—	3	39	39	Trace	Trace	Severe	140/100	—	—	Spont.	L	L	3420	Admitted at 24-5/7 wks. for Investigation and Treatment Elective Mid Forceps Delivery	
7024	23	—	—	39-3	39-5	—	—	—	140/105	2 Days	Stripping of Membranes	Mid Forceps	L	L	2778	Cardiac Disease D.A.A.A.	
7054	32	1	5	39-6	41-1	Trace	Trace	—	160/105	9 "	—	Spont.	L	L	2770		
7056	30	1	4	39-4	39-4	None	None	Moderate	140/105	5 Days	Stripping of Membranes	—	L	L	3420		
7087	26	—	—	35	35-5	—	—	—	145/95	—	—	—	L	L	3062		
7090	38	—	5	39-1	39-1	Cloud	Cloud	—	140/95	—	—	—	L	L	2920		
7116	28	—	—	?	?	Trace	Trace	Slight	185/130	—	—	—	L	L	5230		
7127	27	—	1	44-1	44-1	None	None	Moderate	140/100	—	—	—	L	L	2880		
7175	34	—	5	29-2	29-2	—	—	—	150/100	—	—	Spont.	L	L	2230	Twin Pregnancy	
7183	28	—	2	36-1	38	—	—	—	140/105	13 Days	—	(1) Spont.	L	L	2350	Twin Pregnancy	
7184	32	—	3	36-1	37-2	Trace	Trace	Moderate	150/105	8 Days	A.R.M.	(2) " "	L	L	2090	Twin Pregnancy	
7187	30	—	—	37	37-6	Trace	Trace	Slight	140/105	6 "	—	Spont.	L	L	2630		
7192	34	1	6	?	?	Trace	Trace	—	170/100	—	—	—	L	L	3495		
7299	41	—	7	40-4	40-4	Trace	Trace	—	150/100	—	—	—	L	L	3950		
7318	22	—	—	40-2	40-2	None	None	Moderate	140/100	—	—	—	L	L	3430		
7322	29	—	—	29-5	31-2	Cloud	Cloud	Severe	165/115	11 Days	—	C.C.S.	L	N.D.	3600	P.P.H. 1,000 c.c.	
7356	23	—	2	39-3	39-3	Trace	Trace	—	145/95	—	—	Spont.	L	L	1446	Twin Pregnancy	
7363	22	—	—	37-6	38	—	—	Slight	155/110	—	—	—	L	L	2700		
7372	21	—	—	38	38	None	None	—	140/100	—	—	—	L	L	2650		
7383	20	—	1	36	36	Cloud	Cloud	Moderate	145/100	—	—	—	L	L	2750		
7392	23	—	—	39-4	40-6	Trace	Trace	—	170/105	9 Days	—	(1) Spont.	L	L	2900	Twin Pregnancy. Prolapsed Cord	
7398	38	—	6	41-4	41-4	None	None	Slight	160/100	—	—	(2) " "	L	L	1900		
7422	26	—	—	39	39	Trace	Trace	—	165/95	—	—	Spont.	L	L	2250		
7466	42	—	7	38-6	39-2	—	—	Moderate	185/110	3 Days	—	Spont.	L	L	3380		
7490	33	—	—	38-3	38-2	—	—	—	140/100	—	—	—	L	L	2660		
7531	43	—	—	36-3	36-5	—	—	Slight	140/90	3 Days	—	L.S.C.S.	L	L	2860	P.P.H. 600 c.c.	
7555	45	—	6	36-4	37-1	—	—	—	140/100	4 "	—	Spont.	L	M.C.	3490	Pulmonary T.B.	
7556	20	—	—	37-2	37-3	Trace	Trace	—	170/100	1 Day	Stripping of Membranes	Mid Forceps	L	L	3480	Mid Pelvic Contraction	
7566	32	—	1	28-4	28-4	Cloud	Cloud	—	140/100	—	—	Spont.	L	L	2480		
7592	29	1	1	29	30-2	Trace	Trace	—	180/120	4 Days	—	L.S.C.S.	L	N.D.	1750	Concealed Placenta	
																1200	Accidental Haemorrhage

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks	Maturity On Admission	Albuminuria On Discharge	Oedema	Highest Blood Pressure	No. of Days in Hospital before labour or discharge	If labour induced State method	Method of Delivery	Result M. C.	Weight of Child gm.	Remarks
7615	26	1	36-4	—	Slight	180/100	—	—	Spont.	L.	3232	
7641	37	7	38-2	—	"	140/90	—	—	"	L.	3150	
7699	30	2	40-5	Trace	"	150/100	—	—	"	L.	2380	
7715	21	—	37-5	Cloud	"	155/105	—	—	"	L.	2500	
7723	24	—	37-4	Trace	"	155/100	8 Days	—	"	L.	2800	
7725	26	5	40-1	—	Moderate	155/96	—	—	"	L.	3130	
7740	26	—	33-4	—	Slight	160/100	—	—	"	L.	2420	
7747	23	—	40-6	—	"	160/90	—	—	"	L.	2920	
7796	25	—	40	—	"	140/100	—	—	"	L.	3060	
7863	26	—	35-3	Trace	Severe	145/95	2 Days	—	Low Forceps	L.	2710	P.P.H. 700 c.c. Pulmonary T.B.
7866	32	1	40-2	—	"	140/90	3	—	Spont.	L.	3330	
7888	32	1	39-1	—	"	145/100	2	—	"	L.	2778	
7928	38	6	34-2	Trace	Slight	150/100	—	—	"	L.	3300	
7931	24	2	39-4	"	"	149/95	—	—	Low Forceps	L.	3402	Manual Removal of Placenta Previous L.S.C.S. Done at another Hospital for Pre-eclampsia
6966	28	3	37-6	—	"	160/100	—	—	Spont.	L.	3600	Admitted at 34-6/7 wks. for Investigation and Treatment
7970	30	—	38	Trace	Moderate	160/90	—	—	"	L.	3020	
7992	19	—	39-3	—	Slight	169/100	2 Days	—	"	L.	2650	
8030	27	—	39-6	—	"	140/90	—	—	Spont.	L.	3120	
8104	25	—	39-4	Trace	"	150/100	—	—	"	L.	3050	
8122	33	1	39-1	—	"	140/95	1 Day	—	"	L.	3580	
8253	25	—	39-2	Trace	"	140/100	—	—	"	L.	3000	
8345	24	—	36	Trace	"	140/90	—	—	"	L.	2570	
8188	34	1	39	—	"	155/90	—	—	"	L.	4026	
NON-BOOKED												
190	43	3	37	Cloud	Moderate	200/115	22 Days	—	Spont.	L.	3555	Twin Pregnancy
792	35	1	37-3	Trace	Severe	165/95	22	—	"	L.	1900	
1011	45	7	39-2	—	Slight	190/105	—	—	"	N.P.	920	
1276	19	—	40-3	—	"	140/100	—	—	"	L.	3000	
1613	22	—	40	"	Moderate	140/100	1 Day	—	L.S.C.S.	L.	2948	
1425	22	—	41	Cloud	—	170/110	—	—	Spont.	L.	4400	
1953	34	—	42-6	Trace	Severe	170/110	—	Stripping of Membranes	"	L.	3629	
1958	29	2	35-3	Solid	Slight	190/120	5 Days	—	L.S.C.S.	L.	3680	Caesarean Section and Myomectomy
1998	30	2	39	Trace	—	170/110	7	—	Spont.	L.	3459	Subserous Uterine Fibroid Pulmonary T.B. Trial Labour Successful—1-15/60 Hrs.
2260	36	10	38-2	—	Moderate	160/110	—	—	"	L.	2900	

PRE-ECLAMPSIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks	Maturity On Ad- mission	Albuminuria On Discharge	Oedema	Highest Blood Pressure	No. of Days in hospital before labour or discharge	If labour induced State	Method of Delivery	Result M. C.	Weight of Child gm.	Remarks
2316	24	—	36-3	Trace	Severe	190/110	—	—	M.R. + M.F.	L	3062	
2837	32	2	40-1	"	Slight	140/95	—	—	(1) Vertex-Spont. (2) L.A.P.-In-ternal Version	L	2470 1950	Twin Pregnancy
2983	42	—	40	Solid	Severe	175/100	—	—	Spont.	L	2600	Hypertensive Heart Failure
3129	25	—	42	Trace	Moderate	140/100	5 Days	—	M.F.	L	3400	
3323	42	3	40-6	Cloud	Slight	155/110	—	—	Spont.	L	3400	
3575	38	—	35-6	Trace	Moderate	175/90	1 Day	—	"	L	2480	Twin Pregnancy
3605	37	—	40-4	"	Slight	210/140	—	—	"	L	2800	
3644	42	1	35-2	—	"	150/90	3 Days	—	Internal Version	L	2588	D.A.A.
3782	32	—	39-2	—	"	180/110	3	—	Spont.	L	3060	
3956	28	—	39	Trace	"	170/100	"	—	L.S.C.S.	L	2400	Placenta Praevia Type III
4135	26	—	33-2	Cloud	Moderate	150/115	11 Days	Stripping of Membranes	Spont.	L	3580	
4189	31	1	38-4	Trace	"	180/110	—	A.R.M.	"	L	2800	
4270	38	—	40-4	"	Slight	150/110	7 Days	—	"	L	3000	
4566	44	1	36-3	"	"	190/130	5	—	"	L	2280	
4618	29	—	39-5	Cloud	"	180/115	"	—	"	L	2775	
4997	43	—	36	Trace	"	150/90	—	—	"	L	2400	
5057	31	—	40	"	"	160/100	—	—	"	L	3900	
5100	23	—	33-6	"	Slight	160/100	—	—	"	L	3555	
6041	34	—	35-4	Trace	"	150/110	—	—	"	L	3260	Trial Labour Successful—5-50/g. Hrs.
6306	30	—	28-2	"	Moderate	195/125	3 Days	Stripping of Membranes	"	L	2950	
7483	38	—	35	"	Severe	195/115	—	—	C.C.S.	L	2490	
7526	50	1	39	"	"	150/100	4 Days	—	Spont.	L	3250	
7565	27	—	37	—	"	140/95	"	—	"	L	1750	
7835	52	—	40-5	—	Slight	140/105	"	—	"	L	2680	
8088	35	—	41	Trace	Severe	205/103	—	—	C.C.S.	L	2880	P.P.S.
8112	30	—	36-5	Cloud	"	200/120	1 Day	—	"	L	1850	

TABLE 5
HYPERTENSION—ESSENTIAL

There were 38 cases, an incidence of 0.45%, 36 Booked and 2 Non-booked.

There were no maternal deaths.

There were 2 stillbirths, 1 neo-natal death, and 1 missed abortion.

Reg. No.	Age	Previous Pregnancies Before 28 wks	Maturity On Ad. At De- mission every	Albuminuria At Maximum Discharge	Oedema	Highest Blood Pressure	No. of Days in hospital before Delivery	If labour induced State method	Method of Delivery	Result		Weight of Child gm.	Remarks
										M.	C.		
BOOKED													
86	42	—	6	34.5	—	185/115	—	—	Spont.	L	L	2800	
575	43	3	9	40.2	Trace	150/100	—	—	—	L	L	3020	Admitted at 28 3/7 wks. for investigation and treatment
662	32	1	2	31	—	150/100	1 Day	—	—	L	S.B.	1850	Admitted at 26-1/7 and 31-1/7 wks. for investigation and treatment
1301	37	—	2	41.2	—	150/100	1 "	—	—	L	L	3400	Admitted at 34-1/7 wks. for investigation and treatment
1743	43	—	6	36.4	Trace	170/105	3 "	—	—	L	L	3490	Admitted at 24-2/7 wks. for investigation and treatment
2248	21	—	—	38.4	—	150/100	—	—	Missed Abortion	L	L	2680	Admitted at 35-2/7 wks. for investigation and treatment
2523	34	2	1	25	Trace	210/120	—	—	Aborted	L	Mac.	450	D.A.A.
2778	22	—	—	39	—	180/110	—	—	Spont.	L	L	2900	Accidental Haemorrhage
3117	30	—	2	42.1	—	170/105	—	—	—	L	L	2680	
3334	42	1	6	38.4	—	155/105	1 Day	—	—	L	L	2690	
3664	27	—	2	39.2	—	145/95	—	—	—	L	L	3835	
4072	32	2	6	39.1	—	165/115	—	—	—	L	L	3050	
4333	35	1	6	40.6	Trace	160/110	—	—	—	L	L	3580	
4364	39	1	6	38.4	—	160/100	1 Day	—	—	L	L	3460	
4472	40	—	1	39.5	—	170/110	—	—	—	L	L	2840	
4478	36	—	1	39.4	—	170/90	—	—	L.S.C.S	L	L	3200	Indication for C. S. Contracture I Pelvis, Previous C.S.; Breech Presentation
4679	38	—	7	29.5	Cloud	230/145	7 Days	—	Spont.	L	S.B.	680	
4883	29	—	3	40.3	Trace	180/115	—	—	—	L	L	3040	Previous P.E. 1951, 1954
5124	46	—	8	41.1	—	180/130	—	—	—	L	L	2850	D.A.A.
5125	39	—	11	38.3	—	180/130	—	—	—	L	L	2500	
5362	31	—	—	?	—	175/105	—	O.B.F. & Stripping of Membranes	—	L	L	2800	

HYPERTENSION—ESSENTIAL (Continued)

Reg No	Age	Pregnancies Before 28 wks	Maturity On Ad mission	At De mission	At Maximum	Albuminuria	At Di change	Oedema	Highest Blood Pressure	No of Days in Hospital before Delivery	Method of Delivery	Weight of Child	Remarks
5796	40	—	7	39.3	39.3	Trace	—	—	160/115	—	Spont	2725	
6039	21	—	—	37.5	37.5	—	—	—	150/100	—	—	2,000	
6081	35	—	7	34.5	35.2	—	—	—	140/170	5 Days	—	7440	
6437	35	—	4	38.6	38.6	—	—	—	140/100	—	—	3,700	
6462	26	—	4	39.3	39.3	—	—	—	150/90	2 Days	—	2600	
6482	31	—	—	38.1	38.3	—	—	—	145/90	—	—	2,600	
6696	37	1	6	39.5	40.3	Trace	—	Slight	140/110	6	—	2,300	
6761	24	—	—	40.2	40.7	—	—	—	140/100	—	M1	3320	
6896	33	—	2	39.5	39.5	—	—	—	140/100	—	Spont	4120	
6950	37	3	5	40.6	40.6	—	—	Slight	150/90	—	—	2910	
7210	33	—	6	39.5	39.5	—	—	—	150/100	—	—	3150	
7580	40	—	2	42.2	42.2	Trace	—	—	140/90	—	—	3345	Indication for C.S. Contracted Pelvis and Previous C.S.
8111	29	—	2	38.6	38.6	—	—	—	155/100	—	Spont	3800	
8272	23	—	—	39	39	—	—	—	140/110	—	—	2430	
8287	35	1	3	—	?	—	—	—	150/90	—	—	2200	
NON BOOKED													
2312	39	1	4	26.1	Un delivered	—	—	—	145/115	10 Days	Undelivered	—	Indication for C.S. Severe Hypertension
7494	38	—	5	38	38.3	—	—	—	185/170	3	C.C.S.	3.45	

TABLE 6

CHRONIC NEPHRITIS

There was only one case, Booked No maternal or foetal death

Reg No	Age	Previous Pregnancies Before 28 wks	Maturity On Ad At De	Albuminuria On Maximum Discharge	Oedema	Highest Blood Pressure	No of Days in hospital before labour or discharge	If labour induced State method	Method of Delivery	Result		Weight of Child gms	Remarks			
										M	C					
280	23	1	1	41	41.2	Solid	Solid	Slight	130/95	One Day	---	Spont	L	L	3270	Caets—Hyaline

BOOKED

TABLE 7
ECLAMPSIA

There were 19 cases, an incidence of 0.23%, 15 Booked and 4 Non-booked.
There was one maternal mortality in the Non-booked cases.
There were 2 neo-natal deaths, 1 in Booked and 1 in Non-booked cases.
There were 2 sets of twins.

Reg. No.	Age	Pregnancies Booked After 28 wks. wks.	Condition on Admission	If in labour	Before Adm.	Fls. Ante Partum	Fls. Intra Partum	Fls. Post Partum	Total	Urine Albumin On admission or Maximum	Quantity in first 24 hours	Oedema	Highest Blood Pressure	No. of Days in hospital before Delivery	Treatment in hospital	Induced State method	Method of Delivery	Result M. C.	Remarks
BOOKED																			
181	22	—	42	Good	In Labour	—	1	—	1	Solid	None	++	180/120	—	Pentothal	Spont.	L	I	
881	23	—	45-5	"	"	—	1	—	1	Cloud	—	Slight	140/110	—	" 4.3 gm. " 6.87 gm.	Manual Rotation	L	N.D.	
1250	25	—	42-2	"	"	—	1	—	1	—	—	++	190/125	2 Days	" 7.53 gm.	Mfid Forceps L.S.C.S.	L	L	Right Dermoid Cyst of Ovary
1268	20	—	40-3	"	Not In Labour	—	1	—	1	Trace	—	Slight	160/100	—	Mg SO ₄ Pentothal	D-S	L	L	
1504	35	1	35	"	"	—	1	—	1	Solid	7-	+++	175/110	4 Days	5.4 gm. Cathister	"	L	L	
2318	27	1	38-6	Fair	In Labour	2	1	—	3	Trace	—	+++	100/60	—	" 5.2 gm. " 4.5 gm.	L.S.C.S.	L	L	
2322	27	—	39-3	"	"	—	1	—	1	Solid	10	+++	205/110	—	" 4.5 gm.	Low Forceps	L	L	
4051	21	—	36-2	—	"	—	1	—	1	Trace	—	Slight	175/110	—	" 6.25 gm.	Spont.	L	L	
5039	27	—	37-2	Good	"	—	1	—	1	None	—	++	170/110	—	" 4.2 gm.	"	L	L	
5065	20	—	38	"	"	—	1	—	1	"	—	+++	165/110	—	" 4.9 gm.	"	L	L	Left Heart Failure
5370	32	—	44-3	"	"	—	1	—	1	Solid	Trace	Slight	155/95	7 Days	" 2.8 gm.	(1) Mid Forceps (2) Internal Version	L	L	Uniovular Twin Pregnancy
6352	25	—	39	"	"	—	1	—	1	—	—	++	160/110	—	Mg SO ₄	Mfid Forceps	L	L	
6701	23	—	38-1	"	Not In Labour	10	—	—	10	Cloud	—	+++	155/110	—	"	Spont.	L	L	
7395	36	—	37	"	"	—	1	—	1	Trace	—	+++	190/110	5 Days	" Stripping Membrane	Mfid Forceps	L	L	
8072	20	—	43-6	"	In Labour	—	1	—	1	Severe	—	Slight	160/120	—	Pentothal 9.76 gm.	Low Forceps	L	N.D.	Contracted Mid Pelvis
NON-BOOKED																			
909	24	—	35-2	Good	In Labour	1	—	—	1	—	—	"	145/105	2 Days	Mg SO ₄	L.S.C.S.	L	L	Placenta Praevia Uniovular Twin Pregnancy
1933	32	—	37-5	"	"	2	1	—	3	Trace	—	"	250/160	—	Pentothal 5.3 gm.	Spont.	L	L	
5021	25	—	41-4	"	"	—	2	—	2	Solid	—	"	160/110	—	" 5.4 gm.	Low Forceps	L	L	
3085	27	—	1 28	Please Read Summary Of Maternal Deaths.															

TABLE 8

ACCIDENTAL ANTEPARTUM HAEMORRHAGE

(Includes cases of A.P.H. other than those proven to be due to Placenta Praevia).

There were 38 cases, 24 Booked and 14 Non-booked.

There were no maternal deaths.

There were 19 stillbirths, 11 Booked and 8 Non-booked; 3 neo-natal deaths, one Booked and 2 Non-booked.

Reg. No.	Age	Previous Pregnancies Before 28 wks.	At Ist. After haemor. stage	Maturity On Admission	At Delivery	Condition on Admission	If in Labour	Cause of A.P.H. if known	Treatment	Blood Transfusion c.c.	Amount of Bleeding c.c. Concoaled Mixed Retained	Result	Remarks
BOOKED													
363	30	—	34.4	34.4	34.4	Fair	In Labour	—	Spont. Bipolar Version	1000	(1800)	L	S.B.
407	39	—	30	30	30.1	Good	Not In Labour	—	Craniotomy	1500	(800)	L	S.B.
1504	35	1	34.3	34.3	35	Fair	"	Toxaemia	—	—	100	L	L
1587	28	—	33.5	36.3	36.3	Good	In Labour	—	Spont. A.R.M.	—	(100)	L	L
1663	39	1	33.3	33.3	33.4	"	"	—	A.R.M.	1000	—	L	Mac.
2221	22	—	36.1	36.1	36.1	Ill and Tired	"	—	L.F.	—	(100)	L	L
2512	29	—	44.1	44.1	44.1	Good	"	—	A.R.M. Spont.	—	900	L	S.B.
2541	42	2	36.4	36.4	36.4	"	"	—	A.R.M.	—	(650)	L	L
2713	42	—	37.5	37.5	37.5	Fair	Not In Labour	—	Stripping	1000	(800)	L	S.B.
2771	42	—	35.5	35.5	35.5	Poor	In Labour	—	A.R.M.	1500	(1600)	L	S.B.
3334	42	1	38.4	38.4	38.5	Good	In Labour	Essential Hypertension	Spont.	—	(250)	L	L
4531	27	—	36.2	36.2	36.2	Fair	"	—	"	500	(850)	L	S.B.
4632	27	2	36.1	36.1	36.1	Good	"	—	"	—	(350)	L	L
5912	32	5	38.6	38.6	38.6	"	"	—	"	—	200	L	L
5941	35	—	35.5	35.5	35.5	"	"	—	"	—	700	L	S.B.
6001	32	—	38.3	38.3	38.3	"	"	—	"	500	(850)	L	S.B.
6861	37	—	?	?	?	"	"	—	"	—	200	L	L
7235	28	—	36.6	36.6	36.6	"	"	—	C.C.S.	—	(200)	L	L
7334	33	—	41.2	41.2	41.4	Fair	Not In Labour	—	Spont.	—	(400)	L	L
7592	29	—	29	29	30.2	Fair	"	—	C.C.S.	—	—	L	N.D.

(10 x 6 cm)

_Blood (lot)

ACCIDENTAL ANTEPARTUM HAEMORRHAGE (Continued)

Reg. No.	Age	Previous Pregnancies Before After 28 wks 28 wks	At Lit. haemor- rhage	Maturity On Admission	At Delivery	Condition on Admission	If In labour	Cause of A.P.H. if known	Treatment	Blood Transfusion c.c.	Amount of Bleeding c.c. Concealed Mixed Revealed	Result M. C.	Remarks
7606	35	—	35-3	35-3	35-3	Good	In Labour	—	C.C.S.	1500	(1000)	L	Premature Placenta
7794	32	—	?	?	?	Good	"	—	Spont.	—	(50)	L	
8127	26	—	34-4	34-4	34-4	"	"	—	Spont.	—	200	L	
8325	36	—	33	33	33	"	"	—	Spont.	—	600	L	
NON-BOOKED													
1818	40	—	—	41	41	Poor	In Labour	Unknown	Willitt's Forceps	500	1200	L	S.B. P.P.S.
2669	36	1	?	?	?	Good	"	"	Spont.	—	400	L	
2710	25	—	29	30	30	Fair	"	"	A.R.M.	—	100	L	S.B.
2764	34	1	29	29	29	Good	Not In Labour	—	A.R.M.	500	700	L	S.B.
3067	46	—	?	?	?	Fair	"	"	A.R.M.	1000	(500)	L	S.B.
3323	42	3	40-6	40-6	40-6	Good	"	Toxaemia	Spont.	500	—	L	S.B. P.E.
3603	34	—	40-3	40-3	40-3	Poor	In Labour	—	Spont.	500	—	L	S.B.
4270	38	—	39-6	40-4	41-4	Good	Not In Labour	Toxaemia	Spont.	—	1400	L	S.B.
5904	28	—	39-5	39-6	39-6	"	In Labour	Premature Separation of Placenta	Spont.	—	(100)	L	L
6280	20	—	37-4	37-4	37-4	"	"	—	Spont.	500	(600)	L	L
6311	40	—	32-1	32-1	32-1	"	"	—	C.C.S.	500	(1400)	L	S.B. D.A.A.
7081	21	—	31	31	31	"	"	—	Spont.	—	(50)	L	N.D.
7989	41	1	34-5	34-5	34-5	"	Not In Labour	—	C.C.S.	—	200	L	N.D. Contractin Ring
8236	35	1	40	40	40	Fair	"	—	C.C.S.	1000	1000	L	S.B. P.P.S.

TABLE 9

PLACENTA PRAEVI

There were 48 cases, 33 Booked and 15 Non-booked.
 There were no maternal deaths.
 There were 8 stillbirths, 5 Booked and 3 Non-booked; 11 neo-natal deaths, 9 Booked and 2 Non-booked.
 There were 2 sets of twins.

Reg. No.	Age	Pregnancies Before 28 wks.	At Ist. haemorrhage	Maturity On Admission	At Delivery	Condition On Admission	If in labour	Type 1, 2, 3, 4,	Treatment	Blood Transfusion c.c.	Amount of Bleeding c.c.	Result M. C.	Birth Weight in gram.	Remarks
6949/55	33	—	1 36.0	36.0	38.3	Good	Not In Labour	III	L.S.C.S.	—	470	L L	3370	
506/56	23	—	1 29.3	29.3	30.0	"	"	III	L.S.C.S.	500	900	L N.D.	1250	
624	38	1	4 34.0	34.0	40.1	"	"	I	Spont	—	350	L L	3175	
674	29	—	1 28.3	28.3	28.3	"	In Labour	I	"	—	50	L N.D.	1000	
1186	30	—	3 39.0	39.0	39.0	"	"	I	"	—	100	L L	3000	
1190	33	—	5 37.1	37.1	37.1	"	"	I	"	—	150	L L	2920	
1538	30	—	4 —	39.3	39.3	"	"	I	A.R.M.	—	50	L L	2750	
1649	31	—	4 30.5	36.4	36.5	"	"	II	Rigolac Vergole	—	500	L L	2565	
1918	41	—	4 32.2	34.4	34.4	"	"	III	L.S.C.S.	500	500	L L	2460	
2499	27	—	1 35.1	35.1	37.6	"	Not In Labour	III	A.R.M. & Willett's	—	450	L S.B.	3175	
2507	37	—	2 24.0	33.0	33.4	"	"	IV	Forceps	500	400	L L	2400	
2657	27	—	2 33.5	33.5	33.5	"	In Labour	III	L.S.C.S.	500	1000	L N.D.	2500	
2705	31	—	6 38.5	38.5	42.2	"	Not In Labour	I	Spont.	—	50	L L	3580	
2871	34	—	7 40.2	40.3	40.3	"	In Labour	I	"	—	150	L L	2650	
3086	28	—	4 36.1	36.2	36.4	"	Not In Labour	IV	L.S.C.S.	1500	800	L S.B.	2722	
3532	30	—	2 25.1	28.5	28.6	"	"	III	L.S.C.S.	—	200	L N.D.	1420	
4191	39	1	14 24.4	26.4	26.4	"	In Labour	III	J.S.C.S.	500	1000	L S.B.	960	
4684	26	1	3 34.1	34.1	36.0	"	Not In Labour	I	A.R.M. & Willett's	500	700	L S.B.	3178	
4737	27	1	3 39.4	39.5	40.1	"	"	IV	L.S.C.S.	500	700	L L	2665	Cystic kidney and liver. Hydrocephalus

PLACENTA PRAEVIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	At Ist. haemorrhage	Maturity On Admission	At Delivery	Condition On Admission	If in Labour	Type 1, 2, 3, 4,	Treatment	Blood Transfusion c.c.	Amount of Bleeding c.c.	Result M. G.	Birth Weight in grams	Remarks
5740	25	—	1 24.0	29.0	29.2	Good	In Labour	II	Spont.	—	270	L N.D.	1180	Prolapsed Cord
5663	40	—	7 31.0	?	?	Fair	Not In Labour	II	C.C.S.	1000	800	L N.D.	2570	
5670	23	—	1 33.5	33.5	33.5	Good	In Labour	II	L.S.C.S.	500	1000	L N.D.	2150	Twin Pregnancy
5932	27	—	4 38.2	38.2	38.2	Good	"	I	Spont.	—	—	L	2300	
6402	32	—	4 32.3	33.3	33.3	—	Not In Labour	II	L.S.C.S.	500	400	L N.D.	2090	
6435	25	—	2 35.6	35.6	35.6	Good	"	IV	L.S.C.S.	750	650	L N.D.	1160	
6611	21	—	3 35.3	35.3	35.4	"	"	I	C.C.S.	—	30	L	2707	
6691	30	—	3 34.1	34.1	34.2	"	In Labour	IV	Spont.	—	230	L	2200	
7137	34	—	8 40.4	40.4	40.0	"	"	IV	C.C.S.	—	600	L	2693	
7657	33	—	4 36.0	36.0	36.0	"	"	III	L.S.C.S.	1500	200	L	1885	
7873	26	1	6 29.4	29.4	29.4	"	"	III	Internal Version & Breech Extraction	1500	800	L S.B.	1400	
7964	27	1	3 39.2	?	?	Fair	Not In Labour	III	L.S.C.S.	500	1200	L	2870	
8279	29	1	2 39.2	39.2	39.2	Good	In Labour	I	Spont.	—	50	L	3175	
8284	26	—	1 40.0	40.0	40.2	"	Not In Labour	I	"	—	200	L	3250	
NON-BOOKED														
909	24	—	— 35.0	35.0	35.2	Good	In Labour	I	L.S.C.S.	—	—	L	1630	Antepartum Eclampsia
1191	35	—	5 42.3	43.2	43.5	"	Not In Labour	I	Spont.	—	300	L	1780	Twin Pregnancy
1558	35	—	2 39.4	39.4	39.6	"	In Labour	III	L.S.C.S.	1000	500	L	3260	Uterine Inertia
1706	25	—	2 39.4	?	?	Fair	"	III	Bipolar Version	500	700	L S.B.	1474	
1837	38	—	2 34.3	34.3	34.3	Poor	"	III	L.S.C.S.	1000	1500	L	2080	
2291	38	—	5 30.3	31.2	31.2	Good	"	I	Spont.	—	300	L	1850	
2566	30	—	3 28.6	29.4	32.4	"	Not In Labour	IV	L.S.C.S.	1000	300	L N.D.	1720	
2610	28	—	— 31.6	31.6	32.4	"	In Labour	II	L.S.C.S.	500	500	L	1840	
2634	32	—	1 36.2	36.2	36.2	"	In Labour	III	L.S.C.S.	500	1500	L	3289	
2743	33	—	5 29.4	29.5	32.2	"	Not In Labour	I	Spont.	—	300	L	1670	
2842	30	—	4 31.1	31.4	32.0	"	In Labour	II	"	—	150	L S.B.	1720	Prolapsed Cord
3644	42	1	4 35.2	35.2	35.5	"	Not In Labour	I	"	—	400	L	3060	
3782	32	—	2 37.2	39.2	39.5	"	In Labour	III	L.S.C.S.	—	700	L S.B.	2400	P.E.
4137	24	—	2 29.4	34.4	35.3	"	"	II	Willet's Forceps	500	540	L	2665	
7818	30	—	4 42.3	42.3	42.3	"	"	IV	C.C.S.	—	500	L	3910	

TABLE 10

HYDRAMNIOS

There were 20 cases, 16 Booked and 4 Non-booked.

There were no maternal deaths.

There were 9 stillbirths, 7 Booked and 2 Non-booked; 4 neo-natal deaths, 3 Booked and one Non-booked.

There was one set of twins, hydramnios being with the second twin.

Reg. No.	Age	Pregnancies		Maturity At		Treatment	Result		Remarks
		Before 28 wks.	After 28 wks.	On Admission	Delivery		M.	C.	
662	32	1	2	31	31-1	None	L	S.B.	Pre-Eclampsia—2250 c.c. Hydramnios
897	36	—	4	40-4	40-4	"	L	L	2000 c.c. Hydramnios
1260	28	—	3	42-6	43-1	"	L	S.B.	2460 c.c. Hydramnios
1575	26	—	—	39	42-5	"	L	L	2nd Twin of Uniovular Pregnancy: 2000 c.c. Hydramnios
1752	30	—	4	39	39	None	L	L	2500 c.c. Hydramnios
2534	35	2	7	34-4	34-4	"	L	S.B.	6000 c.c. Hydramnios; Anencephaly
3835	29	2	2	41-5	41-5	"	L	N.D.	2400 c.c. Hydramnios
3926	42	—	1	40-2	40-5	M.R. + M.F.	L	L	1500 c.c. Hydramnios
5599	26	—	2	32-5	32-5	L.S.C.S.	L	N.D.	2000 c.c. Hydramnios
5800	34	—	2	32	32	None	L	S.B.	2000 c.c. Hydramnios
6781	21	—	1	36-3	39-4	"	L	L	3000 c.c. Hydramnios
6869	41	2	4	39-4	40	L.S.C.S.	L	L	1950 c.c. Hydramnios. P.E. P.P.H.—700 c.c.
6863	24	—	1	34-2	37	None	L	N.D.	3000 c.c. Hydramnios; Anencephaly, Abortion
7089	40	—	6	38-5	?	Mid Forceps	L	Mac	2700 c.c. Hydramnios; Anencephaly
7338	40	—	6	38-5	38-5	None	L	S.B.	5000 c.c. Hydramnios; Anencephaly
7838	25	—	1	33	33	"	L	Mac	3000 c.c. Hydramnios
NON-BOOKED									
102	28	1	3	27-2	27-3	None	L	N.D.	6600 c.c. Hydramnios
2710	25	—	—	30	30	"	L	S.B.	(1) 1500 c.c. Hydramnios (2) Accidental Hemorrhage (Revealed Type)
3390	27	—	—	44	44	Mid Forceps	L	L	3000 c.c. Hydramnios
3599	29	—	1	42-6	42-6	L.S.C.S.	L	S.B.	(1) 10920 c.c. Hydramnios (2) Terata Anadidyma Monster

TABLE 11

TUMOURS ASSOCIATED WITH PREGNANCY, LABOUR OR PUERPERIUM

There were 12 cases, 10 Booked and 2 Non-booked.
 There were no maternal deaths.
 There were 10 uterine fibroids and 2 ovarian cysts.
 There was one stillbirth in the Non-booked cases.

Reg. No.	Age	Previous Pregnancies Before 28 wks.	Previous Pregnancies After 28 wks.	Nature of Tumour	Treatment	Result M. C.	Remarks
BOOKED							
273	31	5	2	Cervical Fibroid	Nil	L	
295	36	---	---	Uterine Fibroid	Myomectomy at time of L.S.C.S.	L	
334	32	---	---	"	L.S.C.S.	L	
834	27	---	2	"	Nil	L	
1017	30	---	4	"	Nil	L	
1230	25	---	---	Dermoid Cyst of Ovary, Rt.	Rt. Oophorectomy at time of L.S.C.S.	L	Antepartum Eclampsia
1470	24	---	2	Uterine Fibroid	Nil	L	
2516	29	1	3	Dermoid Cyst of Ovary, Left	Left Oophorectomy	L	Oophorectomy done on the 6th day of puerperium
7255	30	---	1	Uterine Fibroid	Nil	L	
7361	29	---	1	"	Nil	L	
NON-BOOKED							
1958	29	---	2	Uterine Fibroid, Subserous	Myomectomy at time of L.S.C.S.	L	
6649	36	---	2	Cervical Fibroid	Extraperitoneal C.S.	L S.B.	Prolapse of Cord

TABLE 12
PROLONGED LABOUR

(Labour lasting forty-eight hours or more).

There were 43 cases, 39 Booked and 4 Non-booked; 34 in primiparæ and 9 in multiparæ.

There were no maternal deaths.

There were 4 stillbirths, and 3 neo-natal deaths.

There were 2 sets of twins.

Reg. No.	Age	Maturity	Pregnancies Before 28 wks.	Position of Fœtus At Labour	Other Obstetric Abnormality	Cause of Delay	Time of Rupture of Membranes i.e. hours before Delivery	Duration of Labour	1st Stage	2nd Stage	3rd Stage	Treatment	P.P.H. c.c.	Morbidity	Method of Delivery	Weight of Child gm.	Result M. C.
BOOKED																	
277	25	40-5	—	L.O.P.	L.O.P.	Uterine Inertia	21:00	61:00	—	—	—	Sedation	—	—	L.S.C.S.	3147	L
938	28	40-5	—	L.O.T.	L.O.A.	"	4:13	65:50	2:05	0:05	—	"	—	3	M.F.	4430	L
1029	24	42-1	—	L.O.T.	L.O.A.	"	5:25	58:25	2:30	0:05	—	"	—	—	M.F.	4000	L
1089	32	40-3	—	L.O.P.	L.O.P.	O.P.	?	4 days	—	—	—	"	—	—	L.S.C.S.	2800	L
1253	21	41	—	L.S.A.	L.S.A.	Uterine Inertia	0:55	64:15	0:25	0:05	—	"	—	—	Assisted Breech	3580	L
1541	25	40-2	—	L.O.P.	L.O.P.	"	Leaking	53:00	—	—	—	"	—	7	L.S.C.S.	3740	L
1677	28	39	—	L.O.T.	L.O.A.	"	2:45	68:25	3:05	0:02	—	"	—	—	M.R. + M.F.	3800	L
1769	27	42-1	—	R.O.P.	R.O.P.	"	3:25	115:00	0:20	0:10	—	Sedation, Pitocin drip	—	—	L.F.	3175	L
												5 units in glucose	—	—	L.S.C.S.	2200	L
1844	37	38-6	—	L.O.P.	L.O.P.	"	15:00	62:00	0:40	0:03	—	Sedation	—	—	M.F.	2950	L
2016	32	40-4	—	L.O.A.	L.O.A.	"	11:30	65:35	0:45	0:02	—	Sedation, Pitocin drip	—	—	Spont.	2900	L
2199	21	40-6	—	L.O.P.	L.O.A.	"	1:10	66:05	—	—	—	"	—	—	M.F.	3780	L
2268	30	40-1	—	L.O.T.	L.O.A.	"	2:15	60:50	1:35	0:05	—	"	—	—	M.F.	2480	L
2452	34	34-6	1	R.O.P.	R.O.A.	"	0:27	53:30	0:50	0:45	—	"	—	—	Spont.	2730	L
												"	—	—	"	3140	L
2749	39	34-1	—	L.O.A.	L.O.A.	"	0:50	69:15	0:50	0:10	—	"	—	—	"	3140	L
3049	34	41	2	R.O.P.	R.O.A.	"	30:00	73:10	2:00	1:20	—	Sedation, i.v. glucose and saline	600	1	M.R. + M.F.	3550	L

PROLONGED LABOUR (Continued)

Reg. No.	Age	Maturity	Pregnancies Before 28 wks.	Position of Foetus At Labour	Other Obstetric Abnormality	Cause of Delay	Time of rupture of membranes i.e. hrs. before delivery	1st Stage	2nd Stage	3rd Stage	Treatment	P.P.H. Morbidity	Method of Delivery	Weight of Child gm.	Sex of Child
3089	20	35-6	1	R.O.A.	—	Uterine Inertia	0:20	61:40	0:10	0:05	Sedation	—	Spont.	2470	L
3102	26	41.4	—	L.O.A.	—	Inertia	2:05	56:50	2:05	0:25	"	—	L.S.C.S.	3000	L
3121	20	40-6	—	L.O.P.	—	O.P.	66:45	66:45	1:05	0:05	"	—	M.F.	3147	L
3139	25	40-2	—	L.O.A.	P.E.	Uterine Inertia	6:05	73:00	0:35	0:25	"	—	Spont.	3050	L N.D.
3417	19	39-2	—	R.C.A.	—	"	7:00	49:00	3:07	0:03	"	—	L.F.	3190	L
3446	25	38-2	—	L.O.A.	—	"	8:30	63:20	2:15	0:30	"	—	Spont.	3030	L
3512	26	39-4	—	L.S.A.	—	"	8:55	65:15	0:20	0:05	"	—	"	3572	L
3824	30	40-1	2	L.O.T.	—	"	1:30	49:45	0:15	0:10	"	—	L.F.	3629	L
3833	43	39-5	4	R.O.P.	—	"	Leaking	61:45	1:15	0:10	"	—	L.F.	3570	L
4179	28	40-3	—	L.O.A.	—	"	0:35	53:30	0:30	0:05	"	—	L.F.	3062	L
4568	24	42-3	—	L.O.A.	—	"	0:40	49:15	1:25	0:05	"	—	Spont.	3170	L
4892	34	42-4	—	L.O.A.	—	"	Leaking	51:55	0:45	0:05	"	—	"	2850	L S.B.
5048	25	39-5	—	L.O.A.	—	"	0:40	57:25	1:50	0:05	"	—	"	2155	L
5180	20	48-3	—	L.O.A.	—	"	1:40	94:30	0:15	0:15	Sedation, i.v. glucose	800	Forceps Internal Version	3020	L
5570	32	44-3	—	L.O.P.	Twins, A.P.	"	0:30	94:30	0:25	0:03	"	—	"	2350	L
6561	23	38-6	—	R.O.A.	Eclampsia	"	0:30	72:15	0:15	0:03	Sedation	—	Spont.	2950	L
6575	28	39-6	1	L.S.P.	—	"	5:55	78:45	0:55	0:10	"	—	"	2750	L S.B.
6599	22	42	—	L.O.A.	—	"	0:30	52:30	1:30	0:10	"	—	L.F.	3374	L
6670	23	38-1	—	L.O.A.	—	"	1:35	62:45	0:30	0:10	"	—	Spont.	3500	L
6869	41	40	2	L.O.A.	—	"	1:50	77:10	0:50	0:05	"	—	"	2020	L
6930	29	40-5	—	L.O.A.	—	"	10:55	71:40	0:40	0:03	"	—	"	3330	L
7344	34	39-3	1	L.O.A.	—	"	47:00	4 days	—	—	"	Secondary 6 P.P.H.	L.S.C.S.	2400	L
7371	40	40-6	4	L.O.A.	—	"	11:00	80:30	—	—	"	1500 c.c.	L.S.C.S.	3070	L
8211	35	39-4	—	L.S.A.	C.P.	C.P.	8:25	68:00	—	—	"	—	L.S.C.S.	2750	L
NON-BOOKED															
1140	28	39	—	L.S.A.	—	Uterine Inertia	1:35	50:00	0:45	0:05	Sedation	—	L.F.	3000	L N.D.
1425	22	41	—	L.O.P.	P.E.	"	6:05	49:20	0:55	0:10	"	—	Spont.	3629	L
3704	23	35-3	—	R.O.P.	C.P.	"	13:00	51:30	2:15	0:10	"	—	M.F.	2400	L N.D.
4117	33	41	1	L.O.A.	Cervical Polyp	"	0:25	49:30	0:25	0:05	"	—	Spont.	4100	L

TABLE 13

PERSISTENT POSTERIOR POSITION OF THE OCCIPUT AND TRANSVERSE ARREST

There were 139 cases, 131 Booked and 8 Non-booked.
 There were no maternal deaths.
 There were 6 stillbirths and 10 neo-natal deaths.
 Twins pregnancy occurred 9 times, 8 in Booked and one in Non-booked cases.

Reg. No.	Age	Prenatal Pregnancies Before 28 wks.	Maturity	Position of Occiput	Type of Pelvis if Known	Treatment	Result	Weight of child gm.	Remarks
6787/55	24	—	44.1	R.O.T.	Gynaecoid	M.R. + M.F.	L	4000	P.E.
24/56	26	—	40.5	R.O.P.	Normal	M.R. + M.F.	L	3650	
96	23	—	38.4	L.O.P.	"	Spont. Face to Pubes	L	2820	
203	32	—	40.3	R.O.P.	"	"	L	3402	
224	25	—	40	L.O.P.	Gynaecoid	M.R. + M.F.	L	3500	
397	34	—	?	L.O.P.	Anthropoid	Spont. Face to Pubes	L	2900	
439	25	—	40.2	L.O.P.	Gynaecoid	Spont.	L	3140	
404	27	—	41.3	L.O.P.	Normal	Spont. Face to Pubes	L	3200	
597	26	—	38.3	L.O.P.	"	M.R. + M.F.	L	2970	
652	30	—	40.5	R.O.P.	"	M.R. + M.F.	L	3850	
700	26	—	36	L.O.P.	Gynaecoid	M.R. + M.F.	L	3232	
825	28	—	40	R.O.T.	Normal	M.R. + M.F.	L	2985	
881	23	—	45.5	R.O.T.	Gynaecoid	M.R. + M.F.	L	2960	
938	28	—	40.5	L.O.T.	Normal	M.R. + M.F.	L	4430	
995	22	—	43	R.O.T.	"	M.R. + M.F.	L	3162	
1001	23	—	37.6	R.O.P.	"	M.R. + M.F.	L	2850	Baby referred to Q.M.H.
1029	24	—	42.1	L.O.T.	Gynaecoid	M.R. + M.F.	L	4000	
1045	28	—	39.3	R.O.T.	Normal	M.R. + M.F.	L	2665	
1061	30	—	39.5	R.O.P.	Gynaecoid	M.R. + M.F.	L	3340	
1082	34	—	40.2	L.O.T.	Normal	M.R. + M.F.	L	3864	
1093	36	—	35.3	R.O.T.	Normal	M.R. + M.F.	L	2600	
1215	24	—	42.1	L.O.P.	Gynaecoid	Spont. Face to Pubes	L	3110	
1248	25	—	39.6	R.O.P.	"	M.R. + M.F.	L	3200	
1324	27	—	37.5	R.O.T.	Gynaecoid	M.R. + M.F.	L	2820	
1339	25	—	?	R.O.T.	—	M.R. + M.F.	L	2650	D.A.A.
1491	23	—	37.3	L.O.A.	—	Spont. Face to Pubes Vertex Spont.	L	2780	Twin Pregnancy

BOOKED

Mild Contracted Pelvis. P.E.
 Perineal Wound Infection. P.E.
 Intrapartum Eclampsia—One Fit
 P.E.
 Baby referred to Q.M.H.

P.P.H. Laceration of Cervix—repaired

D.A.A.
 Twin Pregnancy

PERSISTENT POSTERIOR POSITION OF THE OCCIPUT AND TRANSVERSE ARREST (Continued)

Reg. No.	Age	Previous Pregnancies		Position of Occiput	Type of Pelvis if Known	Treatment	Result		Weight of child gm.	Remarks
		Before 28 wks.	After 28 wks.				M.	C.		
1677	28	—	—	R.O.T.	—	M.R. + M.F.	L	L	3800	
1716	40	1	9	L.O.T.	Gynaecoid	M.R. + M.F.	L	L	3420	
1769	27	—	—	R.O.P.	Platycephoid	Low Forceps	L	S.B.	3175	
1807	25	—	—	R.O.T.	Gynaecoid	Spont. Face to Pubes	L	L	3289	Successful Trial Labour 29-10/60 hrs.
1873	27	—	—	L.O.T.	Platycephoid	M.R. + M.F.	L	L	3005	
1954	40	2	5	R.O.P.	Gynaecoid	M.R. + M.F.	L	L	3050	
2002	38	3	6	R.O.T.	Anthropoid	M.R. + M.F.	L	L	3520	Right T.B. Hip
2068	27	—	—	L.O.P.	—	Low Forceps	L	L	2590	
2075	27	—	—	R.O.T.	Anthropoid	M.R. + M.F.	L	N.D.	3090	
2088	36	—	2	R.O.P.	Android	Mid Forceps	L	L	3200	
2092	21	—	—	L.O.T.	Gynaecoid	M.R. + M.F.	L	L	3200	P.E.
2123	26	—	2	L.O.P.	—	Spont. Face to Pubes	L	L	2800	
2144	21	—	—	L.O.P.	Gynaecoid	M.R. + M.F.	L	L	3300	
2198	25	—	—	R.O.P.	Anthropoid	Mid Forceps	L	L	2700	
2202	24	—	—	L.O.P.	Anthropoid	M.R. + M.F.	L	L	3200	
2304	36	—	5	R.O.P.	—	Spont. Face to Pubes	L	L	3670	
2352	31	—	4	L.O.P.	—	Spont. Face to Pubes	L	L	3620	
2342	36	—	7	L.O.A.	—	Spont.	L	N.D.	3300	Twin Pregnancy
2599	35	—	8	R.O.P.	—	M.R. + M.F.	L	N.D.	3050	
2671	33	—	3	L.O.T.	Gynaecoid	Spont. Face to Pubes	L	L	3410	
2771	42	—	6	L.O.P.	Anthropoid	M.R. + M.F.	L	L	3450	
				R.S.A.	—	Spont. Face to Pubes	L	S.B.	1899	Twin Pregnancy, Accidental Haemorrhage.
						Spont. (A.R.M.)	L	S.B.	1701	Blood Loss 1000 c.c. Blood Transfusion 1500 c.c.
2946	29	2	4	L.O.T.	Gynaecoid	M.R. + M.F.	L	L	3230	
3021	28	2	4	L.O.T.	—	M.R. + M.F.	L	L	3500	
3049	34	2	5	R.O.T.	—	M.R. + M.F.	L	L	3530	P.P.H. Manual Removal of Placenta
3116	21	—	38	R.O.P.	—	M.R. + M.F.	L	L	2340	
3155	28	—	36-5	R.O.P.	—	M.R. + M.F.	L	L	2466	P.E.
3180	25	—	3	L.O.P.	—	M.R. + M.F.	L	L	3890	
3263	31	1	1	R.O.P.	—	Spont. Face to Pubes	L	L	3800	
3404	24	—	2	R.O.P.	—	Spont. Face to Pubes	L	L	2120	Twin Pregnancy
				L.M.A.	—	Spont.	L	S.B.	2130	
3463	27	—	43	R.O.P.	Anthropoid	M.R. + M.F.	L	L	2880	
3583	24	—	37-4	L.O.P.	—	Spont. Face to Pubes	L	L	2680	
3656	25	—	39-1	L.O.T.	Gynaecoid	M.R. + M.F.	L	L	2960	
3733	23	—	38-4	R.O.P.	—	Spont. Face to Pubes	L	L	2700	
3799	31	—	36	L.O.A.	Gynaecoid	Vertex M.R. + M.F.	L	L	2300	Twin Pregnancy
				R.O.P.	—	P.O.P. M.P. + M.F.	L	L	2300	

PERSISTENT POSTERIOR POSITION OF THE OCCIPUT AND TRANSVERSE ARREST (Continued)

Reg. No.	Age	Pregnancies Before 28 wks., After 28 wks.	Maturity	Position of Occiput	Type of Pelvis if Known	Treatment	Result		Weight of child gm.	Remarks
							M.	C.		
6511	27	—	38-2	L.O.P.	—	Spont. Face to Pubes	L	L	2960	
6680	27	—	41-2	L.O.P.	Gynaecoid	M.R. + M.F.	L	L	2760	
6761	24	—	40-2	R.O.P.	"	Mid Forceps	L	L	3320	
6764	37	—	39-5	R.O.P.	Platyelloid	M.R. + M.F.	L	L	3150	P.E.
6844	32	—	41-1	R.O.P.	Anthropoid	Mid Forceps	L	L	29-48	Contracted Pelvis
6847	40	—	39-3	R.O.P.	—	Spont. Face to Pubes	L	L	3250	
6870	40	—	37-1	L.O.P.	—	M.R. + M.F.	L	L	2700	P.E.
6955	21	—	42-3	R.O.T.	Gynaecoid	M.R. + M.F.	L	L	3000	
6957	26	—	—	R.O.P.	Gynaecoid	Mid Forceps	L	L	2200	
6963	24	—	37	R.O.P.	Anthropoid	Face to Pubes	L	N.D.	3370	P.E. P.P.H. Hydramnios
6970	25	—	40-1	R.O.P.	"	Mid Forceps	L	L	3200	
7011	26	—	39-2	R.O.P.	Gynaecoid	M.R. + M.F.	L	L	3232	Manual Removal of Placenta
7209	23	—	37-1	R.O.P.	—	Spont. Face to Pubes	L	L	2670	Mitral Stenosis—Class II
7320	23	—	39-2	L.O.P.	Anthropoid	Low Forceps	L	L	3800	
7394	26	—	41	R.O.P.	—	Spont. Face to Pubes	L	L	3620	
7636	28	—	38	R.O.P.	—	M.R. + M.F.	L	L	2400	
7720	39	—	39-4	R.O.P.	Gynaecoid	M.R. + M.F.	L	L	3346	Elderly Primigravida. Successful Trial Labour 11-45/60 hours
7743	22	—	41-6	R.O.P.	"	M.R. + M.F.	L	L	3289	Pre-labour L.S.C.S. Successful Trial Labour 4-55/60 hours
7750	40	—	38	L.O.P.	—	Spont. Face to Pubes	L	L	2950	
7767	32	—	38-5	R.O.P.	—	"	L	L	3200	
7811	42	2	39-5	R.O.P.	Gynaecoid	"	L	L	4600	
7888	32	—	39-1	R.O.P.	"	L.S.C.S.	L	L	2778	P.E.
7924	23	—	39	L.O.P.	"	Spont. Face to Pubes	L	L	2750	
7945	31	—	34-6	R.O.T.	Gynaecoid	M.R. + L.F.	L	L	2540	
7971	38	—	43	L.O.T.	"	M.R. + M.F.	L	L	4050	
8008	25	—	40-3	R.O.P.	Platyelloid	Mid Forceps	L	L	2900	Previous C.S. Successful Trial Labour 3-40/60 hrs.
8017	26	—	40-1	R.O.P.	Gynaecoid	M.R. + M.F.	L	L	3657	
8051	42	—	34-1	R.O.P.	"	M.R. + M.F.	L	L	3320	Contracted Pelvis
8087	32	—	38-4	R.O.T.	"	M.R. + M.F.	L	L	3005	P.P.H. Cervical Tear—repaired
NON-BOOKED										
733	22	—	40	R.O.P.	Gynaecoid	M.R. + M.F.	L	L	3260	Prolonged Labour
792	35	—	30-4	R.O.P. → O.A.	Normal	Spont. Face to Pubes	L	L	1900	Term Pregnancy
913	36	—	42-5	R.O.P.	Gynaecoid	"	L	N.D.	920	P.E. Hookworm Infestation
1919	27	—	39-6	L.O.T.	"	M.R. + M.F.	L	L	2550	
2316	24	—	36-3	L.O.T.	"	M.R. + M.F.	L	L	2948	
3110	25	—	39-5	R.O.P.	—	M.R. + M.F.	L	N.D.	3062	P.E.
5021	25	—	41-4	R.O.P.	—	Low Forceps	L	L	2850	Antepartum Felampsia
5022	39	—	38	R.O.P.	—	M.R. + M.F.	L	L	2948	
									3470	

TABLE 14

UNCOMPLICATED BREECH DELIVERIES
PRIMIGRAVID BREECH PRESENTATION AND DELIVERIES
(WITHOUT COMPLICATIONS)

There were 141 cases, an incidence of 1.87%, 129 Booked and 12 Non-booked.

There were 58 primigravidae, and 83 multiparae.

There were 7 stillbirths, 3 macerated foetuses, all in Booked cases; and 9 neo-natal deaths, 5 in Booked and 4 in Non-booked cases.

Reg. No.	Age	Maturity	Obstetric Complication	Treatment	Result	Weight of Child gm.	Episiotomy or Tear	Remarks
BOOKED								
448	38	38-3	None	Assisted	L	2980	Episiotomy	
579	19	41-4	"	Forceps for A.C.H.	L	3120	"	
598	24	38-0	"	"	L	2600	"	
738	28	?	"	"	L	3000	"	
849	27	37-6	"	Assisted	L	2900	"	
1049	25	39-1	"	Spont.	L	3000	"	
1241	20	39-6	"	Assisted	L	3455	"	
1253	21	41-0	"	Spont.	L	3580	"	
1680	28	39-3	"	Assisted	L	2892	"	
1944	31	38-6	"	"	L	3350	"	
2203	25	38-3	"	"	L	2608	"	
2982	24	39-2	"	Spont.	L	3710	"	
3136	19	40-3	"	"	L	3550	"	
3250	24	37-5	"	"	L	2778	"	
3341	26	36-1	"	"	L	2700	"	
3452	28	37-6	"	"	L	2850	"	
3512	26	39-4	"	"	L	3502	"	
3562	21	40-2	"	"	L	3150	"	
3796	29	36-6	"	"	L	2560	"	
4029	28	36-0	"	"	L	3120	"	
4049	19	38-5	"	"	L	3040	"	
4670	20	38-1	"	"	L	1730	"	
4934	25	?	"	"	N.D.	3120	"	
5037	26	40-3	"	"	L	2695	"	

UNCOMPLICATED BREECH DELIVERIES (Continued)

Reg. No.	Age	Maturity	Obstetric Complication	Treatment	Result	Weight of Child gm.	Episiotomy or Tear	Remarks
5130	26	40-4	None	Spont.	L	2815	Episiotomy	
5185	24	38-1	"	"	L	2970	"	
5192	26	39-2	"	"	L	3450	"	
5315	27	41-6	"	Forceps for A.C.H.	L	4060	"	
5410	28	34-1	"	Spont.	S.B.	1550	"	Hydrops foetalis
5455	30	40-6	"	"	L	2680	"	
5471	27	39-3	"	"	L	2300	"	
5651	35	38-4	"	"	L	2860	2nd degree	
6023	37	39-2	"	"	L	2200	"	
6272	32	40-0	"	"	L	2650	Episiotomy	
6725	26	39-6	"	Assisted Spont.	L	2722	"	
6735	24	30-4	"	Forceps for A.C.H.	L	700	"	
6935	27	41-3	"	"	L	3175	"	
7071	21	39-5	"	"	L	3250	"	
7096	24	41-5	"	Spont.	L	2390	Episiotomy	
7169	23	38-3	"	"	L	2750	"	
7332	22	37-4	"	"	L	3350	"	
7546	26	39-3	"	Forceps for A.C.H.	L	2600	"	
7659	19	40-4	"	Spont.	L	2720	"	
7706	35	39-0	"	"	L	3200	"	
7752	29	38-6	"	"	L	2480	"	
7776	20	39-6	"	Assisted	L	3062	1st degree Episiotomy	
7903	26	39-0	"	"	L	2980	"	
8071	25	38-0	"	Spont	L	3005	"	
8164	21	40-3	"	"	L	3580	"	
8194	29	38-5	"	"	L	3100	"	
NON-BOOKED								
1140	28	39-0	None	Forceps for A.C.H.	L	3000	Episiotomy	
1632	30	37-0	"	Spont.	N.D.	2620	"	
4189	31	38-4	"	"	L	2800	"	
4245	27	35-1	"	"	L	1360	"	
4400	21	"	"	"	N.D.	2438	"	
5328	22	35-2	"	"	L	2540	"	
6251	21	35-0	"	"	L	2130	"	
7451	21	?	"	Assisted	N.D.	2540	"	

UNCOMPLICATED BREECH DELIVERIES (Continued)
PAROUS BREECH PRESENTATION AND DELIVERIES
(WITHOUT COMPLICATIONS)

Reg. No.	Age	Prenatal Pregnancies Before 28 wks.,	Maturity	Obstetric Complication	Treatment	Result M. C.	Weight of Child gm.	Ephitomy or tear	Remarks
BOOKED									
82	30	—	37 3	None	Assisted	L	2608	—	
350	27	2	37 1	"	"	L	1820	1st degree	
425	27	3	40 2	"	Spont.	N.D.	5280	—	
466	30	—	38 2	"	"	L	2850	—	
501	31	4	41 4	"	"	L	3470	—	
504	30	1	40 3	"	"	L	3120	1st degree	
501	30	3	37 2	"	"	S.B.	2650	—	V.D.R.L. +; Hydrops foetalis
600	30	2	42 5	"	"	L	2948	1st degree	
726	25	—	42 6	"	"	L	3500	—	
791	29	1	40 2	"	"	L	3540	—	
806	36	5	41 3	"	Assisted Spont.	L	3345	—	
1052	39	9	38 1	"	"	L	3280	—	
1066	31	4	40 3	"	"	L	2850	—	
1319	35	8	36 5	"	"	L	2760	1st degree	
1336	32	3	35 2	"	"	L	1040	—	Anencephaly
1416	36	1	37 5	"	"	S.B.	2977	—	Laceration of Cervix Retained
1447	27	2	39 3	"	Assisted Spont.	L	3000	1st degree	
1486	27	1	38 1	"	Assisted	L	3220	—	
1494	29	1	?	"	"	L	3790	—	
1579	35	6	42 0	"	Spont.	L	2930	—	
1656	32	1	38 1	"	"	L	2977	—	
1696	28	2	39 4	"	"	L	2880	—	
1851	21	1	29 6	"	"	L	1620	—	
1868	26	3	39 6	"	"	S.B.	3175	—	Hydrops foetalis
2040	28	3	39 1	"	"	L	3350	—	
2126	28	2	40 3	"	Spont.	L	3175	—	
2135	41	1	38 6	"	Assisted	L	3515	—	Manual removal of Placenta
2178	31	5	30 6	"	Assisted Spont.	L	2580	—	
2454	39	4	40 0	"	"	L	3350	—	
2465	23	1	34 4	"	"	L	1770	—	
2543	24	1	42 5	"	"	L	3705	—	
2593	32	5	41 1	"	"	L	3430	—	
2763	38	6	40 5	"	"	L	4040	—	
2768	24	1	38 3	"	Assisted Spont.	L	3610	1st degree	
3130	29	3	43 3	"	"	L	3540	—	
3178	27	4	39 1	"	"	L	3067	—	
3272	30	1	38 5	"	"	L	2551	—	
3317	37	3	38 5	"	"	N.D.	1970	—	
3536	23	1	40 5	"	"	L	3550	—	
3546	38	10	36 3	"	"	L	3200	—	Manual Removal of Placenta
3601	32	3	37 5	"	"	L	2950	—	

UNCOMPLICATED BREECH DELIVERIES (Continued)

Reg. No.	Age	Prenatal Pregnancies Before 28 wks.	Prenatal Pregnancies After 28 wks.	Maturity	Obstetric Complication	Treatment	Result M.	Result C.	Weight of Child gm.	Epistomy or tear	Remarks
3870	26	—	1	35.2	None	Spont.	L	L	3450	—	
3909	26	—	1	40.5	"	"	L	L	3090	—	
4185	24	—	1	39.0	"	"	L	L	3770	1st degree	
4312	20	—	1	38.2	"	"	L	L	2800	1st degree	
4373	34	—	5	41.3	"	"	L	L	2900	—	
4391	33	—	6	35.4	"	"	L	S.B.	1800	—	Hydrocephalus
4392	26	—	2	40.1	"	"	L	L	3175	—	
4501	29	—	2	34.1	"	"	L	L	2450	—	
4556	23	—	1	40.2	"	"	L	L	3300	—	
4621	24	—	2	41.0	"	"	L	L	3350	—	
4703	25	—	3	39.1	"	"	L	L	3620	—	
4838	40	—	2	36.0	"	"	L	L	2370	—	
4974	28	—	2	38.5	"	Assisted Spont.	L	L	3100	—	
5020	23	—	1	34.4	"	"	L	N.D.	1820	—	
5189	31	—	2	?	"	"	L	Mac.	2300	—	
5331	26	—	2	41.2	"	"	L	L	2950	—	
5630	35	—	3	41.5	"	"	L	L	3050	—	
5668	28	—	1	36.3	"	"	L	L	3175	—	
5769	31	—	3	41.5	"	"	L	L	3650	—	
5800	34	—	2	32.0	"	"	L	Mac.	2180	—	
5844	30	—	3	42.2	"	Assisted Spont.	L	L	4140	—	
5978	22	—	1	40.6	"	L.S.C.S.	L	L	3175	—	Poor Obs. History with Breech Presentation
6076	32	1	8	40.1	"	"	L	L	4000	—	
6168	34	—	5	37.4	"	Spont.	L	L	1850	—	
6360	26	—	2	38.1	"	"	L	L	3160	—	
6398	32	—	6	43.3	"	"	L	L	3500	—	
6666	32	—	6	?	"	"	L	L	3700	—	
6814	41	—	6	40.0	"	"	L	L	3400	—	
7364	31	2	5	38.0	"	"	L	L	3660	—	
7493	29	—	1	43.0	"	"	L	L	3700	Epistomy	
7579	28	—	1	?	"	"	L	L	2120	—	
7635	33	—	3	40.6	"	"	L	L	2900	—	
7640	37	—	6	38.0	"	"	L	L	3380	—	
7774	40	—	4	35.0	"	"	L	S.B.	1780	—	Monstr
7948	29	—	2	38.5	"	"	L	L	2650	Epistomy	
7975	32	—	2	39.6	"	"	L	L	2948	—	
8061	27	—	2	44.4	"	"	L	N.D.	4005	—	Anencephaly
8121	26	—	1	35.6	"	"	L	L	2850	—	
NON-BOOKED											
523	33	—	7	35.2	"	Spont.	L	L	2470	—	Left Hydronephrosis and Left Ureteric Calculus
1068	33	—	6	?	"	Assisted Spont.	L	L	3530	—	
5400	24	—	1	41.5	"	Spont.	L	L	3550	—	
7825	30	—	2	39.2	"	Assisted	L	L	3440	Epistomy	

TABLE 15
COMPLICATED BREECH DELIVERIES
 (Excluding Breech by Version in Labour)

(Includes breech delivery where some obstetric complication is present, e.g. contracted pelvis, placenta praevia, twins, toxæmia, prolapsed cord, etc.)
 There were 80 mothers in this series, of whom 30 had twin deliveries. Of these 30 mothers, only 5 had both infants delivered by breech presentations.
 Altogether there were 85 babies, 79 in Booked and 6 in Non-booked cases.
 There were 24 primigravidae and 56 multiparæ.
 There were no maternal deaths.
 There were 7 stillbirths, 5 in Booked and 2 in Non-booked cases; 13 neo-natal deaths, 12 in Booked and 1 in Non-booked cases; and 1 macerated foetus in the Booked cases.

PRIMIGRAVID BREECH PRESENTATION AND DELIVERIES
 (WITH COMPLICATIONS)

Reg. No.	Age	Maturity	Obstetric Complications	Treatment	Result	Weight of child gm.	Episiotomy or tear	Remarks
BOOKED								
196	19	40-1	Contracted Pelvis	L.S.C.S.	L	2580	—	
601	19	37-6	Prolapse of Cord	Spont.	L	2930	—	
776	25	37-1	" "	Assisted	L	3000	Episiotomy	Laceration of cervix repaired
1178	22	33-4	Twin Pregnancy	Spont.	L	2100	—	2nd of twins
2150	35	40-2	Elderly Primigravida	L.S.C.S.	L	3147	—	
2694	24	40-4	Twin Pregnancy	Spont.	L	2270	2nd degree	Both twins
				"	L	2140	Episiotomy	1st R.S.A. 2nd L.S.A.
2803	25	38-6	P. E.	"	L	3280	—	
4045	26	?	Contracted Pelvis	L.S.C.S.	L	2725	—	
4500	22	38-1	Uterus Didelphys	"	L	2900	—	
5082	28	36-4	Contracted Pelvis	Spont.	L	2825	Episiotomy	
5447	30	38-1	" "	L.S.C.S.	L	3440	—	Contracted mid pelvic s. outlet
5634	23	40-4	Prolapse of Cord	Spont.	L	2948	Episiotomy	

COMPLICATED BREECH DELIVERIES (Continued)

Reg. No.	Age	Maturity	Obstetric Complications	Treatment	Result	Weight of child gm.	Episiotomy or tear	Remarks
					M. C.			
5778	20	36-5	Contracted Pelvis	L.S.C.S.	L	3200	--	
5832	29	30-0	" " P. E.	" Spont.	N.D.	3250	--	Baby—Bronchopneumonia (20th day)
5913	28	36-4	" " Twin Pregnancy	"	L	1830	Episiotomy	Both twins 1st twin R.S.A.
5967	30	38-1	" " Contracted Pelvis	"	L	2215	"	2nd of twins
6401	22	39-4	" " Prolapse of Cord	L.S.C.S.	L	2628	--	
6525	23	38-4	" " P. E.	"	L	3570	--	
7322	29	31-2	" " Prolapse of Cord	C.C.S.	N.D.	1466	--	
7392	23	40-6	" " Elderly Primigravida	Spont.	L	1990	--	
7419	39	38-3	" " Contracted Pelvis	"	L	2250	--	Both twins 1st twin L.S.A.
8211	35	39-4	" " Contracted Pelvis	L.S.C.S.	L	3110	--	2nd twin R.S.A.
				"	L	2750	--	2nd of twins—prolapse of cord
NON-BOOKED								
4135	26	34-6	P.E. Severe	Spont.	L.	1450	Episiotomy	
6745	22	39-1	Contracted pelvis	"	L.	2003	"	

COMPLICATED BREECH DELIVERIES (Continued)
PAROUS BREECH PRESENTATION AND DELIVERIES
(WITH COMPLICATIONS)

Reg. No.	Age	Prenatal Before 28 wks.	Prenatal After 28 wks.	Maturity	Obstetric Complications	Treatment	Result M. C.	Weight of child gm.	Epistomy or tear	Remarks
BOOKED										
101	30	—	2	38.5	Contracted Pelvis	L.S.C.S.	L	2890	—	
166	30	—	2	39.6	Prolapse of Cord	Assisted	L	3430	—	
221	40	1	2	37.3	Twin Pregnancy	Spont.	L	2637	—	2nd of twins
230	39	—	6	32.6	"	"	L	1950	—	2nd of twins
393	44	—	6	34.6	Prolapse of Cord	Assisted	L	2400	—	
946	25	—	1	38.6	P. E.	J.S.C.S.	L	2863	—	
1702	27	—	3	40.0	Contracted Pelvis	Spont.	L	3220	—	
2184	37	—	5	36.6	P. E.	"	L	2140	—	
2232	24	—	4	32.4	Twin Pregnancy	"	L	1600	—	
2451	31	—	3	39.1	" P. E.	Assisted	L	1460	—	Both twins 1st twin L.S.P., 2nd twin R.S.P.
2507	37	—	2	33.0	Placenta Praevia	Spont.	L	2665	—	
2531	28	—	3	41.0	Contracted Pelvis	L.S.C.S.	L	2400	—	
2638	38	—	3	44.0	Twin Pregnancy	"	L	2835	—	
2752	30	—	4	39.5	P. E.	Spont.	L	2520	—	
2771	42	—	6	35.5	Prolapse of Cord	"	L	2900	—	
2841	42	3	8	40.6	Accidental Haemorrhage	"	L	1701	—	2nd of twins
3086	28	—	4	36.4	Twin Pregnancy	"	L	3890	—	A.P.H., 1600c.c. manual removal of placenta
3119	25	—	3	37.3	Placenta Praevia Type IV	L.S.C.S.	L	2722	—	
3199	28	—	3	37.3	P. E.	Spont.	L	3050	—	
3278	35	—	5	41.2	Prolapse of Cord	"	L	2100	—	Pulmonary Tuberculosis
3532	30	—	2	28.6	Twin Pregnancy	L.S.C.S.	L	3160	—	2nd of twins
3614	37	—	8	39.6	Placenta Praevia Type III	Spont.	L	1420	—	
3708	24	—	2	35.1	P. E.	"	L	2750	—	2nd of twins
3823	28	—	2	36.0	Twin Pregnancy	"	L	2620	—	
4034	30	—	3	29.5	Prolapse of Cord	"	L	3030	—	1st of twins
4497	36	—	8	39.2	Twin Pregnancy	"	L	1240	—	2nd of twins
4461	32	—	5	39.4	"	"	L	2750	—	"
4478	36	—	1	39.4	"	"	L	2180	—	"
4570	22	—	1	44.4	Contracted Pelvis	J.S.C.S.	L	3200	—	"
4602	32	1	3	39.2	Prolapse of Cord	Spont.	L	2110	—	2nd of twins
4610	36	—	4	40.3	Twin Pregnancy	"	L	3090	—	2nd of twins

COMPLICATED BREACH DELIVERIES (Continued)

Reg. No.	Age	Precious Pregnancies Before 28 wks.	Precious Pregnancies After 28 wks.	Maturity	Obstetric Complications	Treatment	Result	Weight of child gm.	Episiotomy or tear	Remarks
4001	25	—	1	37-3	Twin Pregnancy	Spont.	L	1800	—	2nd of twins
4965	38	—	4	43-0	" "	"	L	2700	—	1st of twins
5240	25	—	1	29-0	Placenta Praevia Type II Prolapse of Cord	"	L N.D.	1180	—	
5246	28	—	3	31-1	Twin Pregnancy	"	L N.D.	1440	—	2nd of twins
5571	36	3	5	38-6	Contracted pelvis	"	L	3040	—	
5650	28	—	3	34-3	P.E.	"	L		—	
5932	27	—	4	38-2	Twin Pregnancy Placenta Praevia type I.	"	L	2100	—	2nd of twins
5966	31	—	2	35-6	Twin Pregnancy	"	L	2090	—	" "
6314	22	—	1	40-5	Twin Pregnancy	"	L N.D.	2360	—	2nd of twins
6397	34	—	8	37-4	Contracted pelvis P.E.	L.S.C.S. Spont.	L L	3260	—	2nd of twins
6402	32	—	4	33-3	Twin Pregnancy	"	L	1620	—	
6435	25	—	2	35-6	Placenta Praevia type II Contracted pelvis	L.S.C.S. L.S.C.S.	L N.D. L N.D.	1160	—	Subtotal Hysterectomy
6468	24	—	1	27-4	Placenta Praevia type IV Contracted pelvis	Spont.	L	1005	—	2nd of twins
6528	27	—	2	36-2	Contracted pelvis	"	L	2150	—	
6575	28	1	2	39-6	Prolonged labour Uterine inertia	"	L S.B.	2750	—	
6637	27	—	3	41-0	Twin Pregnancy	"	L L	2950	—	2nd of twins
7175	34	—	5	29-2	Twin P.E.	"	L L	2350	—	" "
7533	26	—	2	39-1	Twin Pregnancy Prolapse of Cord	"	L L	2368	—	
7592	29	1	1	30-2	P.E. Concealed Accidental Haemorrhage	C.C.S.	L L N.D.	1200	—	
7838	25	—	1	33-0	Hydramnios	Spont.	L Mac.	1700	—	2nd of twins
8145	23	—	1	37-0	Twin Pregnancy	"	L L	2910	—	1st of twins
8312	30	—	3	30-1	" "	"	L L	2560	—	
NON-BOOKED										
736	31	—	2	38-4	Ante-partum haemorrhage of unknown origin	Spont.	L L	2810	—	A.P.H. about 300c.c. mixed with urine
1166	26	—	1	33-3	Contracted pelvis Prolapse of Cord	Spont.	L L	1650	1st Degree	Both twins Breach 1st L.S.A. 2nd L.S.A.
2566	30	—	3	32-4	Twin Pregnancy Placenta Praevia type IV	L.S.C.S.	L N.D.	1720	—	Presentation & Prolapsed cord

TABLE 16

FACE AND BROW PRESENTATIONS
(Including Anencephalus)

- I). Face Presentation:—
 There were 14 cases, all Booked.
 There were no maternal deaths.
 There were 5 stillbirths (4 were anencephalic foetuses and 1 second twin), 2 neo-natal deaths, (1 was anencephalic and 1 first of twins).
- II). Brow Presentation:—
 There was one case. It was an emergency admission after failed forceps. Caesarean Section was carried out. The infant died 19 hours after delivery.

Reg. No.	Age	Previous Pregnancies		Maturity	Position	Deliveries	Birth Weight		Results		Remarks
		Before 28 wks.	After 28 wks.				M.	C.			
BOOKED											
790	33	—	5	38.4	L.M.A.	Spont.	2950	L	N.D.	Anencephaly	
2323	32	—	7	44.5	L.M.A.	"	3570	L	S.B.	"	
2388	37	—	9	44.1	R.M.P.	"	3340	L	S.B.	"	
2524	35	2	7	34.4	L.M.A.	Manual Rotation	1750	L	S.B.	"	
2830	28	—	1	39.3	R.M.P.	"	3500	L	L	"	
					→ M.A.						
3404	24	—	3	38.5	L.M.A.	Spont.	2130	L	S.B.	2nd Twin, Proapsed Card. Bronchopneumonia. Prolapse of cord.	
3641	31	—	1	38.4	R.P.P.	Low Forceps	2680	L	L	"	
					→ L.M.A.						
3643	25	—	2	38.2	L.M.A.	Spont.	2180	L	L	"	
4042	24	—	1	38.2	L.M.A.	"	2650	L	L	"	
4077	31	1	3	35.5	R.M.A.	"	3040	L	L	"	
4460	36	—	1	35.5	L.M.A.	"	1300	L	N.D.	1st Twin.	
6159	50	—	2	43.6	L.M.A.	"	3300	L	L	"	
6588	24	—	3	43.6	L.M.A.	"	3830	L	L	"	
7358	40	—	6	38.5	L.M.A.	"	1435	L	S.B.	Hydrannios; Anencephaly	
NON-BOOKED											
6958	35	—	1	?	Brow	L.S.C.S.	3200	L	N.D.	Failed Forceps. Subdural Haemorrhage.	

TABLE 17

TRANSVERSE AND OBLIQUE LIE

There were 30 cases of shoulder presentation in labour, 20 Booked and 10 Non-booked.

There were no maternal deaths.

There were 10 stillbirths, 6 Booked and 4 Non-booked; 6 neo-natal deaths, 3 Booked and 3 Non-booked.

Reg. No.	Age	Pregnancies		Maturity	Obstetric Complication	Size of Os when diagnosed	Treatment	Result		Weight of Child gm.	Remarks	
		Before 28 wks.	After 28 wks.					M.	C.			
BOOKED												
340	29	—	4	40	Contracted Pelvis Shoulder Presentation	9 cm.	Spont. Version	L.	N.D.	2800		
395	34	—	8	35-4	"	F.D.	Internal Version	L.	L.	2470		
407	39	—	3	30-1	Previous L.S.C.S. Mixed Accidental Hemorrhage	3 cm.	Bipolar Version Followed by Craniotomy	L.	S.B.	1913		
734	35	—	2	38 5	Shoulder Presentation Prolapsed Right Arm; P.E.	6 cm.	Internal Version Breech Extraction	L.	L.	2665	Cervical Laceration	
1918	41	—	4	34-4	Placenta Praevia Shoulder Presentation	5 cm.	L.S.C.S.	L.	L.	2460		
3844	28	—	4	35-4	Shoulder Presentation	F.D.	Spont.	L.	Mac.	2100	Baby—Duplicatio Corporis.	
3871	31	—	3	39-1	Oblique Lie Prolapsed Cord	F.D.	Breech Extraction	L.	L.	2850	Prolapsed Cord.	
3874	29	—	4	40	Shoulder Presentation 2nd Twin	F.D.	Internal Version Breech Extraction	L.	L.	3740	Binocular Twin Pregnancy— 2nd Twin.	
3985	21	—	—	34	Contracted Pelvis Shoulder Presentation	2 cm.	L.S.C.S.	L.	L.	2670	P.P.S.	
4737	27	1	3	39-5	Transverse Lie Placenta Praevia IV	—	L.S.C.S.	L.	L.	2665	P.P.S.	
5028	29	—	7	35-6	Transverse Lie Prolapsed Arm & Cord	5 cm.	Internal Version Breech Extraction	L.	S.B.	2200		
5777	23	—	2	?	2nd Twin Transverse Lie	F.D.	Spontaneous Evulsion	L.	N.D.	1390	Binocular Twin Pregnancy— 2nd Twin.	
5952	33	—	3	33-5	Transverse Lie	F.D.	Spont.	L.	Mac.	1200		
6251	21	—	1	39-4	Contracted Pelvis Shoulder Presentation	1 cm.	L.S.C.S.	L.	L.	4260		
6428	23	2	3	38-4	Contracted Pelvis Shoulder Presentation	—	L.S.C.S.	L.	L.	2890		

TRANSVERSE AND OBLIQUE LIE (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Pregnancies After 28 wks.	Maturity	Obstetric Complication	Size of Os when diagnosed	Treatment	Result		Weight of child gm.	Remarks
								M.	C.		
6861	37	—	—	?	Shoulder Presentation Accidental Haemorrhage	—	C.C.S.	L	L	2070	
7873	26	1	6	29-4	Placenta Praevia Type III Transverse Lie	4 cm.	Internal Version Breech Extraction	L	S.B.	1400	
7895	27	—	2	37-3	Shoulder Presentation	F.D.	Internal Version Breech Extraction	L	Mac.	2400	
7944	30	—	1	38-3	Shoulder Presentation Naegele's Pelvis	—	C.C.S.	L	N.D.	3060	Previous C.S. Hepato-splenomegaly.
8312	30	—	3	30-1	Transverse Lie Twin Pregnancy	7 cm.	Internal Version	L	L	2220	Binovular Twin Pregnancy— 2nd Twin.
NON-BOOKED											
1706	25	—	2	?	Shoulder Presentation Placenta Praevia Type III	6 cm.	Bipolar Version	L	S.B.	1474	
1829	41	—	11	32-5	Shoulder Presentation	4 cm.	L.S.C.S.	L	L	1940	P.P.S.
2766	30	—	1	28-4	Shoulder Presentation	2 cm.	Spont.	L	N.D.	1080	
2837	32	—	2	40-1	Pre-clampsia Twin Pregnancy	7 cm.	Internal Version & Breech Extraction	L	N.D.	1950	2nd Twin
3605	37	—	4	40-4	Shoulder Presentation Pre-clampsia	F.D.	Internal Version	L	S.B.	2588	
3862	26	—	3	39-3	Shoulder Presentation Oblique Lie	F.D.	Internal Version Breech Extraction	L	S.B.	1250	
5134	30	—	6	40-4	Shoulder Presentation Prolapse of Cord	F.D.	Internal Version Breech Extraction	L	L	3325	Uniovular Twin Pregnancy— 2nd Twin.
6649	36	—	2	40-4	"	—	Norton's Extra peritoneal C.S.	L	Mac.	2551	Fibroid of Cervix.
7492	21	—	1	?	Contracted Pelvis Shoulder Presentation	4 cm.	L.S.C.S.	L	L	3570	
7999	41	1	9	34-5	Accidental Haemorrhage Transverse Lie Contraction Ring	—	C.C.S.	L	N.D.	1800	Prolapsed Left Arm. P.P.S.

TABLE 18
COMPOUND PRESENTATION

There were 5 cases of compound presentation during the year --- all in Booked cases

There were no maternal deaths

There was one stillbirth.

The condition occurred three times in twin pregnancies

Reg No	Age	Previous Pregnancies Before 28 wks	After 28 wks	Maturity	Complication	Treatment	Result M C	Weight of Child gm	Remarks
BOOKED									
660	39	1	6	33.3	Vertex Plus Left Hand	Bipolar Version Breech Extraction	L S B	1800	Mitral Stenosis Aortic Regurgitation Circular Disease Class III Prolapsed Cord Manual Removal of Placenta
1575	26	—	—	42.5	1st Twin Compound Presentation One Leg and One Hand	Assisted Delivery	L L	2800	L monovular Twins, Hydramnios
2505	34	1	6	33.5	2nd Twin Vertex Plus Right Hand	None	L L	1450	Binoovular Twins
6492	26	—	1	39.6	2nd Twin Vertex Plus Right Foot	Internal Version Breech Extraction	L L	2500	Binoovular Twins
7450	25	—	2	35.4	Vertex, Both Feet and Right Arm	Manual Replacement of Feet and Arm, Spont Delivery	L L	2340	

TABLE 19

MULTIPLE PREGNANCY

There were 83 cases of multiple pregnancy, 82 twin pregnancies and one case of triplets.
 There were 75 Booked and 8 Non-booked cases—14 primigravidae and 69 multigravidae.
 There were 163 babies, including one monster.
 There were 9 stillbirths, all in Booked cases; 29 neo-natal deaths, 25 Booked and 4 Non-booked.
 All the triplets were living.

Reg. No.	Age	Previous Pregnancies		Maturity		Presentation		Method of Delivery		Sex		Weight gm.		Type if known	Result		Remarks
		Before	After	1st	2nd	1st	2nd	1st	2nd	1st	2nd	1st	2nd		M.	F.	
6583/55	35	—	6	40.0	R.O.A.	Spont.	Spont.	M	F	3120	2950	Binovular	L	L	L	P.E.	
221/56	40	1	2	37.3	L.O.A.	"	"	M	M	2730	2637	Uniovular	L	L	L	Prolapsd Cord, 2nd of	
230	39	—	8	32.6	L.O.A.	"	"	M	M	1850	1950	"	L	L	L	Twins	
326	22	—	1	37.6	R.O.P.	"	"	M	M	2730	2500	Binovular	L	L	L	P.E.	
649	38	1	2	41.0	R.O.A.	"	"	M	F	2980	2100	"	L	L	L	P.E.	
1378	22	—	—	32.4	L.S.P.	"	"	F	F	1750	2100	"	L	L	N.D.		
1279	36	—	7	41.6	R.O.A.	"	"	F	M	2110	2360	"	L	L	L		
1491	23	—	—	37.3	L.O.P.	"	"	M	M	2780	2120	Uniovular	L	L	L	P.P.H. 800 c.c.	
1575	26	—	—	42.5	Compound R.S.P. & hand	Assisted	Mid-Forceps	M	F	2800	3572	"	L	L	L	Hydramnios	
2184	37	—	5	36.6	R.S.P.	Spont.	Spont.	F	F	2140	1600	"	L	L	L	P.E.	
2252	24	—	4	32.4	L.S.P.	"	Assisted	M	M	1520	1400	"	L	N.D.	N.D.		
2452	34	1	1	34.6	R.O.P.	"	Spont.	F	M	2480	2730	Binovular	L	L	L	P.E.	
2505	34	1	6	33.5	L.O.P.	"	"	M	F	1750	1450	"	L	N.D.	L		
2542	36	—	7	37.2	L.O.A.	"	M.R. + M.F.	M	M	3300	3050	Uniovular	L	N.D.	N.D.		
2638	34	—	3	43.0	L.O.A.	"	Spont.	M	M	2715	2520	Binovular	L	L	L	P.E.	
2694	24	—	—	40.4	R.S.P.	"	"	F	M	2270	2140	"	L	L	L	Accidental Haemorrhage.	
2771	42	—	6	35.5	L.O.P.	"	"	M	M	1899	1701	Uniovular	L	L	S.B.	1600 c.c.	
2784	34	—	2	38.2	L.O.A.	"	"	M	F	2350	2150	Binovular	L	L	L	Triplets	
2812	30	—	5	42.1	R.O.A.	Spont.	Spont.	M	M	2590	2070	Binovular	L	L	L	3rd	
											2830					L	

BOOKED

MULTIPLE PREGNANCY (Continued)

Reg. No.	Age Before 28 wks., 28 wks.,	Previous Pregnancies	Maturity		Presentation		Method of Delivery		Sex		Weight		Type if known	M.	Result		Remarks
			1st	2nd	1st	2nd	1st	2nd	1st	2nd	1st	2nd			1st	2nd	
3148	26	1	2	36 0	L.O.A.	R.O.A.	Spont.	Spont.	F	F	2000	2180	Uniovular	L	L	L	P.E.
3278	38	—	5	41-2	R.O.A.	L.S.A.	"	"	F	F	3340	3160	Binovular	L	S.B.	L	
3449	28	—	1	38 2	R.O.A.	L.O.A.	"	"	F	F	2820	2720	Uniovular	L	L	L	
3404	24	—	2	38-5	R.O.P.	L.M.A.	"	"	F	F	2120	2130	"	L	L	S.B.	Prolapsd Cord, 2nd of Twins
3614	37	—	8	39-6	R.O.A.	L.S.A.	"	"	F	F	2560	2750	"	L	L	L	P.E.
3621	23	—	—	34-3	R.O.A.	Papiraceus	"	"	M	M	2630	—	"	L	Mac	—	Foetus Papiraceus
3799	31	—	—	36-0	L.O.A.	R.O.P.	Mid-Forceps	Mid-Forceps	M	M	2300	2180	"	L	L	L	Prolapsd Cord, 2nd of Twins
3823	28	—	6	36-0	L.S.A.	L.O.A.	Spont.	Spont.	M	M	3030	2670	Binovular	L	L	L	
3874	29	—	4	40-0	R.A.A.	R.A.A.	Internal Version	Internal Version	F	F	3180	3740	"	L	L	L	
3905	32	—	2	36-5	L.O.A.	R.O.A.	Spont.	Spont.	M	M	2450	1980	Uniovular	L	L	N.D.	P.F.
4034	30	—	3	29-5	L.O.A.	R.S.A.	"	"	M	M	1190	1240	"	L	N.D.	N.D.	
4204	24	—	4	41-5	L.O.A.	R.O.P.	"	"	M	M	2650	2500	"	L	Mac	L	
4217	30	—	4	41-5	L.O.A.	R.O.A.	"	"	M	M	3317	3345	Binovular	L	L	L	
4397	36	—	8	39-2	L.O.A.	R.S.A.	"	"	M	M	2370	2730	Uniovular	L	L	L	
4460	36	—	1	35-5	L.M.A.	L.S.A.	"	"	F	F	1300	1660	"	L	N.D.	N.D.	
4461	32	—	5	39-4	R.O.A.	L.S.P.	"	"	F	F	2460	2150	"	L	L	L	
4570	22	—	2	44 4	L.O.A.	L.S.A.	"	"	F	F	1770	2110	"	L	N.D.	N.D.	Prolapsd Cord, 2nd of Twins
4610	36	—	4	40 3	L.O.A.	R.S.P.	"	"	F	F	2680	2710	Binovular	L	L	L	P.E.
4656	32	—	1	39-5	L.O.A.	R.O.A.	"	"	F	F	2080	2630	Uniovular	L	L	L	
4733	29	—	3	43-2	L.O.A.	R.O.A.	"	"	M	M	2910	2930	"	L	L	L	
4797	27	—	—	39 5	R.O.T.	L.O.A. →	Internal Version	Internal Version	M	F	3180	1900	Binovular	L	L	N.D.	P.F.
4901	25	—	1	37-3	R.O.A.	L.S.A.	Spont.	Spont.	M	M	2600	1800	Uniovular	L	S.B.	S.B.	
4985	38	—	4	43-0	R.S.A.	L.O.A.	"	"	F	F	2700	2575	Binovular	L	L	L	
5231	37	1	4	37-0	R.O.P.	R.O.A.	"	"	F	F	2280	1800	Uniovular	L	N.D.	N.D.	Prolapsd Cord
5246	28	—	3	31-0	L.O.A.	R.S.A.	"	"	F	F	1400	1440	"	L	N.D.	N.D.	
5370	32	—	—	44-3	L.O.P.	R.O.P.	Mid-Forceps	Mid-Forceps	M	M	3020	2350	"	L	L	L	Antepartum Eclampsia. P.P.H. 700 c.c.—800 c.c.
5615	32	—	3	37-0	L.O.A.	R.O.P.	Spont.	Spont.	M	M	2740	3140	"	L	L	L	P.E.; P.P.H. 640 c.c.
5650	28	—	2	34-3	L.O.A.	R.S.P.	"	"	M	M	2400	2100	Binovular	L	N.D.	N.D.	
5777	23	—	2	?	R.O.A.	L.A.A.	Spont.	Spont.	M	M	1330	1390	"	L	L	L	
5913	28	—	—	36-4	R.S.A.	L.S.A.	"	"	F	F	1830	1800	Uniovular	L	L	L	P.F.
5932	27	—	4	38 2	L.O.A.	R.S.A.	Spont.	Spont.	F	F	2300	2090	"	L	L	L	Placenta Praevia
5966	31	—	2	35-6	R.O.A.	R.S.P.	"	"	M	M	2680	2360	Binovular	L	L	N.D.	
5967	30	—	—	38-1	R.O.A.	R.S.A.	"	"	F	F	2730	2215	"	L	L	L	
6117	22	—	3	34 6	L.O.A.	R.O.A.	"	"	M	M	1700	2100	Uniovular	L	N.D.	N.D.	P.E.
6133	24	—	—	35-0	L.O.A.	R.O.P.	L.S.C.S.	L.S.C.S.	F	F	1900	1600	"	L	L	L	P.F.

MULTIPLE PREGNANCY (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks. 28 wks.	Maturity	Presentation		Method of Delivery		gm.		Weight Sex		Type if known	Result		Remarks
				1st	2nd	1st	2nd	1st	2nd	M.	F.		1st	2nd	
6232	28	—	34.5	L.O.A.	R.O.A.	Spont.	Spont.	F	F	1575	2200	Uniovular	L	L	
6275	35	5	37.1	L.O.A.	R.O.A.	"	"	F	F	2500	2520	"	L	L	
6286	28	2	40.1	L.O.A.	R.O.A.	"	"	M	M	3175	3320	Binovular	L	L	Pulmonary Tuberculosis
6378	22	—	42.5	L.O.A.	R.O.A.	"	"	M	M	2700	2450	Uniovular	L	L	
6397	34	—	37.4	L.O.A.	R.S.A.	"	"	F	F	2160	2370	"	L	L	P.F.
6492	26	—	39.6	L.O.P.	Compound	Mid-Forceps	Internal Version	M	M	2150	2500	Binovular	L	L	
					Vertex & Rt. Foot										
6578	27	—	36.2	L.O.A.	R.S.A.	Spont.	Spont.	M	M	2340	2150	Uniovular	L	L	
6637	27	3	41.0	R.O.A.	L.S.A.	"	"	M	M	3090	2950	Binovular	L	L	
6807	35	1	37.0	L.O.A.	R.O.A.	"	"	M	M	2800	2150	Uniovular	L	L	
7175	34	5	29.2	L.O.A.	R.S.A.	"	"	F	F	2230	2350	"	L	L	P.P.H., 1230 c.c.
7183	28	—	38.0	L.O.A.	L.O.A.	"	"	M	M	2150	2090	"	L	L	P.F.
7184	32	—	37.2	L.O.A.	R.O.A.	"	"	F	F	2230	2630	Binovular	L	L	P.F.
7356	23	—	39.3	R.O.P.	L.O.A.	"	"	M	M	2470	2700	Uniovular	L	L	P.L.
7392	23	—	40.6	L.S.A.	R.S.A.	"	"	M	M	1990	2250	"	L	L	P.L., Prolapsed Cord, 2nd of Twins
7591	27	—	37.4	R.O.P.	L.O.A.	"	"	M	M	2670	2670	"	L	L	
7681	25	1	36.4	L.O.A.	R.O.A.	"	"	M	M	2240	2250	"	L	L	
7845	28	—	38.2	R.O.A.	L.O.P.	"	"	F	F	2960	4500	"	L	L	Conjoined Twins
8001	29	3	35.6	L.O.A.	R.O.A.	L.S.C.S.	Spont.	F	F	2000	1630	Binovular	L	L	
8077	22	1	38.4	R.O.A.	L.O.A.	"	"	F	F	2680	2910	Uniovular	L	L	
8145	23	—	37.0	L.O.A.	R.S.A.	"	"	M	M	2500	2220	Binovular	L	L	
8312	30	—	30.1	L.S.A.	L.A.A.	"	Internal Version	F	F			"	L	L	
NON-BOOKED															
792	35	—	30.4	R.O.P.	R.O.P.	Spont.	Spont.	F	F	1900	920	Uniovular	L	L	P.F., Hookworm Infestation
909	24	—	35.2	L.O.P.	R.O.P.	L.S.C.S.	L.S.C.S.	F	F	1640	1780	"	L	L	Antepartum Eclampsia, Placenta Praevia Type 1, Uterine Inertia
1166	26	—	33.3	L.S.A.	L.S.A.	Spont.	Spont.	F	F	1650	2220	"	L	L	Prolapsed Cord Both Twins
2837	32	—	40.1	L.O.A.	L.A.P. → L.S.A.	Internal Version	Internal Version	M	M	2470	1950	Binovular	L	L	N.D.
3575	38	—	35.6	L.O.A.	R.O.A.	"	"	M	M	2480	2800	"	L	L	P.E., Terata Anadidyma
3599	29	—	42.6	L.O.A.	L.O.A.	"	L.S.C.S.	F	F	2700	2700	Uniovular	L	L	Monster, Hydramnios
5134	30	—	40.4	L.O.A.	L.A.A. → L.S.A.	Internal Version	Internal Version	F	F	2890	3325	"	L	L	Prolapsed Cord, 2nd of Twins
7988	38	2	37.0	R.O.A.	L.O.A.	"	"	M	M	2750	2700	Binovular	L	L	N.D.

TABLE 20

LABOUR FOLLOWING PREVIOUS CAESAREAN SECTION

(THIS TABLE DOES NOT INCLUDE ELECTIVE REPEAT CAESAREAN SECTION.)

There were 64 cases of labour following previous section, 63 Booked and 1 Non-booked cases.

Vaginal delivery occurred in 33 cases.

There were no maternal deaths.

There were 3 stillbirths, and 4 neo-natal deaths.

Reg. No.	Age	Pregnancies Before After 28 wks. 28 wks.	Maturity	Indication	Previous Caesarean Section		Type of Caesarean Section	Method of Delivery	Weight of child gm.	Result M. C.	Remarks
					Weight of Baby gm.	Caesarean Section					
BOOKED											
1	23	—	38-4	Contracted Pelvis	2800	L.S.C.S.	L.S.C.S.	2350	L	L	
101	32	2	38-5	"	2850	"	"	2890	L	N.D.	
407	39	—	30-1	Concealed Accidental Haemorrhage	1850	"	Bipolar " Version with Craniotomy	1310	L	S.B.	P.E.
479	25	—	40	Cephalo-pelvic Disproportion	3820	"	"	3402	L	L	
507	29	—	32-3	Severe P.E., Maternal Distress	3175	"	Spont.	2977	L	L	P.E.
894	24	—	?	Contracted Pelvis	3000	"	L.S.C.S.	3000	L	L	
946	25	—	38-6	Maternal Distress, Failed Trial Labour, Contracted Pelvis	2900	"	"	2865	L	L	
988	29	—	39-2	Placenta Praevia	?	"	Spont.	3000	L	L	Previous C.S. done at another hospital
1098	25	—	?	T.B. Hip. Contracted Pelvis	2800	"	"	3332	L	L	Trial Labour Successful
1142	24	—	39-5	Eclampsia	?	"	"	3200	L	L	Previous C.S. done at another hospital
1414	27	—	38-3	Contracted Pelvis	2950	"	"	3090	L	L	
1720	29	2	39-5	Placenta Praevia Type III	3050	"	L.S.C.S.	2972	L	L	
1806	40	—	40-6	Uterine Inertia, Foetal Distress	3030	"	Spont.	2870	L	L	
1954	40	2	40	Rt. T.B. Hip with Arthrodesis	3160	"	"	3050	L	L	Rt. T.B. Hip
2165	29	—	40-2	Ante-partum Eclampsia	2730	"	M.R. + M.F. Spont.	3260	L	L	
2286	22	—	32-6	Severe P.E., Breech	3000	"	"	2370	L	S.B.	
2405	25	—	40-1	Cephalo-pelvic Disproportion, Contracted Pelvis	2920	"	"	2657	L	L	Trial Labour Successful
2471	25	—	38-5	Contracted Pelvis	2650	"	"	2830	L	L	
2531	28	3	41	Breech Presentation, Contracted Pelvis	2575	"	L.S.C.S.	2835	L	L	
2807	26	—	41	Contracted Pelvis	3500	"	"	4111	L	L	
2864	24	1	33	"	3175	"	"	2608	L	L	

LABOUR FOLLOWING PREVIOUS CAESAREAN SECTION (Continued)

Reg. No.	Age	Pregnancies Before 28 wks., 28 wks.	Maturity	Indication	Previous Caesarean Section		Type of Caesarean Section	Method of Delivery	Weight of child gm.	Result	Remarks
					Weight of Baby gm.	M. C.					
2996	28	—	31-5	Contracted Pelvis, Previous C.S.	2800	I.S.C.S.	Spont.	2040	L	2	Previous C.S.
2996	24	—	35-4	Cephalo-pelvic Disproportion, Failed Forceps	3300	"	"	2800	L		
3087	25	—	34-3	Vaginal Septum	?	"	L.S.C.S.	1830	L		Uterus Dilelphys
3413	32	—	40-3	Failure of Dilatation of Os.	2722	"	Spont.	2720	L		
3810	28	—	40-5	Elderly Primigravida	3500	"	L.S.C.S.	3780	L		
4437	39	—	44	Cephalo-pelvic Disproportion Brow Presentation, Contracted Pelvis, Early Rupture of Membranes	2850	"	Mid Forceps	3200	L		Trial Labour Successful
4478	36	—	39-4	Contracted Pelvis, Elderly Primigravida	2600	"	L.S.C.S.	3200	L		Essential Hypertension
4712	21	—	39-1	Eclampsia	?	"	Spont.	2890	L		Previous C.S. done at another hospital
4795	25	—	43-2	Severe P.E.	2750	"	Spont.	3000	L		
4863	27	—	37-5	Contracted Pelvis	2900	"	"	2637	L		Previous C.S. done at another hospital
4885	20	—	41	Contracted Pelvis	?	"	"	3075	L		Previous C.S. done at another hospital
5018	28	—	?	Cephalo-pelvic Disproportion	3630	"	L.S.C.S.	2637	L		Previous C.S. done at another hospital
5045	35	—	41-3	Cephalo-pelvic Disproportion, P.F.	3500	"	M.R. + M.F.	3175	L		Trial Labour Successful
5053	27	—	39-5	Cephalo-pelvic Disproportion, Bad Obs. History	3232	"	Spont.	2863	L		
5233	26	—	39-6	Contracted Pelvis	?	C.C.S.	L.S.C.S.	2900	L		Previous C.S. done at another hospital
5429	33	—	39-3	?	3100	L.S.C.S.	"	3150	L		Occult Rupture of previous C.S. Scar. Previous C.S. done at another hospital
5547	39	1	34-3	?	?	"	"	3100	L		
5670	23	—	33-5	Borderline Contracted Pelvis	2950	"	"	2150	L	N.D.	
5715	31	—	38	Contracted Pelvis	2750	"	Spont.	2850	L		
5731	29	—	40-1	Cephalo pelvic Disproportion, Contracted Pelvis	3800	"	L.S.C.S.	3850	L		
5804	38	—	40	Elderly Primigravida, Non-engagement of Head at 41 weeks	2950	"	M.R. + M.F.	3430	L	S.B.	Trial Labour 29 hours
5983	26	—	35-5	Contracted Pelvis	3350	"	L.S.C.S.	3130	L		
6251	21	—	39-4	Cephalo-pelvic Disproportion	2855	"	"	3260	L		
6314	22	—	40-5	Contracted Pelvis	2700	"	"	3260	L		
6428	23	2	38-2	Transvers Lie; Contracted Pelvis	2013	"	"	2890	L		
6431	21	—	38-3	Contracted Pelvis	2950	"	C.C.S.	3120	L		
6796	38	—	43	Placenta Praevia Type IV.; P.F.	3650	"	Spont.	3300	L		
6824	25	1	40-3	Contracted Pelvis	3515	"	L.S.C.S.	3570	L		2 Previous C.S., P.P.S.

LABOUR FOLLOWING PREVIOUS CAESAREAN SECTION (Continued)

Reg. No.	Age	Pregnancies Before 28 wks. 28 wks.	Maturity	Previous Caesarean Section Indication	Weight of Baby gm.	Type of Caesarean Section	Method of Delivery	Weight of child gm.	Result		Remarks
									M.	C.	
6929	23	—	39	Antepartum Icthmia	2100	L.S.C.S.	Low Forceps	2608	L	L	Manual Removal of Placenta
7011	26	—	39 2	Prolonged Labour, Foetal Distress	2840	"	Mtd Forceps	3232	L	L	Previous C.S. done at another
7145	30	—	35-1	"	"	"	"	2850	L	L	hospital
7224	24	—	40 1	P.F., Accidental Haemorrhage	2810	"	Spont.	3005	L	N.D.	Trial Labour Successful
7231	27	—	38-2	Placenta Praevia Type III	2150	"	"	3070	L	L	
7562	38	—	38-5	Foetal Distress	3000	"	L.S.C.S.	3948	L	L	
7586	23	1	38	"	3050	"	"	3100	L	L	
7630	27	—	39 2	Contracted Pelvis—Naegele's	3050	"	C.C.S.	2850	L	L	Contracted Pelvis
7742	22	—	41 6	Contracted Pelvis	2608	"	M.R. + M.F.	3289	L	L	2. Previous C.S., P.P.S.
7931	24	—	39 5	Contracted Pelvis	2608	"	Low Forceps	3402	L	L	Trial Labour Successful
7944	30	—	38 3	Cephalo pelvic Disproportion	?	"	C.C.S.	3060	L	N.D.	Previous C.S. done at another
8008	25	—	40 3	Cephalo-pelvic Disproportion.	2600	"	Mtd Forceps	2900	L	L	Previous C.S. done at another
8080	25	—	39 5	Failed Trial Labour	2760	"	"				Naegele's Pelvis. Hepato Spleno
8213	25	—	38	Contracted Pelvis	3200	"	L.S.C.S.	2580	L	L	Mtd Contracted Uterus
				Contracted Pelvis		"	Spont.	3118	L	L	Labour Successful
NON-BOOKED											Trial Labour Successful
643	20	—	44 4	Contracted Outlet	3150	L.S.C.S.	L.S.C.S.	2750	L	L	

TABLE 21

CONTRACTED PELVIS

There were 264 cases of contracted pelvis, 249 in Booked and 15 in Non-booked cases.

There were no maternal deaths.

There were 5 stillbirths, 3 in Booked and 2 in Non-booked cases; 10 neo-natal deaths, all in Booked cases.

There was no induction of labour in this series.

Trial labour was successful in 22 out of 37 cases.

Morbidity was noticed in 44 cases.

There was one set of twins in the Non-booked cases.

Reg. No.	Age	Previous Pregnancies Before 28 wks.	Maturity	Type of Pelvis	Pelvic Measurements		Surgical Induction	Management		Method of Delivery	Length of Labour		Weight of Child	Result	Morbidity	Previous C.S.	Remarks
					O.C.	I.D.		I.S.	Trial of Labour		1st Stage	2nd Stage					
1	23	—	38.4	Gynaecoid	9.6	11.5	—	—	—	L.S.C.S.	—	4:40	2350	L	—	—	—
91	24	—	39.5	Anthropoid-Gynaecoid	10.7	11.3	—	—	—	Spont.	—	—	2268	L	—	—	—
101	32	—	38.5	Android	10	11.8	—	—	—	L.S.C.S.	—	—	2890	L	N.D.	—	P.P.S.
116	28	—	40.1	Gynaecoid	9.5	11.9	8.6	—	—	Spont.	—	15:43	2580	L	—	—	—
150	23	—	38.4	Anthropoid-Gynaecoid	11.6	11.3	—	—	—	"	—	13:30	3750	L	—	—	—
174	36	—	37.4	"	11.0	11.3	—	—	—	"	—	9:05	2500	L	—	—	—
177	21	—	40.5	"	10.0	11.7	—	—	—	"	—	39:40	2820	L	—	—	—
237	18	—	38.4	Platyphelloid-Gynaecoid	9.9	12.6	—	Failed	—	L.S.C.S.	—	—	2948	L	—	—	P.F.
330	24	—	45.4	Platyphelloid-Gynaecoid	8.1	12.5	—	—	—	"	—	—	2900	L	—	—	Plective C.S.
334	32	—	39.4	Gynaecoid	10.9	11.0	—	—	—	"	—	—	2982	L	—	—	Plective C.S.
340	29	—	40	"	9.8	11.8	—	—	—	Spont.	—	12:50	2800	L	N.D.	—	Uterine Fibroid, Shoulder Presentation, Spontaneous Version to Breech, to Breech, Assisted, Plective C.S.
346	23	—	41.3	"	9.8	11.0	—	—	—	L.S.C.S.	—	—	3629	L	—	—	—
352	25	—	39.2	"	10.0	11.7	—	—	—	Spont.	—	6:00	2700	L	—	—	—
400	34	—	38.6	"	10.1	12.7	8.6	—	—	L.S.C.S.	—	—	2700	L	—	—	—

BOOKED

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Maturity	Type of Pelvis	Pelvic Measurements		Surgical Indication	Trial of Labour	Method of Delivery	Length of Labour		Weight of Child	Result	Morbidity	Previous C.S.	Remarks	
					O.C.	T.D.				1st Stage	2nd Stage						
445	29	1	42-1	Platypelloid	9.2	12.2	—	—	L.S.C.S.	—	2855	L	—	—	—		
450	38	6	39.4	Anthropoid	12.2	10.8	—	—	Spont.	12:20	0:05	2810	L	—	—		
479	25	1	40	Gynaecoid	10.7	10.9	—	—	L.S.C.S.	—	3402	L	—	1	—		
495	21	—	39.2	"	11.0	11.4	—	—	Spont.	16:00	0:04	3100	L	—	—		
545	34	—	39	"	9.7	11.6	—	—	L.S.C.S.	—	2887	L	—	—	P.E.		
607	30	—	43	Anthropoid	12.3	11.4	9.1	—	Spont.	6:20	0:20	3450	L	—	—	—	
668	32	—	37.3	Gynaecoid	10.8	11.2	—	—	"	17:45	0:10	2760	L	—	—	—	
700	26	—	36	"	9.8	12.2	—	—	Manual Rotation & Mid Forceps	7:05	3:00	3232	L	—	—	P.E.	
735	26	—	38	Platypelloid	9.2	12.3	10.7	—	Spont.	16:15	1:25	2000	L	—	—	—	
772	31	—	38.4	"	9.9	13.1	—	—	"	5:25	0:05	3100	L	—	—	—	
894	24	—	?	Gynaecoid	10.7	11.2	—	—	L.S.C.S.	—	3000	L	—	—	—		
902	27	—	41-6	"	10.2	11.2	—	—	Spont.	5:45	0:05	3200	L	—	1	—	
946	25	—	38-6	"	9.8	12.2	—	—	L.S.C.S.	—	2863	L	—	—	—		
955	22	—	?	Andriod-Gynaecoid	10.6	11.4	—	—	Spont.	1:15	1:50	3000	L	—	—	—	
969	32	—	39.2	Gynaecoid	9.4	11.7	—	—	L.S.C.S.	—	3062	L	—	—	2	Electric C.S., P.P.S.	
987	23	1	41-5	Platypelloid	9.4	13.5	—	Successful	Spont.	3:00	1:15	3317	L	—	—	—	
1005	18	—	36.4	Gynaecoid	9.9	12.2	—	—	L.S.C.S.	—	2980	L	—	3 days	—	—	
1061	30	—	39.5	"	10.0	11.5	—	Successful	Manual Rotation & Mid Forceps	18:00	0:55	3340	L	—	—	P.E.	
1270	24	—	41-3	Naegale's Gynaecoid-Platypelloid	10.5	9.6	—	—	"	—	3400	L	—	—	2	Electric C.S.	
1290	35	—	43	"	9.7	13.3	—	—	L.S.C.S.	—	3317	L	—	—	—	Electric C.S.	
1293	38	—	38.4	Anthropoid	11.0	10.9	—	—	Spont.	5:35	0:05	3450	L	—	—	—	
1392	23	—	39.3	Gynaecoid	10.7	11.2	—	—	"	10:50	0:10	3657	L	—	—	—	
1414	27	—	38.3	Platypelloid	8.5	12.4	—	—	L.S.C.S.	—	3090	L	—	—	—	—	
1421	30	—	38.1	Anthropoid	13.1	11.1	—	—	"	—	2807	L	—	1 day	—	Electric C.S.	
1431	33	—	38.2	Anthropoid-Gynaecoid	11.0	11.2	—	—	Spont.	11:10	0:05	2880	L	—	—	—	P.E., T.B.
1456	28	—	36.4	Gynaecoid-Anthropoid	10.8	10.5	—	—	"	16:35	1:00	2550	L	—	—	—	—
1511	21	—	40	Anthropoid	10.6	11.1	9.3	—	Mid-Forceps	11:5	2:00	2892	L	—	—	—	—
1597	19	—	41	"	12.0	9.2	—	—	Spont.	5:55	0:05	2665	L	—	6 days	—	—
1652	24	—	39	Gynaecoid	10.0	11.3	—	—	"	7:25	0:05	2609	L	—	—	—	—
1696	28	—	39.4	"	9.7	12.4	—	—	"	5:00	0:05	2880	L	—	—	—	Breech Presentation
1717	32	—	36.6	Anthropoid	12.2	11.0	—	—	"	4:10	0:10	3120	L	—	—	—	—
1719	22	—	40.6	Gynaecoid	10.0	11.9	—	Unsuccessful	L.S.C.S.	—	2750	L	—	—	—	—	
1735	29	—	40.4	"	9.5	12.0	—	Successful	Spont.	10:10	0:20	3750	L	—	1 day	—	—
1771	22	—	39.3	"	9.9	11.7	—	—	"	14:17	0:03	2396	L	—	—	—	—
1782	31	—	39.5	Gynaecoid-Platypelloid	9.8	12.4	—	—	"	3:50	0:05	3220	L	—	—	—	—

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Pregnancies Before 28 wks. 28 wks.	Maturity	Type of Pelvis	Pelvic Measurements		Management	Length of Labour	Weight of Child	Result	Morbidity	Previous C.S.	Remarks
					O.C.	I.S.							
1833	32	—	5	Gynaecoid-Anthropoid	10.4	11.0	Spont.	13:16	3013	L	L	—	—
1951	26	—	2	Platyellipsoid	9.7	13.0	Successful	16:10	3800	L	L	—	—
1964	21	—	—	"	9.1	12.7	Failed	—	3175	L	L	—	—
1966	23	—	—	"	11.7	11.2	L.S.C.S.	24:10	3140	L	L	—	—
1982	25	—	1	Anthropoid	9.8	12.3	Spont.	5:10	2810	L	L	—	—
2000	27	—	—	Gynaecoid	9.5	13.4	"	5:20	3200	L	L	—	—
2009	38	—	—	Platyellipsoid-Gynaecoid	9.8	12.5	"	13:30	2900	L	L	—	—
2014	24	—	—	Gynaecoid	10.9	11.9	Mid-Forceps	17:00	2948	L	L	—	—
2016	32	—	—	"	9.8	12.4	"	65:35	0:40	L	L	—	—
2071	19	—	—	"	9.9	12.2	Spont.	13:00	2780	L	L	—	—
2091	26	1	2	Platyellipsoid-Gynaecoid	9.7	11.0	L.S.C.S.	—	2560	L	L	2	Elective C.S.
2138	28	—	—	Gynaecoid	10.8	11.4	Spont.	14:50	3190	L	L	—	—
2144	21	—	—	Gynaecoid	11.5	11.8	Manual Rotation & Mid-Forceps	29:30	3300	L	L	—	—
2212	33	—	—	Anthropoid	9.6	12.5	Unsuccessful	—	3175	L	L	—	—
2272	24	—	1	Platyellipsoid	9.6	12.4	Successful	16:30	3062	L	L	—	—
2335	42	—	—	Gynaecoid	9.9	12.9	Spont.	23:20	2885	L	L	—	—
2356	27	—	—	Platyellipsoid	11.6	10.6	Mid-Forceps	29:40	3409	L	L	—	—
2391	26	1	2	Anthropoid	10.0	11.6	Spont.	15:25	3005	L	L	—	Syphilis
2405	25	—	1	Gynaecoid	10.6	11.4	"	13:50	2637	L	L	—	—
2471	25	—	1	"	9.5	11.3	L.S.C.S.	—	2830	L	L	—	—
2531	28	—	3	Platyellipsoid	9.3	12.8	"	—	2835	L	L	—	Breach Presentation
2675	27	—	—	"	9.5	12.4	Spont.	11:15	2740	L	L	—	—
2778	23	—	—	Gynaecoid	9.9	12.4	"	8:45	2900	L	L	—	—
2807	26	—	2	Platyellipsoid	9.5	12.1	L.S.C.S.	—	4111	L	L	—	Elective C.S.
2864	24	1	2	"	9.0	12.4	"	—	2608	L	L	—	—
2922	23	—	1	Anthropoid-Gynaecoid	8.2	11.8	"	—	2722	L	L	—	—
2945	29	—	—	Platyellipsoid	9.0	13.1	"	—	3632	L	L	—	—
2956	28	—	3	"	9.4	12.1	Spont.	2:40	2040	L	L	2	—
2978	25	—	1	Gynaecoid	9.9	12.0	"	2:40	3650	L	L	—	—
3060	20	—	—	"	9.8	11.3	"	3:30	3000	L	L	—	—
3070	30	—	4	Platyellipsoid	9.2	13.9	"	—	3020	L	L	—	1 day
3109	23	—	—	"	9.0	12.1	L.S.C.S.	—	3370	L	L	—	—
3171	23	—	—	Gynaecoid	9.9	11.2	Spont.	1:40	2850	L	N.I.D.	—	—
3174	33	2	7	Platyellipsoid	9.4	12.9	"	15:58	0:02	2910	L	—	—
3186	28	—	3	Gynaecoid	10.8	10.8	"	9:00	1:15	3240	L	—	—
3210	34	—	1	Platyellipsoid	10.5	11.4	"	9:45	0:05	3060	L	—	—
3264	22	—	3	Anthropoid-Gynaecoid	9.8	11.7	"	11:15	0:20	2650	L	—	—
3295	30	—	—	"	9.9	13.5	"	27:55	0:10	3850	L	—	—
3369	27	—	2	Platyellipsoid-Gynaecoid	11.0	11.8	"	3:55	0:25	3270	L	—	—

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Pregnancies After 28 wks.	Type of Pelvis	Pelvic Measurements		Surgical Induction	Management		Length of Labour 1st Stage	Weight of Child	Result		Precious C.S.	Remarks	
					O.C.	I.D.		T.S.	Trial of Labour			Method of Delivery	M.			C.
3339	21	—	—	40-4	10.7	11.0	—	Failed	L.S.C.S.	20:45	3100	L	L	—	Severe P.E.	
3373	29	—	—	Anthropoid	12.4	11.2	—	—	Spont.	—	2700	L	L	—	—	
3376	38	1	4	10-6	9.7	13.2	—	—	Spont. Abortion	—	—	L	L	1	—	
3413	32	—	—	Platyphelloid	11.9	11.3	—	—	Spont.	—	2720	L	L	—	—	
3422	24	—	—	Anthropoid	10.4	11.0	—	—	"	—	2600	L	L	—	—	
3440	26	—	—	Gynaecoid	10.9	11.1	9.2	—	"	—	6:30	2730	L	L	—	—
3448	28	—	—	Gynaecoid-Android	9.5	12.1	—	Successful	"	—	22:00	2920	L	L	—	—
3491	26	—	—	Gynaecoid	9.6	12.4	—	—	"	—	0:38	3000	L	L	—	—
3500	26	—	2	Platyphelloid	10.8	11.1	—	—	"	—	7:35	3420	L	L	—	—
3501	26	—	—	Gynaecoid	—	—	—	—	—	—	—	—	—	—	—	—
3651	27	—	1	Anthropoid	9.8	13.7	—	—	"	—	6:55	3050	L	L	—	—
3661	27	—	2	Platyphelloid	9.7	11.2	—	—	"	—	20:00	3150	L	N.D.	—	—
3663	21	—	2	Gynaecoid	9.7	13.2	—	—	"	—	6:30	2600	L	L	—	—
3674	30	—	2	Platyphelloid	9.7	12.5	—	Successful	"	—	8:00	3430	L	L	—	—
3695	26	—	1	Platyphelloid	10.8	11.4	—	—	"	—	10:15	2650	L	L	—	—
3698	41	—	2	Gynaecoid	9.7	13.8	—	—	"	—	7:50	3160	L	L	—	—
3724	32	—	8	Platyphelloid	10.7	11.8	—	—	"	—	7:00	3000	L	L	1 day	—
3744	30	—	30-1	Gynaecoid	9.5	11.8	9.0	—	"	—	3:48	3400	L	L	—	—
38-2	38-2	—	5	Anthropoid	10.7	11.8	—	—	"	—	3:45	3629	L	L	—	—
3805	35	—	2	Gynaecoid	12.2	11.0	8.9	—	L.S.C.S.	—	1:10	3780	L	L	—	P.E.
3810	28	—	2	Platyphelloid	10.5	11.4	—	—	Spont.	—	11:30	3100	L	L	—	—
3840	28	—	3	Anthropoid	9.7	14.2	—	—	"	—	—	2030	L	L	—	—
3903	19	—	—	Gynaecoid	11.3	11.4	—	—	"	—	—	—	—	—	—	—
3985	21	—	—	Gynaecoid	10.3	11.0	—	—	L.S.C.S.	—	—	2670	L	L	—	Shoulder Presentation
3995	20	—	—	Anthropoid	12.2	11.0	—	—	Spont.	—	13:30	2630	L	L	—	—
4007	33	—	—	Gynaecoid	10.6	11.0	—	—	"	—	15:45	2450	L	L	—	P.E.
4014	26	—	2	Platyphelloid	9.5	12.9	—	—	"	—	13:00	2540	L	L	—	—
4022	25	—	1	Gynaecoid	9.7	11.0	—	—	L.S.C.S.	—	—	2820	L	L	1	O-cult Rupture of Uterus, repaired, Elective C.S.
4026	24	—	—	Anthropoid	11.5	11.4	—	—	Spont.	—	46:00	2500	L	L	—	—
4045	26	—	—	Gynaecoid	8.8	11.1	—	—	L.S.C.S.	—	—	2725	L	L	—	—
4069	32	—	4	"	9.8	12.6	—	—	Spont.	—	3120	L	L	—	—	
4154	26	—	—	Anthropoid	11.4	11.2	—	—	"	—	9:40	2670	L	L	—	—
4216	24	—	2	Platyphelloid	9.7	12.6	—	—	L.S.C.S.	—	12:55	2910	L	L	2	Elective C.S.; P.P.S.
4243	26	—	2	Gynaecoid	9.9	10.1	—	Unsuccessful	"	—	—	2525	L	L	—	—
4246	24	—	1	"	11.0	11.2	—	—	Spont.	—	10:35	3060	L	L	—	—
4309	31	—	1	Platyphelloid	9.3	11.3	—	—	L.S.C.S.	—	—	2510	L	L	—	—
4319	29	—	2	"	9.8	12.7	—	—	Spont.	—	10:15	3570	L	L	—	P.E.

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Prenous Pregnancies Before 28 wks.	Type of Pelvis	Pelvic Measurements		Management		Length of Labour 1st Stage	Weight of Child	Result	Morbidity	Prenous P.S.	Remarks
				O.C. T.D. I.S.	Surgical Induction	Trial of Labour	Method of Delivery						
4321	37	—	Gynaecoid	10.2	11.4	—	L.S.C.S.	—	3270	L. S.B.	—	1	Incomplete Rupture of Uterus
4361	23	—	Platypteloid	9.2	12.9	—	Spont.	8.25	2750	L. L.	—	—	—
4448	24	—	Gynaecoid	10.4	11.4	—	"	17.15	2807	L. L.	—	—	—
4456	21	—	"	11.0	12.3	—	"	8.30	3340	L. L.	—	—	—
4478	36	—	Platypteloid	9.9	12.4	—	L.S.C.S.	—	3200	L. L.	—	—	—
4497	30	1	"	8.2	11.4	—	"	—	2830	L. L.	1 day	—	—
4554	26	1	41-2	9.9	13.2	—	Spont.	11.50	3040	L. L.	—	—	—
4558	19	—	Gynaecoid	11.3	11.4	—	"	9.30	3300	L. L.	—	—	—
4580	28	—	"	10.7	11.1	—	"	11.40	3925	L. L.	—	—	—
4589	30	—	"	9.9	12.7	—	"	26.55	2948	L. L.	—	—	—
4635	30	—	"	11.2	11.2	—	L.S.C.S.	—	2950	L. L.	2 days	—	—
4705	25	1	"	9.5	11.7	—	"	—	2350	L. L.	—	—	—
4769	29	—	Platypteloid	9.3	12.2	—	Spont.	7.10	3175	L. L.	—	—	P.F.
4773	35	—	Anthropoid	11.8	11.4	—	"	30.30	2700	L. L.	—	—	P.F.
4782	24	—	Platypteloid	9.6	12.6	—	Successful	5.25	2950	L. L.	—	—	P.F.
4785	22	—	Gynaecoid	10.3	11.0	—	"	31.05	3220	L. L.	—	—	—
4832	22	—	"	11.5	11.0	—	"	18.35	3000	L. L.	—	—	—
4871	26	—	Gynaecoid	9.7	11.9	—	"	—	3060	L. L.	—	—	—
4873	23	—	Platypteloid	9.0	Net	—	Successful	25.40	3450	L. L.	—	—	—
				Obtain-able									
4885	20	1	"	9.6	12.4	—	"	10.50	3075	L. L.	—	1	—
4906	21	—	Gynaecoid	9.7	11.9	—	Successful	21.15	3005	L. L.	—	—	—
4920	28	—	"	10.9	11.3	—	"	37.45	2750	L. L.	—	—	—
4929	24	—	Anthropoid	11.7	11.3	—	"	9.30	3050	L. L.	—	—	P.F.
4930	24	—	Platypteloid	9.4	12.4	—	"	10.10	3360	L. L.	—	—	—
5045	35	—	Anthropoid	12.4	11.4	—	Successful	18.00	3175	L. L.	—	1	—
5053	27	—	Gynaecoid	10.0	10.4	—	Manual Rotation & Mid Forceps	16.04	2863	L. L.	—	1	—
5082	28	—	Platypteloid	8.3	12.2	—	Spont.	10.40	2825	L. L.	—	—	Breech Presenta- tion
5210	27	—	Anthropoid	11.8	11.1	—	"	33.0	2670	L. L.	—	—	—
5212	34	—	Platypteloid	9.6	12.7	—	"	1.50	2950	L. L.	—	—	—
5233	26	1	"	8.8	11.8	—	L.S.C.S.	—	2900	L. L.	1 day	—	—
5251	35	—	Gynaecoid	9.9	12.2	—	"	—	2495	L. L.	1 "	—	P.F.
5285	24	—	"	9.8	12.6	—	Spont.	10.45	2820	L. L.	—	—	—
5322	20	—	"	10.3	11.4	—	"	40.10	3170	L. L.	1 day	—	—
5352	19	—	"	10.3	11.1	—	"	6.40	3030	L. L.	—	—	—
5360	25	—	"	9.4	11.8	—	L.S.C.S.	—	2500	L. L.	—	—	P.F.; Flective (C.S.
5364	24	—	Platypteloid	9.9	13.2	—	Spont.	18.00	2980	L. L.	—	—	—
5372	24	—	Gynaecoid	10.7	11.4	—	"	14.40	3000	L. L.	—	—	—

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Prenatal Pregnancies Before 28 wks. 28 wks.	Maturity	Type of Pelvis	Pelvic Measurements		Surgical Induction	Management		Length of Labour	Weight of Child	Result	Morbidity	Previous C.S.	Remarks
					O.C.	I.D.		I.S.	Trial of Labour						
5387	30	—	1	Gynaecoid	9.5	12.1	—	—	—	9:55	2950	L	—	—	—
5429	30	—	2	"	9.2	12.3	—	—	—	—	3150	L	—	1	—
5439	22	—	37-3	Andrioid	9.5	12.5	—	—	—	6:40	2700	L	—	—	—
5447	30	—	38-1	Gynaecoid	10.3	12.7	9.4	—	—	—	2740	L	1 day	—	—
5489	40	—	38-1	Anthropoid	11.3	11.1	—	—	—	15:18	2600	L	—	—	—
5513	30	—	39-3	Gynaecoid	11.3	11.4	—	—	—	9:30	3450	L	—	—	—
5571	36	3	38-6	Anthropoid	11.4	11.1	—	—	—	10:50	3040	L	—	—	Breech Presentation
5581	39	—	41	Gynaecoid	9.6	12.2	—	Unsuccessful	—	—	3150	L	—	—	—
5599	26	—	32-5	Anthropoid-Gynaecoid	11.1	11.3	—	—	—	—	2300	L	N.D.	2	—
5665	21	—	38-4	Gynaecoid	9.7	11.9	—	—	Spont.	10:20	3570	L	L	—	—
5670	23	—	33-5	"	9.8	13.3	—	—	L.S.C.S.	—	2150	L	N.D.	1	—
5715	31	—	38	"	9.5	12.1	—	—	Spont.	18:45	2850	L	L	1	—
5778	20	—	36-5	Platypelloid	9.7	12.7	—	—	L.S.C.S.	7:30	3200	L	L	4 days	Post-Partum Psychosis
5832	29	—	39	Gynaecoid	10.8	11.4	—	—	"	—	3250	L	N.D.	2	—
5896	27	—	40-3	Gynaecoid	10.0	11.9	—	—	Spont.	16:55	2980	L	L	—	—
5897	28	—	40-1	"	11.2	11.6	—	—	Mid Forceps	21:30	3005	L	L	—	—
5976	36	—	40-6	"	10.6	11.6	—	—	Spont.	1:25	2660	L	L	—	—
5983	26	—	35-5	Anthropoid	12.0	9.8	—	—	L.S.C.S.	—	3130	L	L	1 day	—
6027	23	—	41	Gynaecoid	9.8	12.4	—	—	Spont.	8:25	2570	L	L	3 "	—
6077	22	—	39	"	10.0	11.9	—	—	"	6:50	3050	L	L	—	—
6101	32	—	39-5	"	9.8	12.3	—	—	"	11:00	3110	L	L	—	—
6144	30	—	37-1	"	11.5	12.9	9.0	—	Mid Forceps	23:00	3420	L	L	2 days	—
6166	38	—	38	"	12.1	12.6	9.0	—	Low Forceps	2:15	2750	L	L	—	—
6197	32	—	41-2	"	10.2	11.4	—	—	L.S.C.S.	—	2680	L	L	—	P.E.; Elective C.S.
6204	24	—	42	"	10.2	11.0	—	—	Spont.	21:20	2320	L	L	—	—
6233	26	—	41-5	"	9.7	12.3	—	—	"	22:40	3000	L	L	—	—
6276	36	—	40-5	Naegle's	11.9	10.7	6.8	—	L.S.C.S.	—	2977	L	L	2 days	Elective C.S.; P.P.S.
6251	21	—	39-4	Gynaecoid	9.6	12.0	—	—	"	—	3260	L	L	2 "	1
6298	31	—	40-4	"	10.6	11.1	—	—	Spont.	11:00	2500	L	L	2 "	—
6314	22	—	40-5	"	11.0	11.1	—	—	L.S.C.S.	—	3260	L	L	3 "	1
6381	36	—	38-6	"	9.2	11.9	—	—	"	—	3147	L	L	—	Elective C.S. P.E.; Elective C.S.; P.P.S.
6428	23	2	38-2	Platypelloid	9.0	13.1	—	—	"	—	2890	L	L	1 day	1
6431	21	—	38-3	Gynaecoid	9.5	11.5	—	—	"	—	3120	L	L	—	1
6435	25	—	35-6	"	9.1	11.2	—	—	"	—	1160	L	N.D.	—	2
6468	24	—	37-4	Platypelloid	8.4	11.3	—	—	Spont.	5:50	1005	L	S.B.	—	Subtotal Hysterectomy
6487	19	—	39-4	Gynaecoid	9.2	11.9	—	—	L.S.C.S.	—	2590	L	L	—	P.E.
6491	22	—	39-4	Platypelloid	9.3	12.6	—	—	"	—	3629	L	L	—	Elective C.S.
6538	34	—	39	"	9.2	12.0	—	—	"	—	3400	L	L	1 day	—

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks. 28 wks.	Type of Pelvis	Pelvic Measurements		Surgical Induction	Management		Length of Labour 1st Stage 2nd Stage	Weight of Child	Result	Morbidity	Previous C.S.	Remarks
				O.C. T.D. I.S.	Radiological		Trial of Labour	Method of Delivery						
6574	25	—	Gynaecoid	9.4	11.6	—	—	Spont.	4:20	0:10	L	—	—	Placenta Praevia
6611	21	1	Naegle's	10.3	10.6	8.2	—	C.S.S.	—	—	L	1 day	—	Type I
6617	28	—	Platyelloid	8.8	12.2	—	—	L.S.C.S.	32:10	1:00	L	1 day	—	Effective C.S.
6622	20	—	Gynaecoid	9.6	11.6	—	—	Spont.	10:20	1:30	L	—	—	P.E.
6681	26	—	"	11.0	11.2	—	—	Low Forceps	13:55	0:05	L	—	—	—
6720	37	3	Platyelloid	9.7	12.7	—	—	Spont.	4:00	0:20	L	—	—	—
6732	18	—	Gynaecoid	9.5	11.9	9.5	—	—	—	—	L	—	—	—
6738	25	—	"	9.5	11.8	—	Failed	L.S.C.S.	—	—	L	3 days	—	—
6764	37	4	Platyelloid	8.5	12.2	—	—	Manual Rotation & Mid Forceps	23:10	2:15	L	—	—	—
6824	25	1	Gynaecoid	9.8	11.7	—	—	Spont.	—	—	L	1 day	2	P.P.S.
6844	32	1	Anthropoid	10.7	10.7	8.6	—	L.S.C.S.	10:30	1:20	L	—	—	—
6867	25	2	Gynaecoid	9.6	11.1	—	—	Mid Forceps	—	—	L	—	—	—
6925	23	—	Anthropoid	12	11.4	9.0	—	L.S.C.S.	19:15	0:15	L	—	—	—
6957	26	—	Gynaecoid	10.5	11.2	—	—	Low Forceps	8:15	0:35	L	—	—	—
6962	23	—	"	9.9	12.2	—	—	Manual Rotation & Mid Forceps	6:00	0:15	L	—	—	—
6975	26	—	"	10.0	11.8	—	—	Spont.	12:50	0:40	L	1 day	—	—
6976	25	—	"	10.7	11.1	—	—	Mid Forceps	13:25	0:35	L	1 "	—	P.E.
7024	23	—	"	9.6	11.5	—	—	Spont.	29:30	0:30	L	—	—	Elective Mid Forceps
7207	34	4	Platyelloid	9.5	12.8	—	Successful	Spont.	7:30	0:15	L	—	—	—
7226	33	2	Gynaecoid	8.8	11.0	—	—	—	4:00	0:15	L	—	—	—
7385	26	—	"	9.3	11.9	—	Failed	L.S.C.S.	24:00	—	L	—	—	—
7390	28	—	"	11.0	12.0	8.7	—	Low Forceps	40:25	2:10	L	—	—	—
7395	36	—	"	11.4	12.4	9.0	—	Mid Forceps	19:30	2:45	L	2 days	—	F. Lumpsia 1 ft (Antepartum)
7478	27	—	"	9.3	11.7	—	—	Spont.	19:30	0:10	L	—	—	—
7507	29	2	"	9.3	10.9	—	—	—	6:00	0:20	L	—	—	—
7536	20	—	Anthropoid	14.0	11.9	8.9	—	Mid Forceps	14:20	2:00	L	2 days	1	—
7550	26	1	Platyelloid	9.5	12.4	—	Failed	L.S.C.S.	9:45	0:10	L	1 day	—	—
7573	23	1	Gynaecoid	9.4	11.7	—	—	Spont.	—	—	L	—	—	—
7586	23	1	Platyelloid	9.1	12.8	—	—	L.S.C.S.	—	—	L	—	—	—
7630	27	3	Naegle's	9.0	11.0	—	—	C.C.S.	5:35	0:05	L	—	—	Prolapsed Cord
7676	24	1	Gynaecoid	11.0	12.5	9.3	—	Spont.	3:15	2:23	L	—	2	P.P.S.
7679	21	—	Platyelloid	10.3	13.1	9.3	—	Mid Forceps	11:50	1:05	L	1 day	—	—
7743	22	—	Gynaecoid	9.4	13.2	—	Successful	Manual Rotation & Mid Forceps	6:55	0:35	L	—	—	—
7768	19	—	Platyelloid	9.3	12.3	—	—	Spont.	12:30	2:00	L	1 "	—	—
7824	27	—	Gynaecoid	9.7	11.8	—	—	—	15:20	3:30	L	2 days	—	—
7827	33	—	"	9.9	12.0	—	—	Mid Forceps	8:45	0:45	L	—	—	—
7850	21	—	"	9.8	12.2	—	—	Spont.	15:30	0:05	L	—	—	—
7933	22	—	"	10.2	10.6	—	—	—	—	—	L	—	—	S.B.

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Maturity	Type of Pelvis	Pelvic Measurements		Management		Length of Labour	Weight of Child	Result	Novelty	Previous C.S.	Remarks
					O.C.	T.D.	Surgical Induction	Trial of Labour						
7944	30	—	38-3	Naegele's	11.7	11.6	—	—	—	3060	L	N.D.	1	
7960	32	7	40-3	Anthropoid	11.8	11.6	8.5	—	—	10:30	0:55	3350	L	L
7971	38	7	43	Gynaecoid	11.5	11.7	9.0	—	—	23:50	1:20	4050	L	L
8008	25	—	40-3	Platypelloid	9.8	13.8	—	—	—	8:15	0:25	2900	L	L
8020	27	—	38-1	Gynaecoid	9.4	11.1	—	—	—	7:30	1:30	2800	L	L
8031	41	—	31-4	"	9.6	12.8	—	—	—	17:35	0:05	3175	L	L
8037	23	—	39-1	Platypelloid	9.7	13.1	—	—	—	17:00	0:30	3005	L	L
8041	24	—	43-1	Gynaecoid	9.7	11.7	—	—	—	31:20	0:40	3240	L	L
8051	42	—	34-1	"	10.9	12.4	8.6	—	—	5:15	1:55	3320	L	L
8057	26	—	40-5	"	11.2	12.4	9.1	—	—	19:45	1:15	2250	L	L
8072	20	—	43-6	"	10.5	12.0	9.2	—	—	6:15	2:00	3000	L	N.D.
8080	25	—	39-5	Gynaecoid-Platypelloid	9.2	11.7	—	—	—	—	—	—	—	—
8204	17	—	39	Gynaecoid	8.7	10.8	—	—	—	41:10	0:45	2720	L	L
NON-BOOKED														
537	17	—	—	—	—	—	—	—	—	—	—	—	—	—
638	22	—	40-3	Android-Gynaecoid	9.0	11.9	—	—	—	—	—	—	—	—
1094	28	—	41-2	Gynaecoid	10.0	11.6	—	—	—	—	—	—	—	—
1166	26	—	39-6	Platypelloid	8.2	13.0	—	—	—	—	—	—	—	—
		1	32	Gynaecoid	8.5	11.7	—	—	—	—	0:35	1650	L	L
		—	—	—	—	—	—	—	—	—	—	—	—	—
1276	19	—	40-3	Naegele's	9.0	9.5	—	—	—	—	—	—	—	—
1453	38	—	43-5	Platypelloid & Gynaecoid	9.7	11.9	—	—	—	—	—	—	—	—
1998	30	—	39	Gynaecoid	9.6	12.1	—	—	—	—	—	—	—	—
2473	35	—	40-4	Platypelloid	9.3	13.7	—	—	—	—	—	—	—	—
3129	23	—	42	Gynaecoid	11.1	11.1	—	—	—	—	—	—	—	—
3746	21	—	39-2	Android	10.5	11.2	—	—	—	25:00	3:05	3400	L	L
3899	28	—	43.5	Gynaecoid	10.4	11.4	—	—	—	—	—	—	—	—
4323	23	—	39-5	"	9.7	11.7	—	—	—	—	—	—	—	—
6745	22	—	39-1	"	8.4	11.0	—	—	—	—	—	—	—	—
7492	21	—	?	Platypelloid	9.6	13.5	—	—	—	—	—	—	—	—
7955	27	—	42-4	Gynaecoid	9.3	11.8	—	—	—	—	—	—	—	—

Hepato-Splenomegaly; Uterus absent; Right Kidney P.P.S.

P.P.H. 600 c.c.

Antepartum Eclampsia

Flective C.S. T.B. Hip (Left) Twin Pregnancy

Pre-Eclampsia Ruptured Uterus Caesarean Hysterectomy Bilateral Oophorectomy

Pre-Eclampsia Elective C.S. Elective C.S. Breech Presentation

Shoulder Presentation

TABLE 22

FAILED FORCEPS

There was one case of failed forceps in the Non booked cases

The case was delivered by Caesarean Section

The infant was stillborn

Reg No	Age	Prenatal Before 28 wks	After 28 wks	Maturity	Failed Before Admission	Forceps After Admission	Cause of Failure	Treatment and Method of Delivery	Height of Child	Result	Remarks	
										M	C	
NON BOOKED 6958	35	---	1	?	Yes	---	Head Floating Too High For Forceps	L S C S	3200	I	N D	Blow Presentation

TABLE 23

PROLAPSE AND PRESENTATION OF CORD

There were 33 cases of prolapse and presentation of cord, 28 Booked and 6 Non-Booked.
There were no maternal deaths.

There were 8 stillbirths, 6 in Booked and 2 in Non-booked cases; 4 neo-natal deaths, 3 in Booked and one in Non-booked cases. Twin pregnancy occurred 8 times, 6 Booked and 2 Non-booked.

Reg. No.	Age	Previous Pregnancies Before 28 wks.	Maturity	Size of Os when Diagnosed	Cause of Prolapse or Presentation of Cord	Treatment	Result		Remarks
							M.	G.	
166	30	—	39.6	F.D.	L.S.A. Malpresentation	Assisted breech delivery	L	L	
230	39	—	32.6	"	Twin pregnancy	"	L	L	2nd Twin
601	19	—	37.6	8 c.m.	R.S.A. Malpresentation	Breech extraction	L	L	
660	39	1	33.3	7 c.m.	L.S.A. Malpresentation Compound presentation R.O.A. with left hand	Bipolar podalic version breech extraction	L	S.B.	Mitral stenosis with aortic regurgitation, class IV. Manual removal of placenta. Laceration of cervix—repaired.
776	25	—	37.1	8 c.m.	R.S.P. Malpresentation	Breech extraction	L	L	
1702	27	—	40	F.D.	"	Assisted breech delivery	L	L	
2480	33	7	44	"	R.O.A. Unknown	Nil, spontaneous delivery	L	L	
2752	30	—	39.5	"	L.S.A. Malpresentation	Immediate breech extraction	L	L	
3116	25	3	37.3	"	"	"	L	L	
3205	24	—	41.1	"	L.O.A. Unknown	Nil, spontaneous delivery	L	L	P.E. Pulsation absent, foetal heart not heard on admission
3404	24	—	38.5	"	Twin pregnancy	Digital reposition	L	S.B.	2nd Twin
3641	31	—	38.4	"	L.M.A. Malpresentation Brow, R.F.P. → Face Malpresentation	Digital reposition, immediate forceps delivery	L	L	
3708	24	—	35.1	8 c.m.	L.S.A. Malpresentation	Breech extraction	L	L	
3799	31	—	36	F.D.	Twin pregnancy	Digital reposition, immediate forceps delivery	L	L	2nd Twin
3871	31	—	39.3	"	R.O.P. Malpresentation R.A.A. Oblique lie	Internal version, immediate breech extraction	L	L	
4570	22	—	44.3	"	Twin pregnancy	Nil, spontaneous delivery	L	N.D.	2nd Twin
5028	29	—	35.6	5 c.m.	L.S.A. Malpresentation Shoulder R.A.A. Malpresentation	Internal version and breech extraction	L	S.B.	
5240	25	—	8 c.m.	29	L.S.A. Malpresentation	Breech extraction	L	N.D.	

BOOKED

PROLAPSE AND PRESENTATION OF CORD (Continued)

Reg. No.	Age	Previous Pregnancies Before After 28 wks. 28 wks.	Maturity	Size of Os when Diagnosed	Cause of Prolapse or Presentation of Cord	Treatment	Result		Remarks
							M.	C.	
5246	28	—	31-1	F.D.	Twin Pregnancy	Assisted breech delivery	L	N.D.	2nd Twin
5597	26	—	40-3	F.D.	R.S.A. Malpresentation L.O.A. Unknown.	Assisted breech delivery	L	L	Prolapsed Cord on Admission
5634	23	—	40-4	"	L.S.A. Malpresentation	Digital reposition, spontaneous delivery	L	L	
6158	26	—	41-2	"	L.O.A. Unknown	Assisted breech delivery	L	L	
6875	25	—	38-4	7 c.m.	L.S.A. Malpresentation	Immediate forceps delivery	L	L	
6831	31	—	39	F.D.	L.O.A. Unknown	Nil	L	S.B.	
7392	23	—	40-6	"	Twin Pregnancy	Assisted breech delivery	L	L	2nd Twin, P.E.
7533	26	—	39-1	"	R.S.A. Malpresentation	Nil	L	L	
7575	27	—	42-2	"	L.S.A. Malpresentation	"	L	L	
7647	40	1	38-2	"	L.O.A. Contracted Inlet R.O.A. Unknown	"	L	S.B.	
NON-BOOKED									
1166	26	—	33-3	F.D.	Twin Pregnancy	Immediate breech extraction	L	L	Prolapse in both Twins
2610	28	—	32-4	3 c.m.	(1) L.S.A. (2) L.S.A. Both Malpresentation L.O.A.	L.S.C.S.	L	N.D.	
2842	30	—	32	5 c.m.	Placenta Praevia Type II R.S.A.	Manual replacement, spontaneous delivery	L	S.B.	
5134	30	—	40-4	F.D.	Placenta Praevia Type II Twin Pregnancy Shoulder Presentation L.A.A.	Internal version and breech extraction	L	L	2nd Twin
6649	36	—	40-4	5 c.m.	Shoulder Presentation L.A.A.	Norton's extraperitoneal C.S.	L	S.B.	Fibroid of Cervix

TABLE 24

POST-PARTUM HAEMORRHAGE

There were 74 cases of post-partum haemorrhage, 68 Booked and 6 Non-booked.
 There was one maternal death.
 Postpartum haemorrhage occurred 5 times in twin pregnancy.
 There were 71 cases of primary post-partum haemorrhage and 3 cases of secondary post-partum haemorrhage.

Reg. No.	Previous Pregnancies		Maturity	Method of Delivery	Duration of Labour	Amount of Haemorrhage (c.c.)		Cause or predisposing factor	Treatment State drug, dose and routine	Result		Remarks		
	Age	Before 28 wks.				After 28 wks.	Before delivery of placenta			Secondary P.P.H. of placenta	M.		C.	
BOOKED														
385	36	—	4	39-2	Spont.	10:05	—	680	—	2nd degree tear	Repair of Laceration	L	L	
750	32	—	2	39-2	"	5:48	—	—	—	Laceration of Cervix	Repair of Laceration; Vaginal Packing	L	L	
1082	34	—	2	40-2	M.R. + M.F.	8:12	700	—	—	"	Repair of Laceration	L	L	
1098	25	—	1	?	Spont.	7:40	700	—	—	Retained Placenta	Manual Removal of Placenta; Ergometrine 0.5 mg. I.M.I.	L	L	
1328	25	—	3	37-5	"	5:15	700	—	—	"	Ergometrine 0.5 mg. I.V.I.	L	L	
1489	24	—	2	37-6	"	12:05	—	1000	—	Unknown	Blood Transfusion 500 c.c. Manual Removal of Placenta; Ergometrine 0.5 mg. I.V.I.	L	L	
1540	19	—	—	40-5	"	19:45	(650)	—	—	Retained Placenta	Ergometrine 0.5 mg. I.V.I.	L	L	
1575	26	—	—	42-5	(1) Compound-Assisted	10:25	—	800	—	Atony	Ergometrine 0.5 mg. I.V.I.	L	L	Twin Pregnancy Hydramnios
1716	40	1	9	41	(2) Vertex-M.F. M.R. + M.F.	14:45	—	500	—	Retained Placenta	Manual Removal of Placenta; Blood Transfusion 500 c.c.	L	L	
2137	20	—	2	40-6	Spont.	5:55	—	500	—	"	Manual Removal of Placenta; Ergometrine 0.5 mg. I.V.I.	L	L	
2304	36	—	5	41	"	11:52	—	900	—	Laceration of Cervix	Repair of Laceration; Vaginal Packing	L	L	
2771	42	—	6	35-5	"	1:05	1600	—	—	Adherent Placenta	Manual Removal of Placenta; Blood Transfusion 1500 c.c.;	L	S.B.	Twin Pregnancy
2980	29	1	4	39-3	"	12:40	—	600	—	"	Ergometrine 0.5 mg. I.V.I.	L	S.B.	
2999	35	2	5	40-4	"	30:15	—	700	—	"	Manual Removal of Placenta; Ergometrine 0.5 mg. I.M.I.	L	L	Mitral Stenosis Class IV
3049	34	2	5	41	M.R. + M.F.	76:30	—	600	—	"	"	L	L	
3120	37	—	6	41-2	Spont.	7:30	—	600	—	Atony	Manual Removal of Placenta; Ergometrine 0.5 mg. I.V.I.	L	L	Prolonged Labour
											Ergometrine 0.5 mg. I.M.I.	L	L	

POST-PARTUM HAEMORRHAGE (Continued)

Reg. No.	Age	Pregnancies Before After 28 wks. 28 wks.	Maturity	Method of Delivery	Duration of Labour	Amount of Haemorrhage (c.c.)		Causa or predisposing factor	Treatment State drug, dose and routine	Result	Remarks
						Before delivery of placenta	After delivery of placenta				
3293	22	—	37-1	Spont.	12:43	—	—	Perineal Laceration	Repair of Laceration	L	
3635	26	2	41-5	"	4:40	—	500	Atony	Ergometrine 0.5 mg. I.M.I.	L	N.D.
4332	27	1	39-2	"	4:25	—	800	Retained Placenta	Manual Removal of Placenta, Ergometrine 0.5 mg. I.V.I., Pitocin 10 units I.M.I. & Pitocin drip 5 units Ergometrine 0.5 mg. I.V.I.	L	L
4333	38	1	40-6	"	4:15	—	520	Atony		L	L
4556	23	—	40-2	"	2:50	—	600	"	"	L	L
4716	36	2	38-6	"	38:25	—	1400	"	"	L	L
4752	24	—	41-2	"	2:05	—	660	"	"	L	L
4925	43	—	39-6	M.R. + M.F.	18:40	—	700	"	"	L	L
5210	27	—	?	Spont.	33:55	—	950	"	"	L	L
5240	25	—	29	"	5:15	—	(1200)	Placenta Praevia	Ergometrine 0.5 mg. I.V.I. Blood Transfusion 500 c.c.	L	N.D.
5370	32	—	44-3	(1) M.F. (2) Int.-Version + M.F.	95:0	—	700-800	Manual Remove of Placenta	Ergometrine 0.5 mg. I.V.I.	L	L
5472	27	1	42	Spont.	8:15	—	780	Retained Placenta Cervical Laceration	Manual Removal of Placenta Repair of Laceration	L	L
5630	35	—	41-5	Spont.	6:55	—	600	Atony	Ergometrine 0.5 mg. I.M.I.	L	L
5650	28	2	34-3	(1) Vx.-Spont. (2) Br.	5:10	—	640	"	Ergometrine 0.5 mg. I.V.I. Ergometrine 0.5 mg. I.M.I.	L	N.D.
5832	29	—	39	L.S.C.S.	—	—	800	C.S.	Ergometrine 0.5 mg. I.V.I.	L	N.D.
5942	40	—	40-3	Spont.	5:20	—	1000	Manual Remove of Placenta	Ergometrine 0.5 mg. I.V.I. Blood Transfusion 500 c.c.	L	L
6039	21	—	37-5	"	14:40	—	770	Atony	Ergometrine 0.5 mg. I.V.I.	L	L
6109	24	—	43-2	"	11:30	—	1200	"	"	L	L
6266	34	—	39-5	"	6:35	—	500	"	Ergometrine 0.5 mg. I.M.I.	L	L
6411	19	—	?	"	7:00	—	650	"	"	L	L
6427	19	—	39-1	"	20:00	—	760	Inadequate Retraction of Uterus	Ergometrine 0.5 mg. I.V.I.	L	L
6468	24	—	27-4	"	6:16	—	800	Manual Remove of Placenta	"	L	S.B.
6493	36	—	40-4	M.F.	16:55	—	650	Atony	"	L	L
6506	26	—	42-4	Spont.	2:33	—	900	"	Pitocin 10 units I.M.I.; Ergometrine 0.5 mg. I.V.I.	L	L
6596	25	—	42-6	"	10:55	—	900	"	Methergin 0.2 mg. I.V.I.	L	L
6625	33	—	43-2	"	9:25	—	700	"	"	L	L

POST-PARTUM HAEMORRHAGE (Continued)

Reg. No.	Age Before 28 wks.	Age After 28 wks.	Previous Pregnancies Before 28 wks.	Parity	Method of Delivery	Duration of Labour	Amount of Haemorrhage (c.c.) Before delivery	Amount of Haemorrhage (c.c.) After delivery	Secondary P.P.H.	Cause or predisposing factor	Treatment and routine	Result	Remarks
6669	34	—	4	39-6	Spont.	1:05	—	650	—	Atony	Methergin 0.2 mg. I.M.I. Methergin 0.2 mg. I.V.I. Repair of Lacerations	L	
6681	26	—	—	39-6	L.F.	12:55	—	1200	—	Episiotomy	Blood Transfusion 500 c.c. Ergometrine 0.5 mg. I.V.I.	L	
6709	30	—	4	?	Spont.	12:40	—	500	—	Atony	"	L	
6807	25	—	2	37	Spont.	3:15	—	1230	—	"	"	L	Twin Pregnancy
6963	24	—	2	37	M.F.	12:12	—	700	—	"	Methergin 0.2 mg. I.V.I. Ergometrine 0.5 mg. I.V.I.	L	P.E. Hydramnios
7023	26	—	1	40	Spont.	14:15	—	600	—	"	D, and C.	L	Secondary P.P.H.; D. & C. done in Q.M.H. 4 weeks after delivery
7056	30	1	4	39-4	"	9:10	—	—	Unknown	Retained Cotyledon of Placenta	"	L	
7112	25	—	3	37-3	"	2:00	—	500	—	Atony	Ergometrine 0.5 mg. I.M.I.	L	
7128	25	—	2	39-1	"	3:04	—	540	—	"	"	L	
7145	30	—	1	35-1	M.F.	29:05	—	600	—	"	Methergin 0.2 mg. I.V.I. Ergometrine 0.5 mg. I.M.I.	L	
7182	37	—	6	42	"	9:40	—	600	—	"	"	L	
7286	25	—	—	38-1	Spont.	7:20	—	600	—	"	Methergin 0.2 mg. I.V.I.	L	
7318	22	—	—	40-2	"	36:45	—	1000	—	"	"	L	
7371	40	—	3	40-6	L.S.C.S.	—	—	—	1500	"	Blood Transfusion 2500 c.c.	L	Secondary P.P.H. on 11th day
7381	24	—	—	38-5	Spont.	23:50	—	700	—	"	Methergin 0.2 mg. I.M.I. Ergometrine 0.5 mg. I.V.I.	L	
7444	26	1	3	41-5	"	4:40	—	1500	—	Cervical Tear; Manual Removal of Placenta	Repair of Laceration	L	
7464	21	—	—	41-4	"	45:15	—	—	550	Retained Placenta	Methergin 0.2 mg. I.V.I. D. & C.	L	Secondary P.P.H. on 13th day
7468	42	—	6	39-2	"	2:58	—	600	—	Atony	Ergometrine 0.5 mg. I.V.I.	L	P.E.
7696	23	—	2	39-3	"	4:10	—	750	—	"	Ergometrine 0.5 mg. I.M.I.	L	
7747	28	—	—	40-6	"	19:38	—	700	—	Retained Placenta	Methergin 0.2 mg. I.V.I. Manual Removal of Placenta	L	P.E.
7928	38	3	6	35-3	"	24:30	—	550	—	"	"	L	P.E.
8008	25	—	1	40-3	M.F.	8:45	—	600	—	Atony	Ergometrine 0.5 mg. I.V.I. Ergometrine 0.5 mg. I.M.I.	L	
8024	27	—	—	39-4	L.F.	22:10	—	660	—	Flabby Uterus Vaginal Tear	Methergin 0.2 mg. I.V.I.	L	
8053	25	—	—	40-5	"	7:55	—	500	—	Retained Placenta	Manual Removal of Placenta	L	Vaginal Septum
8056	24	—	3	40	Spont.	7:55	—	850	—	"	Methergin 0.2 mg. I.V.I. Manual Removal of Placenta;	L	
8087	32	—	—	38-4	M.R. + M.F.	17:20	—	1000	—	Cervical Tear	Methergin 0.2 mg. I.M.I. Repair of Laceration; Ergometrine 0.5 mg. I.M.I. Blood Transfusion 1000 c.c.	L	

POST-PARTUM HAEMORRHAGE (Continued)

Reg. No.	Age	Previous Pregnancies Before After 28 wks., 28 wks.	Maturity	Method of Delivery	Duration of Labour	Amount of Haemorrhage (c.c.)		Cause or predisposing factor	Treatment State drug, dose and routine	Result	Remarks	
						Before delivery of	After delivery of					
841	21	—	?	Spont.	5:48	1600	—	Retained Placenta	I.V. Pitocin drip 5 units Blood transfusion 500 c.c.	L	L	Retained Placenta, Manual Removal of Placenta done in Eastern Maternity Home, P.P.H. 55 ozs. before Admission
1907	42	—	9	"	?	—	650	Unknown	Ergometrine 0.5 mg. I.M.I.	L	Not Adm.	B.B.A. Forceps Delivery outside admitted in severe shock
3834	26	1	?	—	—	—	—	—	Blood Transfusion 850 c.c.	D	L	Blood loss during C.S.
7818	30	—	4	42-3	—	—	500	Placenta Praevia	—	L	L	
7961	26	—	2	39-1	15:53	—	650	Atony	Methergin 0.2 mg. I.M.I. Methergin 0.2 mg. I.V.I.	L	L	
8148	35	—	3	40-2	19:55	—	1200	Secondary Uterine Inertia	Ergometrine 0.5 mg. I.M.I. Blood Transfusion 1000 c.c.	L	L	

NON-BOOKED

TABLE 25
MANUAL REMOVAL OF PLACENTA

There were 99 cases in which manual removal of the placenta was carried out, 97 Booked and 2 Non-booked.
There were no maternal deaths.
Morbidity was noted in 19 cases.

Reg. No.	Age	Pregnancies Before 28 wks.,	Maturity	Method of Delivery	Length of Labour			Indication	Amount of Bleeding c.c.	Morbidity Days	Result	Remarks
					1st Stage	2nd Stage	3rd Stage					
BOOKED												
181	22	—	42	Spont.	38:08	0:35	1:00	Retained Placenta	80	—	L	Intra-partum Eclampsia
198	27	2	30-3	"	24:00	0:10	1:10	"	80	—	L	
222	31	4	?	"	1:50	0:20	1:30	Partially Adherent	400	2	L	N.D.
378	29	1	38-6	"	12:00	0:10	1:40	Placenta Retained	300	—	L	L
612	36	5	37-3	"	5:30	0:05	1:25	Retained Placenta	100	—	L	L
640	38	10	42-6	"	20:05	0:05	1:35	Adherent Placenta	300	—	L	L
660	39	1	33-3	Bipolar "	5:40		0:35	Adherent Placenta	30	—	L	S.B.
663	44	10	39	Spont.	22:45	0:25	1:25	Retained Placenta	—	—	L	L
697	23	—	40-2	"	15:30	2:00	1:00	"	50	—	L	L
718	32	3	38-3	"	4:20	0:10	2:15	"	150	—	L	L
836	32	2	39-6	"	7:00	0:20	0:05	"	20	—	L	S.R.
955	25	—	38-1	"	4:10	0:25	1:15	"	200	—	L	L
955	22	—	43	M.R. + M.F.	15:15	2:25	0:22	"	100	—	L	L
963	36	—	39-3	Spont.	10:25	0:50	1:30	Adherent Placenta Retained	200	—	L	L
999	25	2	38-1	M.R. + L.F.	6:30	0:01	1:00	"	400	—	L	L
1001	23	—	37-6	"	15:40	2:20	0:35	"	100	—	L	L
1098	25	1	3	Spont.	5:30	2:20	1:35	"	700	—	L	L
1328	25	3	37-5	"	3:45	0:05	1:25	"	700	—	L	L
1361	18	—	33-2	"	7:40	0:15	1:20	"	100	—	L	L
1427	24	—	40-1	"	12:20	0:05	1:05	"	300	5	L	L
1540	19	—	40-5	"	18:10	0:25	1:10	"	650	7	L	L
1582	24	1	35	"	16:35	0:15	1:30	"	250	—	L	L
1716	40	1	41	M.R. + M.F.	11:45	1:45	1:15	"	500	—	L	L
1806	29	1	40-6	Spont.	5:45	1:15	1:10	"	50	—	L	L
1835	29	2	43-4	"	7:35	0:10	1:15	"	60	—	L	L
1851	23	4	29-6	"	6:20	0:10	0:30	"	540	—	L	S.B.
1873	27	1	42-5	M.R. + M.F.	5:50	0:45	1:05	"	250	—	L	L
1899	27	1	31-1	Spont.	3:40	0:15	1:00	"	60	—	L	L
2126	28	2	40-3	"	19:40	0:15	0:50	"	—	—	L	L

P.P.S.
Mitral Stenosis & Aortic Re-
gurgitation Class IV
P.E.
P.L.
P.F.
Filaria
Endometritis
Previous L.S.C.S.
Breech, Hydrops Foetalis
Severe P.E.
Breech

(Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	Maturity 28 wks.	Method of Delivery	Length of Labour			Indication	Amount of Bleeding c.c.	Morbidity Days	Result		Remarks
					1st Stage	2nd Stage	3rd Stage				M.	C.	
2137	20	—	40 6	Spont.	5:00	0:05	0:50	Retained Placenta	500	—	L		
2154	26	—	40-1	"	6:30	0:20	1:52	"	300	—	L		
2254	19	—	38	"	5:10	0:15	1:20	"	200	—	L		
2334	31	—	39-5	"	8:15	0:20	1:30	"	100	—	L		
2376	24	—	35	"	4:35	1:00	1:15	"	100	—	L	Breech	
2378	31	—	30 6	"	2:20	0:05	1:15	"	100	—	L		
2495	20	1	44	"	10:30	0:10	1:30	"	60	1	L		
2505	34	1	33-5	"	3:10	0:15	1:30	"	40	—	L	Bivovular Twin Pregnancy	
2588	22	—	39-3	"	34:30	0:35	1:15	"	60	1	L		
2675	27	—	36-2	"	11:45	0:15	1:10	"	50	—	L		
2688	25	—	39	"	17:00	0:45	1:15	Adherent Placenta	200	—	L	P.E. Uniovular Twin Pregnancy	
2771	42	—	35-5	"	0:30	0:10	0:15	"	1600	1	L	2nd Twin Breech Blood Transfusion 1,500 c.c.	
2806	39	—	32 4	"	9:10	0:05	1:45	Retained Placenta	240	1	L	Breech, P.E.	
2841	42	3	40 6	"	13:20	0:15	0:15	Adherent Placenta	150	—	L		
2845	25	—	39	"	3:05	0:10	1:15	"	30	—	L		
2860	29	—	39-3	"	10:50	0:05	1:45	"	600	2	L		
2999	35	—	40 4	"	28:45	0:05	1:25	"	700	1	L		
3049	34	2	41	M.R. + M.F.	73:10	2:00	1:20	"	600	1	L		
3062	28	—	40-5	Spont.	11:15	0:20	1:55	Retained Placenta	150	—	L		
3469	29	—	42 3	"	5:02	0:03	1:25	Adherent Placenta	200	—	L		
3482	25	—	35	"	6:25	0:05	1:00	"	100	—	L		
3512	26	—	39-4	"	6:50	0:30	0:30	"	300	—	L		
3540	36	—	36-4	"	5:00	0:20	1:10	"	100	—	L		
3541	27	—	39-5	"	6:10	0:20	1:15	"	200	—	L		
3601	32	—	37-5	"	10:25	0:05	1:55	"	150	—	L		
3722	21	—	39-3	"	7:45	0:55	1:25	"	200	—	L		
3933	40	—	43	"	3:12	0:02	1:10	"	60	—	L		
3951	27	—	41-1	"	6:35	0:05	1:20	"	60	—	L		
3965	32	2	37-2	"	22:00	0:05	1:25	"	400	—	L		
3993	28	—	40 5	"	2:25	0:10	1:45	"	150	—	L		
4025	35	—	37-6	"	3:00	0:15	1:10	"	200	—	L		
4213	24	—	41-4	"	3:45	0:05	1:35	"	100	—	L		
4228	29	—	43 4	"	6:10	0:30	1:00	"	200	—	L		
4332	27	1	39 2	"	2:45	0:05	1:35	Adherent Placenta	800	—	L		
4451	25	—	39-3	"	35:01	1:30	1:05	Retained Placenta	50	1	L		
4622	31	—	40 6	"	24:15	0:05	2:00	Adherent Placenta	200	—	L		
4797	27	—	39 5	(1) Mid. Forceps (2) Internal Version	18:25	4:05	2:45	Retained Placenta	300	1	L	Breech, Hydrois 1 oculus	
4819	24	—	38-5	Spont.	6:25	0:45	1:00	Adherent Placenta	200	—	L		
4958	26	—	41	Mid forceps	10:10	2:20	1:45	"	450	—	L		

MANUAL REMOVAL OF PLACENTA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Maturity	Method of Delivery	Length of Labour			Indication	Amount of Bleeding c.c.	Morbidity Days	Result	Remarks
					1st Stage	2nd Stage	3rd Stage					
4998	23	—	38-5	Spont.	3:05	0:05	1:00	Adherent Placenta	400	—	L.	
5288	31	—	35-4	"	6:05	0:05	1:30	"	200	2	L.	
5388	32	—	39-1	"	7:55	0:10	1:50	Retained Placenta	700	—	L.	
5370	32	—	44-3	(1) Mid Forceps (2) Internal Version	94:30	0:15	0:15	"	700 800	1	L.	Uniovular Twin Pregnancy Eclampsia—1 Fit.
5421	30	1	38-1	"	11:40	1:10	1:50	Adherent Placenta	300	—	L.	
5472	27	1	42	Spont.	6:25	0:10	1:40	Retained Placenta	780	1	L.	
5494	30	—	39-2	"	11:45	2:30	0:40	Adherent Placenta	200	—	L.	
5498	24	—	38-6	Low Forceps	3:35	0:15	1:25	"	400	—	L.	One Pint Blood Given
5607	30	—	38	"	1:25	0:05	1:35	"	120	—	L.	
5697	27	—	42-2	"	13:30	0:05	1:25	"	180	—	L.	
5694	23	—	40-4	"	3:00	0:40	1:30	"	150	—	L.	
5942	40	—	40-3	"	4:00	0:40	0:40	"	1,000	1	L.	One Pint Blood Given
6210	26	—	38-1	"	7:00	0:35	1:10	"	150	—	L.	
6408	23	—	39-4	"	10:30	0:40	1:20	"	400	—	L.	
6463	24	—	27-4	"	5:50	0:25	1:00	"	800	—	L.	
6492	26	—	39-1	(1) Mid Forceps (2) Internal Version	12:40	1:45	0:48	"	200	—	L.	Bimovular Twin Pregnancy
6709	30	—	?	"	11:30	0:15	0:55	"	500	—	L.	
6875	25	—	43-6	Spont.	2:50	0:10	1:30	"	250	—	L.	
7011	26	—	39-2	M.R. + M.F.	19:15	1:30	0:10	"	450	—	L.	
7413	36	—	38-4	"	8:50	0:05	1:10	"	400	1	L.	Pulmonary T.B. Previous C.S.
7444	26	1	41-5	Spont.	4:00	0:20	0:20	"	1,500	—	L.	Cellulitis Left Forearm
7655	38	3	39-4	"	6:15	0:03	0:45	"	400	—	L.	Cellulitis Left Forearm
7701	26	—	41-4	"	24:20	1:30	1:30	"	350	—	L.	
7807	24	—	39-2	"	24:00	1:20	1:45	"	200	—	L.	
7928	38	3	35-2	"	22:35	0:10	1:45	"	550	—	L.	
8053	25	—	40-5	Mid Forceps	7:10	0:25	0:20	Retained Placenta	500	—	L.	P.F. Vaginal Septum
8056	24	—	40	"	7:00	0:30	0:25	Adherent Placenta	850	—	L.	
8146	20	—	40-2	Spont.	6:30	0:50	1:40	"	130	—	L.	
8301	30	—	37-4	"	10:05	1:00	1:30	"	300	—	L.	
NON-BOOKED												
913	36	—	42-5	M.R. + M.F.	9:15	4:15	0:15	Retained Placenta	300	—	L.	Anæmia
4139	27	—	38-4	Spont.	14:55	0:05	1:40	"	—	—	L.	

TABLE 26

EXTERNAL VERSION BEFORE LABOUR

This table includes those cases in which version was successfully performed in the Antenatal clinic. External cephalic version was attempted on 49 cases; in 3 patients version was attempted on more than one occasion. The indications for external cephalic version were:—breech presentation—35 cases; transverse and oblique lies—14 cases. There were 28 cases of successful version in the series. Of the 21 cases in whom version was unsuccessful, 11 cases were subsequently delivered as vertex—spontaneous version.

No anaesthesia was used in any of the cases.

There were no maternal deaths, stillbirths or neo-natal deaths.

Reg. No.	Age	Pregnancies Before 28 wks.	After 28 wks.	Maturity At Version	Indication Flexed breech Extended breech Transverse lie	Position of legs confirmed by X-ray Yes, No	Anaesthetic employed	Presentation at Delivery	Result M. C.	Remarks.
14	30	—	2	32.5	Flexed Breech	No	..	R.O.A.	L	Version Failed. Spontaneous Version Later.
29	38	3	5	32.5	"	"	..	L.O.A.	L	Forceps Delivery.
42	29	—	6	29.6	"	"	..	R.O.A.	L	"
55	38	1	5	36.3	"	"	..	R.O.A.	L	"
95	36	—	4	41	Transverse Lie	Yes	..	R.O.A.	L	"
99	33	2	5	36.1	Flexed Breech	No	..	R.O.A.	L	"
173	32	—	3	36.3	"	"	..	L.O.A.	L	"
178	40	2	12	31.4	"	"	..	L.O.A.	L	"
234	27	—	2	34.2	"	"	..	L.O.A.	L	"
243	36	—	2	31.2	"	"	..	R.O.A.	L	"
294	25	—	1	34.5	"	"	..	R.O.A.	L	"
365	32	—	3	37.1	"	"	..	R.O.A.	L	"
365	36	—	4	24.3	Transverse Lie	"	..	L.O.A.	L	"
415	28	1	3	34.1	Flexed Breech	"	..	L.O.A.	L	"
425	27	—	3	27	Extended Breech	Yes	..	L.S.A.	L	"
448	36	1	—	36.3	Flexed Breech	Yes	..	R.S.P.	L	"
501	31	1	4	28	"	No	..	L.S.P.	L	"
510	35	—	7	34.4	"	"	..	R.O.A.	L	"
517	24	—	—	?	Transverse Lie	"	..	R.O.A.	L	"
577	23	—	1	26.6	Flexed Breech	Yes	..	L.O.A.	L	Version Failed. Spontaneous Version Later.
579	19	—	—	34.6	"	"	..	L.S.A.	L	Version Failed. Twic.
598	24	—	—	32.6	"	"	..	L.S.A.	L	Version Failed.
607	30	—	5	29	"	"	..	R.O.A.	L	Version Failed. Spontaneous Version Later.
751	31	1	5	33.5	"	No	..	L.O.A.	L	"

BOOKED

EXTERNAL VERSION BEFORE LABOUR (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	Maturity At Version	Indication Flexed breech Extended breech Transverse lic	Position of legs confirmed by X-ray		Anaesthetic if employed	Presentation at Delivery		Result	Remarks.
					Yes	No		M.	C.		
776	25	—	33-6	Flexed Breech	Yes	—	—	R.S.P.	L	L	Version Failed.
791	29	—	37-5	"	"	—	—	R.S.A.	L	L	"
808	26	—	26-5	"	No	—	—	R.O.A.	L	L	"
828	23	—	35-5	"	Yes	—	—	R.O.A.	L	L	Version Failed.
867	34	1	36-2	"	No	—	—	L.O.A.	L	L	Version Failed.
889	36	2	30-5	"	"	—	—	L.O.A.	L	L	P.P.S.
902	27	2	36-5	Transverse Lic	Yes	—	—	L.O.A.	L	L	Version Failed.
961	26	3	31-6	Flexed Breech	No	—	—	R.O.A.	L	L	Version Later.
988	29	1	33-3	Transverse Lic	"	—	—	R.O.A.	L	L	"
1152	38	8	30-6	"	"	—	—	L.O.A.	L	L	Version Failed.
1272	29	4	?	Flexed Breech	Yes	—	—	R.O.A.	L	L	"
1494	29	1	?	Oblique Lic	"	—	—	R.S.A.	L	L	"
1696	28	2	37-9	"	Yes	—	—	R.S.A.	L	L	"
1942	31	1	31-3	Flexed Breech	No	—	—	R.O.A.	L	L	"
2264	25	—	36-2	"	Yes	—	—	L.O.A.	L	L	"
2676	23	—	33-2	"	"	—	—	L.O.A.	L	L	"
2788	38	1	34-1	Transverse Lic	No	—	—	L.O.A.	L	L	"
3112	32	—	33	"	Yes	—	—	R.O.A.	L	L	"
3186	28	3	35	Flexed Breech	"	—	—	R.O.A.	L	L	"
3272	30	1	37	Extended Breech	"	—	—	R.S.A.	L	L	"
3508	32	4	33-3	Transverse Lic	No	—	—	R.O.A.	L	L	"
3631	23	2	32-2	"	"	—	—	L.O.A.	L	L	"
3994	36	4	31	"	"	—	—	R.O.A.	L	L	"
4435	23	—	28-5	"	"	—	—	R.O.P.	L	L	"
4508	32	3	35-3	Flexed Breech	Yes	—	—	L.O.A.	L	L	"

TABLE 27

INDUCTION OF LABOUR

There were 91 cases of induction of labour, 81 Booked and 10 Non-booked; 31 in primiparae and 60 in multiparae; 86 cases were successfully induced and 5 failed.

There were no maternal deaths.

There were two cases of induction in multiple pregnancy.

Reg. No.	Age	Previous Pregnancies Before 28 wks.	After 28 wks.	Maturity	Indication	Method of Induction	Induction Delivery Interval	Method of Delivery	Duration of Labour 1st Stage 2nd Stage	Weight of Child gm.	Result M. C.	Remarks
BOOKED												
6787/55	24	—	—	44-1	P.E. Postmaturity	O.B.E.	—	—	—	—	L	Failed
407/56	39	—	3	30-1	P.E. Mixed Accidental Haemorrhage	O.B.E. Pitocin	20:45	Bipolar Version Craniotomy	27:50	1013	L S.B.	
1261	33	1	3	36-5	P.E.	A.R.M.	22:10	Spont.	0:25	2150	L	
1455	33	—	1	38-6	"	D-S Catheter	17:55	"	1:15	2480	L	
1504	35	1	3	35	Eclampsia Accidental	"	16:35	"	15:18	1990	L	
1572	40	—	3	42-1	Haemorrhage T.R. Pleural	O.B.E.	7:15	"	2:14	3188	L	
2087	23	—	1	40-5	Overdist. & Large Baby	Stripping of Membranes	8:55	"	8:00	4080	L	Cystitis
2224	36	1	3	41	Mitral Stenosis	D-S Catheter	24:05	"	23:50	3884	L	
2230	30	1	2	42-5	Postmaturity	D-S Catheter	4:00	L.S.C.S.	14:35	3200	L	
2286	22	—	1	32-6	P.E.	" A.R.M.	5:40	Spont.	5:20	2370	L S.B.	
2418	34	—	2	39-4	Pulmonary T.B.	O.B.E.	2:55	"	2:25	3175	L	
2451	31	—	3	39-1	P.E.	"	17:00	"	3:00	2665	L	
2520	29	1	1	38-1	Breech Presentation	D-S Catheter	—	L.S.C.S.	—	3350	L	Failed
2541	42	2	6	36-4	Bronchial Asthma Accidental	A.R.M.	7:00	Spont.	14:55	2645	L	
2543	28	1	2	42-5	Haemorrhage Postmaturity	O.B.E.	3:20	"	6:20	3705	L	
2705	31	—	6	42-2	Placenta Praevia Type 1	D-S Catheter Stripping of Membranes	8:55	"	8:00	4080	L	Cyrtus
2711	37	—	—	45-6	Postmaturity Postmaturity	Stripping of Membranes O.B.E.	—	L.S.C.S.	—	3500	L	Both Failed

INDUCTION OF LABOUR (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	Previous Pregnancies After 28 wks.	Indication	Method of Induction	Induction Delivery Interval	Method of Delivery	Duration of 1st Stage	Duration of Labour 2nd Stage	Weight of Child gm.	Result M. C.	Remarks
3053	38	—	7	35-3 P.E.	Stripping of Membranes	29:25	Spont.	4:00	0:05	1900	L L	
3186	28	—	3	39-5 Contracted Pelvis	Stripping of Membranes	10:15	"	9:00	1:15	3340	L L	
3515	38	2	3	41 Overdue	O.B.E.	1:55	"	1:55	0:05	3175	L L	
3537	42	—	5	37 P.E. (Severe)	Stripping of Membranes	7:45	"	4:27	0:03	2640	L L	
3572	39	—	3	38-5 " "	"	5:00	"	5:25	0:05	3317	L L	
3600	31	—	5	38-5 " P.E.	"	14:25	"	3:40	0:05	3110	L L	
3675	23	—	—	39-2 Aortic Aneurysm	"	9:20	"	8:40	1:00	2480	L L	
3691	26	1	1	40 Bleeding Piles	"	4:30	"	4:00	0:05	3180	L L	
3906	29	—	3	43-1 Postmaturity	Stripping of Membranes	4:35	"	13:00	1:35	4030	L L	
3958	35	1	5	42 D-S Catheter	O.B.E.	17:00	"	11:30	0:10	3600	L L	Stripping of Membranes
4092	23	—	—	40-2 Pulmonary T.B.	D-S Catheter	17:00	"	16:00	0:30	2050	L S.B	O.B.E., Both Failed
4213	24	—	1	41-4 " "	O.B.E. Stripping of Membranes	18:20	"	3:45	0:05	3580	L L	
4243	26	—	2	41-4 Overdue	O.B.E. Stripping of Membranes	—	L.S.C.S.	—	—	2725	L L	
4349	26	—	—	43-5 Postmaturity	D-S Catheter	41:40	Spont.	39:30	0:30	3150	L L	
4389	31	—	4	41-2 P.E.	O.B.E.	2:15	"	2:10	0:05	3200	L L	
4506	22	2	—	40-5 P.E.	Stripping of Membranes	9:00	"	6:45	0:30	2675	L L	
4531	27	—	4	36-2 Accidental Haemorrhage	A.R.M.	5:05	"	8:45	0:05	2920	L L	
4598	26	—	2	? P.T.	Stripping of Membranes	7:10	Spont.	1:35	0:05	3100	L L	
4739	26	—	2	38-4 Hypertension	O.B.E.	20:59	"	10:35	0:20	3350	L L	
5362	31	—	3	? Essential Hypertension	Stripping of Membranes	21:45	"	8:40	0:25	2800	L L	
5393	21	—	—	37 P.E.	A.R.M.	8:00	Low Forceps	4:45	2:15	3170	L L	
5650	28	—	2	34-3 " "	Stripping of Membranes	9:20	Spont.	4:45	(1) 0:05	(1) 2400	L N.D.	
5692	38	1	3	39 " "	"	9:45	Spont.	7:05	(2) 0:20	(2) 2100	L L	Twin Pregnancy
5701	24	—	1	38-4 " "	"	24:00	"	5:10	0:15	3317	L L	
										3940	L L	

INDUCTION OF LABOUR (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	After 28 wks.	Maturity	Indication	Method of Induction	Induction Delivery Interval	Method of Delivery	Duration of 1st Stage	Duration of Labour 2nd Stage	Weight of Child gm.	Result M. C.	Remarks
5803	25	—	1	39-3	Mitral Stenosis & Aortic Regurgitation	Stripping of Membranes	12:45	Spont.	13:30	0:25	3380	L	
5986	27	—	—	42	P.E.	"	30:00	"	14:00	0:15	2950	L	
6022	20	—	—	39-4	"	"	9:00	"	7:00	1:15	2948	L	
6052	21	—	—	35-1	Mitral Stenosis	"	8:55	"	5:00	0:30	2900	L	
6067	22	—	—	39	P.E.	"	61:30	"	6:50	0:30	3050	L	
6080	32	2	2	40-3	P.E.	"	8:15	"	6:00	0:05	3330	L	
6108	34	—	—	39-6	"	A.R.M.	15:35	"	4:45	0:25	3370	L	
6100	24	—	—	43-2	"	Stripping of Membranes	33:35	"	10:20	1:00	2500	L	
6164	31	—	—	41-4	Postmaternity	"	13:05	"	2:30	0:20	2950	L	
6218	24	—	—	37-6	Mitral Stenosis	"	6:35	"	4:35	0:35	2610	L	
6357	21	—	—	38-4	Elective	A.R.M.	3:55	"	3:30	0:25	3400	L	
6493	36	—	—	40-4	P.E.	Stripping of Membranes	24:45	Mid-Forceps	16:30	1:15	3550	L	
6498	28	1	1	40-1	Mitral Stenosis	"	4:30	Spont.	3:00	0:05	3000	L	
6501	25	—	—	38-6	Elective	"	78:00	"	72:15	0:15	2950	L	
6596	25	—	—	42-3	P.E.	"	24:45	"	10:15	0:35	3220	L	
6610	39	—	—	38	"	"	30:25	"	7:30	0:25	4030	L	
6644	40	—	—	40	"	"	19:30	Low Forceps	8:20	2:10	4040	L	
6673	23	—	—	44-1	"	"	1:00	Spont.	7:00	0:45	3115	L	
6791	37	—	—	38-5	"	"	9:35	"	4:50	0:05	4020	L	
6827	32	—	—	41-1	Elective	"	13:35	"	9:25	0:05	2680	L	
6844	32	—	—	41-1	"	"	29:30	"	10:30	1:20	2948	L	
6850	43	—	—	38-2	Contracted Pelvis	"	15:50	"	4:20	0:10	2500	L	
6868	36	—	—	38-4	Elective	"	15:30	"	18:30	0:20	3400	L	
6907	37	2	4	38-5	"	"	8:15	"	19:55	0:05	2800	L	
6902	23	—	—	41	Elective	"	14:30	"	6:00	0:15	2300	L	
6968	22	—	—	38-3	P.E.	"	3:05	Spont.	2:00	0:45	4200	L	
7024	23	—	—	39-5	Contracted Pelvis	"	19:50	Mid Forceps	29:30	0:30	2778	L	
7087	26	—	—	35-5	P.E.	"	6:50	Spont.	3:30	0:30	4062	L	
7156	23	1	1	38-1	Elective	A.R.M.	7:30	"	3:30	0:50	4500	L	
7184	32	—	—	37-2	P.E.	"	52:30	(1)	18:00	(1) 0:15	(1) 2230	L	Twin Pregnancy
								(2)		(2) 0:30	(2) 2640	L	
7219	32	1	5	39-3	Elective	Stripping of Membranes	14:40	"	13:00	0:10	3900	L	
7334	33	—	3	41-3	Accidental Haemorrhage	"	10:50	"	2:45	0:05	3040	L	
7346	27	—	—	40	Elective	Stripping of Membranes	105:10	"	10:00	0:40	3180	L	
7395	36	—	—	37	Eclampsia	A.R.M.	9:00	Mid Forceps	19:10	2:25	2750	L	

INDUCTION OF LABOUR (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	Previous Pregnancies After 28 wks.	Maternity	Indication	Method of Induction	Induction Delivery Interval	Method of Delivery	Duration of 1st Stage	Duration of Labour 2nd Stage	Weight of Child gm.	Result M. C.	Remarks
7421	37	—	7	38-2	Elective	A.R.M.	7:35	Spont.	11:20	1:05	3030	L	
7556	20	—	—	37-3	P.E.	Stripping of Membranes	23:20	Mid Forceps	14:20	2:00	2480	L	Mid pelvic Contraction
7621	33	—	6	40-6	Elective	A.R.M.	26:25	Spont.	2:40	0:15	3340	L	
7791	32	—	5	38-6	"	"	6:15	"	18:25	0:05	3100	L	
7843	17	—	—	43-1	"	"	10:40	"	6:30	1:10	4000	L	
8278	24	—	—	40	"	Stripping of Membranes	20:45	"	12:00	0:10	2900	L	
NON-BOOKED													
1232	26	—	—	43	Postmaturity	O.B.E.	—	—	—	—	—	L	Failed
1953	34	—	—	42-6	P.E.	"	21:15	Spont.	12:15	0:45	3680	L	
2182	28	—	1	41	Postmaturity	"	5:15	"	2:10	0:05	3345	L	
3067	46	—	12	?	Overdue Accidental Haemorrhage	A.R.M.	0:30	"	4:00	0:05	1250	L	S.B.
3323	42	7	3	40-6	P.E. Accidental Haemorrhage	"	2:45	"	0:25	0:10	3400	L	S.B.
3358	32	1	6	42	Postmaturity	Stripping of Membranes D-S Catheter	9:40	"	1:50	0:05	3250	L	L
4135	26	—	—	34-6	P.E.	A.R.M.	16:10	"	6:20	0:05	1450	L	S.B.
4566	44	1	12	36-3	P.E.	Stripping of Membranes D-S Catheter	124:45	"	4:25	0:05	2280	L	L
6306	30	—	2	38-3	P.E. (Severe)	Stripping of Membranes	39:15	"	28:40	0:15	2950	L	L
6307	26	—	1	39-1	Elective	"	42:20	"	6:30	0:20	2790	L	L

TABLE 28

CAESAREAN SECTION

Caesarean Section was performed in 174 occasions—in 141 Booked and 33 Non-booked cases; 155 were lower uterine segment sections and 19 classical sections.

There were no maternal deaths.

There were 9 stillbirths, 4 in booked and 5 in Non-booked cases; 18 neo-natal deaths, 14 in Booked and 4 in Non-booked cases.

Caesarean Section was performed on twin pregnancy on 4 occasions, including one for conjoined twins and one for one head double body monster.

Reg. No.	Age Before After Maturity 28 wks. 28 wks.	Previous Pregnancies	Duration of Labour since 1st Stage	No. of hours since rupture of membranes	Indication	Anaesthetic Used	Type of Operation	Peri-Bladder Inj. Inj. Irid. Irid.	If Steri- lized Child	Weight of Child gm.	Weight Result M. C.	Morbidity (Days)	Remarks
BOOKED													
6949/55	33	—	—	—	Placenta Previa III	Spinal	L.S.C.S.	—	—	3370	L L	—	Elective C.S.
1/56	23	—	19:00	—	Contracted Pelvis	"	"	—	—	2350	L L	—	
101	32	—	2:00	0:20	Previous L.S.C.S. Contracted Pelvis	"	"	—	—	—	L L	—	
170	29	—	44:00	18:40	Previous L.S.C.S. Incoordinate Action of Uterus	"	"	—	Yes	2890	L N.D.	—	Failed Trial Labour
					Maternal Distress					3175	L L	—	
196	19	—	22:00	—	Uterine Inertia	"	"	—	—	2580	L L	2	
237	18	—	7:00	6:00	Breech Presentation Contracted Pelvis, Mild	"	"	—	—	2948	L L	1	
277	25	—	61:00	21:00	Pre-clampsia	"	"	—	—	3147	L L	—	
295	36	—	—	—	Primary Uterine Inertia Elderly Primigravida	"	"	—	—	2950	L L	—	Uterine I fibroid, Myometomy, Polycystic, Flective C.S.
330	24	—	—	—	Contracted Pelvis	"	"	—	—	2900	L L	—	Flective C.S.
334	32	—	—	—	"	"	"	—	—	2962	L L	5	Flective C.S. Uterine Fibroid
337	27	—	25:00	—	Cephalo—Pelvic Disproportion	"	"	—	—	2750	L L	—	
346	23	—	—	—	Contracted Pelvis	"	"	—	—	3629	L L	—	Flective C.S.
400	34	—	19:00	1:30	Mid Pelvic Contraction	"	"	—	—	2700	L L	—	
445	29	1	8:00	—	Contracted Pelvis	"	"	—	—	2855	L L	—	
479	25	—	?	—	Previous C.S. Contracted Pelvis	"	"	—	—	3402	L L	—	

CAESAREAN SECTION (Continued)

Reg. No.	Age	Previous Pregnancies Before After 28 wks. 28 wks.	Maturity	Previous C.S.	Duration of Labour	No. of hours since rupture 1st 2nd of membranes	Indication	Anaesthetic Used	Type of Operation	Peri-Bladder Injured	Weight of Sterilised Child gm.	Result M. C.	Morbidity (Days)	Remarks
506	23	—	1	30	—	—	Placenta Previa Type III	Spinal	L.S.C.S.	—	1250	L N.D.	3	
545	34	—	—	39	6:00	—	Contracted Pelvis	"	"	—	2887	L L	—	
878	31	—	—	43-6	21:00	44:40	Postmaturity Foetal Distress, Uterine Inertia	"	"	—	3175	L L	2	
894	24	—	1	?	20:00	—	Previous L.S.C.S. Contracted Pelvis	"	"	—	3000	L L	1	
946	25	—	1	38-6	1:40	—	Contracted Pelvis Breech Presentation	"	"	—	2863	L L	—	
969	32	—	2	39-2	—	—	Previous Cesarean Section Two Previous L.S.C.S.	"	"	—	3062	L L	—	Elective C.S.
1005	18	—	—	36-4	41:00	—	Contracted Pelvis Pre-eclampsia	"	"	—	2980	L L	3	
1089	32	—	—	40-3	4 days	?	Cervical Dystocia Uterine Inertia	Local	"	—	2800	L L	1	
1230	25	—	—	42-2	—	—	Severe Pre-eclampsia Ante-Partum Eclampsia Dermoid Cyst Rt. Ovary	Spinal Pentothal Local	"	—	3170	L L	—	Elective C.S. Right Oophorectomy.
1270	24	—	3	41-3	—	—	Naegle's Pelvis Poor Obs. History	Spinal	"	—	3345	L L	—	
1290	35	—	2	43	—	—	Contracted Pelvis 2 Previous Cesarean Sections, Postmaturity	"	"	—	3317	L L	—	
1410	19	—	—	39-3	20:00	—	Uterine Inertia	"	"	—	3005	L L	—	
1414	27	—	1	38-3	5:00	Leaking	Contracted Pelvis Previous L.S.C.S.	"	"	—	3090	L L	—	
1421	30	—	—	38-1	—	—	Non-descent of Foetal Head Contracted Pelvis	Local and Spinal	"	—	2807	L L	1	Elective C.S.
1541	25	—	—	40-2	53:00	Leaking	Severe Pre-eclampsia Uterine Inertia	Spinal	"	—	3740	L L	7	
1719	22	—	—	40-6	8:00	Leaking	Foetal Distress	"	"	—	2750	L L	—	Failed Trial Labour
1844	37	—	—	38-6	62:00	15:00	Contracted Pelvis Uterine Inertia Foetal Distress	"	"	—	2200	L L	—	
1918	41	—	4	34-4	—	—	Elderly Primigravida Placenta Previa Type III Transverse Presentation	"	"	—	2460	L L	1	
1964	21	—	—	41-2	6:00	—	Contracted Pelvis	"	"	—	3175	L L	—	

CAESAREAN SECTION (Continued)

Reg. No.	Age	Pregnancies Before After 28 wks. 28 wks.	Maturity	Previous C.S.	Duration of Labour	No. of hours since rupture	Indication	Anaesthetic Used	Type of Operation	Peri-Bladder Inj. Inj. Inj.	If Ser. Inj. Used	Wt. Child gm.	Weight Result	Morbidity (Days)	Remarks
					1st Stage	2nd Stage							M. C.		
2091	26	1	2	38-3	---	---	Two Previous Contracted Pelvis T.B. Hip (left)	Spinal	L.S.C.S.	---	Yes	2650	L. I.	---	Elective C.S.
2150	35	---	---	---	---	---	Elderly Primigravida Breech Presentation	"	"	---	---	3147	L. L.	---	"
2212	33	---	---	---	16:50	---	Contracted Pelvis	"	"	---	---	3175	L. L.	---	Failed Trial Labour
2230	30	1	2	42-5	14:35	3:25	Postmaturity Prolonged 2nd Stage Maternal and Foetal Distress	"	"	---	---	3260	L. I.	---	"
2318	27	---	---	---	---	---	Ante-partum Eclampsia	Local	"	---	---	2600	L. L.	---	Elective C.S.
2471	25	---	---	---	---	---	Previous L.S.C.S. Contracted Pelvis	Spinal	"	---	---	2870	L. L.	---	"
2520	29	1	1	38-1	---	---	Maternal Distress	"	"	---	---	3350	L. I.	---	Bronchial Asthma, Induced by Low Rupture of Membranes & Pilo-cin Drip, both Failed.
2531	28	---	3	41	6:00	---	Contracted Pelvis Breech Presentation	"	"	---	---	2875	L. I.	---	"
2657	27	---	2	33-5	---	---	Previous L.S.C.S. Placenta Praevia Type III	Local	"	---	---	2500	L. N.D.	---	"
2711	37	---	---	45-6	46:00	22:00	Elderly Primigravida Postmaturity	Spinal	"	---	---	2500	L. I.	---	"
2807	26	---	2	41	---	---	Uterine Inertia Contracted Pelvis	Local	"	---	---	4111	L. L.	---	Elective C.S.
2864	24	1	2	33	27:30	---	Previous L.S.C.S. Contracted Pelvis	Spinal	"	---	---	2608	L. I.	---	"
2922	23	---	1	40	---	---	Previous L.S.C.S. Contracted Pelvis	"	"	---	---	2722	L. L.	---	"
2945	29	---	---	44-3	---	---	Contracted Pelvis	"	"	---	---	3652	L. L.	---	Elective C.S.
3057	25	---	1	34-3	18:00	---	Uterus Didelphys Previous L.S.C.S.	"	"	---	---	1830	L. L.	---	Vaginal Septum
3086	28	---	4	36-4	---	---	Uterine Inertia Placenta Praevia Type IV	"	"	---	---	2722	L. S.B.	2	Total Blood Loss 800 c.c. Blood Transfusion 1500 c.c.
3109	23	---	---	38-2	8:00	---	Contracted Pelvis	"	"	---	---	3370	L. L.	1	"
3121	20	---	---	40-6	66:20	---	Uterine Inertia	"	"	---	---	3147	L. L.	---	"
3277	28	---	---	41	---	---	"	"	"	---	---	3062	L. L.	---	"
3339	21	---	---	40-4	24:00	---	Severe Pre-eclampsia	"	"	---	---	3100	L. L.	1	Failed Trial Labour
3532	30	---	2	28-6	---	---	Contracted Pelvis Placenta Praevia Type III	"	"	---	---	1420	L. N.D.	---	"
3682	24	---	---	39-1	16:00	---	Leaking Clinical Disproportion	"	"	---	---	3570	L. L.	---	"

CAESAREAN SECTION (Continued)

Reg. No.	Age Before After Maturity 28 wks. 28 wks.	Previous Pregnancies	Duration of Labour	No. of hours since rupture 1st Stage of mem- branes	Previous C.S.	Indication	Anesthetic Used	Type of Operation	Peri- tonium der In- jured	Blad- der In- jured	If Steri- lized Child	Weight of Child gm.	Morbidity (Days)	Remarks	
															M.
3810	28	—	2	40-5	L.S.C.S.	Previous L.S.C.S. Contracted Pelvis	Spinal	L.S.C.S.	—	—	—	3780	L	L	—
3985	21	—	—	34	—	Contracted Pelvis Shoulder Presentation	"	"	—	—	—	2670	L	L	—
3988	28	—	2	40-6	—	Diabetes Mellitus; Postmaturity	"	"	—	—	—	4150	L	L	Elective C.S.
4022	25	—	1	40-2	C.C.S.	Large Baby Contracted Pelvis Previous C.C.S.	"	"	—	—	—	2820	L	L	Elective C.S. Incomplete Rupture of Previous Cesarean Section Scar.
4045	26	—	—	?	—	Breech Presentation Contracted Pelvis	"	"	—	—	—	2725	L	L	—
4163	32	—	—	36	—	Severe Pre-eclampsia	"	"	—	—	—	2720	L	L	Elective C.S.
4191	39	1	—	26-4	—	Placenta Praevia Type IV	Local	"	—	—	Yes	900	L	S.B.	1
4198	34	—	—	39-2	—	Pre-eclampsia; Foetal Distress	Spinal	"	—	—	—	3150	L	L	—
4216	24	—	2	39-4	2 L.S.C.S.	2 Previous L.S.C.S.; Contracted Pelvis	"	"	—	—	Yes	2910	L	L	Elective C.S.
4243	26	—	2	42-6	—	Contracted Pelvis Postmaturity	"	"	—	—	—	2525	L	L	—
4309	31	—	1	38-2	—	Contracted Pelvis	"	"	—	—	—	2510	L	L	—
4321	37	—	4	39-6	C.C.S.	Pathological Retraction Ring; Foetal Distress Contracted Pelvis	"	"	—	—	—	3270	L	S.B.	—
4336	27	—	—	39	—	Congenital Rectovaginal Fistula; Imperforate Anus	"	"	—	—	—	3856	L	L	—
4478	36	—	1	39-4	L.S.C.S.	Contracted Pelvis; Previous L.S.C.S.;	"	"	—	—	—	3200	L	L	—
4497	30	1	—	40-3	—	Breech Presentation Contracted Pelvis	"	"	—	—	—	2830	L	L	1
4500	20	—	—	38-1	—	Uterus Didelphus; Breech Presentation	"	"	—	—	—	2900	L	L	—
4534	27	1	2	38-1	2 L.S.C.S.	2 Previous L.S.C.S.	"	"	—	—	Yes	2900	L	L	Elective C.S.
4655	30	—	—	40-2	—	Leaking Primary Uterine Inertia	"	"	—	—	—	2950	L	L	2
4705	25	1	—	37	—	Contracted Pelvis	"	"	—	—	—	2350	L	L	—
4737	27	1	3	40-1	—	Placenta Praevia Type IV Transverse Lie	"	"	—	—	Yes	2665	L	L	—
4769	29	—	—	38-5	—	Contracted Pelvis; Pre-eclampsia	"	"	—	—	—	3175	L	L	—

Incomplete Rupture of
Uterus; Subtotal
Hysterectomy and Rt.
Salpingo Oophorectomy
P.F.

CAESAREAN SECTION (Continued)

Reg. No.	Age Before 28 wks.	Previous Pregnancies After 28 wks.	Previous C.S.	Duration of Labour		No. of hours since rupture of membranes	Indication	Anesthetic Used	Type of Operation	Peri-Bladder Injured	H. Sterilized	Weight of Child gm.	M. C. Result	Norbidity (Days)	Remarks
				1st Stage	2nd Stage										
5018	28	—	1	?	9:00	—	Previous L.S.C.S.; Foetal Distress	Spinal	L.S.C.S.	—	—	2637	L. I.	—	
5233	26	—	1	1	5:00	—	Contracted Pelvis; Previous C.C.S.	"	"	—	—	2900	L. I.	1	
5251	35	—	—	30-3	5:10	—	Elderly Primigravida Foetal Distress; Pre-eclampsia	"	"	—	—	2495	L. I.	1	
5360	25	—	—	39	—	—	Contracted Pelvis Pre-eclampsia	"	"	—	—	2500	L. I.	1	Ilece C S
5429	30	—	2	39-3	2:30	—	Contracted Pelvis; Previous L.S.C.S.	"	"	—	—	3150	L. I.	—	
5447	30	—	—	38-1	6:00	—	Contracted Pelvis; (Mid and Outlet); Breech Presentation	"	"	—	—	3440	L. I.	1	Incomplete Vaginal Septum.
5547	39	1	1	34-3	21:00	2:00	Previous L.S.C.S.; Prolonged 2nd Stage Without Progress	"	"	—	—	3100	L. I.	2	Occult Rupture of Previous C.S. Scar
5581	32	—	—	41	—	—	Contracted Pelvis; Foetal Distress	"	"	—	—	3150	L. I.	—	
5599	26	—	2	32-5	10:00	—	Previous L.S.C.S.; Contracted Pelvis	"	"	—	Yes	2300	L. N.D.	—	
5621	24	—	—	36	—	—	Severe Pre-eclampsia	"	"	—	—	2495	L. I.	1	D.A.A. Flective C S
5663	40	—	7	?	—	—	Placenta Praevia Type II	Local	C.C.S.	—	—	2570	L. N.D.	1	Flective C.S.
5670	23	—	1	33-5	—	—	Placenta Praevia Type II 2 Previous L.S.C.S. for Contracted Pelvis	Local Spinal	L. S.C.S.	—	—	2150	L. N.D.	5	
5731	29	—	1	40-1	17:00	—	Contracted Pelvis; Previous L.S.C.S.	"	"	—	—	3850	L. I.	—	
5778	20	—	—	36-5	7:30	3:00	Contracted Pelvis; Prolonged 2nd Stage; Breech Presentation	"	"	—	—	4200	L. I.	4	Postpartum Psychosis
5832	29	—	—	39	13:00	6:00	Contracted Pelvis; Breech Presentation	"	"	—	—	3250	L. N.D.	2	
5856	26	—	—	39-2	33:30	3:00	Severe Pre-eclampsia; Primary Uterine Inertia	Local	"	—	—	2800	L. I.	2	
5983	26	—	1	35-5	10:30	—	Previous L.S.C.S.; Contracted Pelvis	Spinal	"	—	—	3180	L. I.	1	
6076	32	1	8	40-1	—	—	Breech Presentation; 2nd Previous S.P. with Severe Pre-eclampsia	"	"	—	—	4000	L. I.	5	Flective C.S.
6133	24	—	—	35	—	—	Breech Delivery	Local	"	—	—	1900	L. I.	1	Flective C.S., Uniovular. Twin Pregnancy.

CAESAREAN SECTION (Continued)

Reg. No.	Age Before After	Previous Pregnancies 28 wks. to 48 wks.	Maturity	Previous C.S.	Duration of Labour 1st Stage	No. of hours since rupture of membranes	Indication	Anæsthetic Used	Type of Operation	Peri-toneum for Inflation	If Sterilised Child	Weight Child gm.	Result		Morbidity (Days)	Remarks
													M.	C.		
6197	32	—	41-2	—	—	—	Pre-eclampsia; Contracted Pelvis	Spinal	L.S.C.S.	—	—	2680	L.	L.	—	Elective C.S.
6251	21	—	39-4	L.S.C.S.	2:00	—	Previous L.S.C.S.	"	"	—	—	4200	L.	L.	2	
6276	36	5	40-5	2 L.S.C.S.	—	—	Naegle's Pelvis; 2 Previous L.S.C.S.	"	"	—	Yes	2977	L.	L.	2	Elective C.S.
6314	22	—	40-5	L.S.C.S.	—	—	Previous C.S.; Contracted Pelvis; Breech Presentation	"	"	—	—	3200	L.	L.	3	Elective C.S.
6381	36	—	38-6	L.S.C.S.	—	—	Previous C.S.; Contracted Pelvis	"	"	—	Yes	3147	L.	L.	—	Elective C.S.
6391	31	2	40-3	—	—	—	Pre-Eclampsia	"	"	—	—	3760	L.	L.	1	
6402	32	4	33-3	—	—	—	Poor Obs. History	"	"	—	—	1620	L.	N.D.	1	Elective C.S.
6428	23	2	38-2	L.S.C.S.	6:30	—	Præcæta Prævia Type II Contracted Pelvis	Local Spinal	"	—	—	2890	L.	L.	1	
6431	21	—	38-3	L.S.C.S.	7:00	—	Shoulder Presentation Contracted Pelvis	"	C.C.S.	—	—	3120	L.	L.	—	
6435	25	—	35-6	2 L.S.C.S.	—	—	Previous L.S.C.S. (1) Placenta Prævia Type IV	Spinal	L.S.C.S.	—	—	1160	L.	N.D.	—	Subtotal Hysterectomy
6487	19	—	39-4	—	10:00	—	2 Previous C.S.; (2) Contracted Pelvis	Spinal	"	—	—	2590	L.	L.	—	
6491	22	—	30-4	—	—	—	Contracted Pelvis	"	"	—	—	3629	L.	L.	—	Elective C.S.
6525	25	—	38-4	—	4:30	1:30	Breech Presentation Prolapsed Cord	"	"	—	—	3590	L.	L.	3	
6538	34	—	39	—	2:30	4:30	Breech Presentation Contracted Pelvis	"	"	—	—	3400	L.	L.	1	
6611	21	—	35-4	—	—	—	Pathological Retraction Ring; Fœtal Distress History of S.B.; Naegle's Pelvis;	"	C.C.S.	—	—	2707	L.	L.	1	Placenta Prævia Type I.
6617	28	—	36-1	—	—	—	Placenta Prævia Type II	"	"	—	—	3317	L.	L.	1	Elective C.S.
6701	23	—	38-1	—	—	—	Contracted Pelvis Ante-partum Felampnia; 10 Convulsions	Local	L.S.C.S.	—	—	2875	L.	L.	1	Elective C.S.
6738	25	—	39-6	—	10:30	—	Contracted Pelvis	Spinal	"	—	—	3400	L.	L.	3	Failed Trial Labour
6781	36	—	38	—	19:30	—	Elderly Primigravida Sterility 11 Years; Uterine Inertia	"	"	—	—	2930	L.	L.	—	Pre Felampnia
6824	25	1	40-3	2 L.S.C.S.	5:30	—	Contracted Pelvis; 2 Previous C.S.	"	"	—	Yes	3570	L.	L.	1	

CAESAREAN SECTION (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Age at 28 wks.	Maturity	Previous C.S.	Duration of Labour	No. of hours since rupture	Indication	Anaesthetic Used	Type of Operation	Peri-toneum injured	Bled-injured	Used	Weight of Child gm.	Result	Morbidity (Days)	Remarks
						1st Stage	2nd Stage										
6861	37	—	—	?	—	—	—	Accidental Haemorrhage, Revealed;	Spinal	C.C.S.	—	—	—	2070	L L	—	Bicornuate Uterus
6867	25	—	2	44.3	—	6:20	0:50	Elderly Primigravida Contracted Pelvis;	"	L.S.C.S.	—	—	—	3572	L L	—	
7026	29	—	4	41	—	—	—	Poor Obs. History Previous Manchester Operation	"	"	—	—	Yes	2722	L L	—	
7082	42	2	5	40.5	—	11:45	4:30	Foetal Distress; Excessive Sized Infant	"	"	—	—	—	4451	L L	—	
7137	34	—	8	40	—	—	—	Placenta Praevia Type IV	"	C.C.S.	—	—	—	2693	L L	—	Elective C.S.
7322	29	—	—	31.2	—	—	—	Severe Pre-eclampsia	Local	"	—	—	—	1446	L N.D.	—	Elective C.S.
7344	34	1	5	39.3	—	4 days	47:00	Uterine Inertia; Foetal Distress	Spinal	L.S.C.S.	—	—	—	2400	L L	—	
7371	40	—	4	40.6	—	80:30	11:00	Foetal Distress	"	"	—	—	—	3070	L L	6	Secondary p.p.H. 1,500 c.c.
7385	26	—	—	40.3	—	24:00	—	Foetal Distress Contracted Pelvis;	Local	"	—	—	—	3050	L L	—	Failed Trial Labour
7419	39	—	—	36.3	—	15:00	10:00	Foetal Distress Elderly Primigravida	Spinal	"	—	—	—	3110	L L	—	Failed Trial Labour
7490	33	—	—	38.3	—	21:00	—	Breech Presentation Kyphosis	Local	C.C.S.	—	—	—	3490	L L	—	Pre-Eclampsia; Old T.B. Spine & Lungs
7550	26	1	1	42.2	—	—	3:45	Foetal Distress Contracted Pelvis; Excessive Sized Infant	"	L.S.C.S.	—	—	—	4120	L L	1	Failed Trial Labour
7562	38	1	1	38.5	L.S.C.S.	20:00	—	Bad Obs. History; Previous L.S.C.S.	Spinal	"	—	—	—	3948	L L	—	
7586	23	1	1	38	L.S.C.S.	12:00	—	Contracted Pelvis; Previous L.S.C.S.	"	"	—	—	—	3100	L L	—	
7592	29	1	1	29	—	—	—	Severe Pre-eclampsia	Local	C.C.S.	—	—	—	1200	L N.D.	—	Elective C.S. (Unsuccessful Accidental Hemorrhage found at Operation)
7606	35	—	8	35.2	—	—	—	Accidental Hemorrhage	Spinal	"	—	—	—	2600	L S.B.	—	Elective C.S.
7630	27	—	3	39.2	2 L.S.C.S.	15:30	—	Mixed Naegele's Pelvis;	"	"	—	—	Yes	2850	L L	—	
7657	33	1	4	36	—	—	—	2 Previous L.S.C.S. Placenta Praevia Type III	"	"	—	—	—	1880	L J	—	
7811	42	2	2	39.5	—	15:00	10:00	Excessive Sized Baby With C.P. Disproportion	"	L.S.C.S.	—	—	—	4600	L L	—	
7845	28	—	1	38.2	—	13:00	3:10	Conjoined Twins With Obstructed Labour	"	"	—	—	—	4500	L N.D.	—	

CAESAREN SECTION (Continued)

Reg. No.	Age Before After 28 wks. 28 wks.	Pregnancies Before After 28 wks. 28 wks.	Maternity	Previous C.S.	Duration of Labour 1st Stage 2nd Stage	No. of hours since rupture of membranes	Indication	Anaesthetic Used	Type of Operation	Peri-Operative Infection	Used Steril-If	Weight of Child gm.	Result M. C.	Morbidity (Days)	Remarks
7870	30	—	1	41-5	—	—	Previous C.S. with Febrile Postoperative Course	Spinal	C.C.S.	—	—	2600	L L	—	Elective C.S.
7944	30	—	1	38-3	L.S.C.S.	9:00	Naegele's Pelvis; Transverse Lie Previous L.S.C.S.	"	"	—	—	3060	L N.D.	2	Hepato splenomegaly, Absence of Right Kidney; Septate Uterus
7964	27	1	3	—	—	—	Placenta Praevia Type III; Local Transverse Lie	Local	L.S.C.S.	—	Yes	2670	L L	3	Elective C.S.
8080	25	—	1	39-5	L.S.C.S.	11:00	Previous L.S.C.S.; Contracted Pelvis	Spinal	"	—	—	2580	L L	—	
8211	35	—	—	39-4	—	68:00	Elderly Primigravida Prolonged Labour	"	"	—	—	2750	L L	—	
NON-BOOKED															
361	21	—	—	40-3	—	50:30	Uterine Inertia	"	"	—	—	3300	L L	1	
537	17	—	—	40-3	—	15:00	Contracted Pelvis;	"	"	—	—	2807	L L	—	
638	22	—	—	41-2	—	—	"	"	"	—	—	3289	L L	3	Elective Caesarean Section
643	20	—	1	44-4	L.S.C.S.	5:00	Previous L.S.C.S.; Foetal Distress	"	"	—	—	2750	L L	—	
909	24	—	—	35-2	—	89:00	Ante-partum Eclampsia; Uterine Inertia	"	"	—	—	1630	L L	—	Placenta Praevia Type I. Uniovular Twin Pregnancy. 1 Fit before Admission.
1054	28	—	—	39-6	—	15:00	Contracted Pelvis T.B., Left Hip Joint	"	"	—	—	2523	L L	—	C.S. Done on admission.
1232	26	—	—	43	—	—	Postmaturity; Failed Medical Induction; Unengaged Head	"	"	—	—	3345	L L	—	Elective C.S.
1276	19	—	—	40-3	—	60:00	Uterine Inertia; Pre-eclampsia; Foetal Distress; Prolonged Labour;	"	"	—	—	2948	L L	—	In Labour for 2 days before Admission.
1558	35	—	2	39-6	L.S.C.S.	—	Naegele's Pelvis; Placenta Praevia Type III	"	"	—	—	3260	L L	—	Blood Loss 500 c.c. Blood Transfusion 1000 c.c.
1829	41	—	11	32-5	—	3:00	Shoulder Presentation; Prolapse of Cord and Arm	"	"	—	Yes	1940	L L	—	
1837	38	—	2	34-3	—	—	Placenta Praevia Type III	Local	L.S.C.S. &	—	—	2050	L L	1	Flective C.S., Subserous Uterine Fibroid
1958	29	2	—	36-1	—	—	Severe Pre-eclampsia	Spinal	Myomectomy	—	—	2520	L L	—	

CAESAREN SECTION (Continued)

Reg. No.	Age Before 28 wks. 28 wks.	Pregnancies Before After 28 wks. 28 wks.	Maturity	Previous C.S.	Duration of Labour		No. of hours since rupture of membranes	Indication	Anaesthetic Used	Type of Operation	Peri- In- junct In- junct	Blad- tonem der In- junct	If Seri- Used	Wt. Child gm.	Result		Morbidity (Days)	Remarks
					1st Stage	2nd Stage									M.	C.		
2473	35	—	40.4	—	24:00	—	Contracted Pelvis; Foetal Distress	Spinal	L.S.C.S.	—	—	—	—	3400	L	L	—	Failed Trial Labour
2566	30	—	32.4	—	—	—	Elderly Primigravida	"	"	—	—	—	—	1720	L	N.D.	—	
2610	28	—	32.4	—	—	—	Placenta Praevia Type IV Proapsed Cord	"	"	—	—	—	—	1840	L	N.D.	—	
2634	32	1	36.2	—	—	—	Placenta Praevia Type II	"	"	—	—	—	—	3289	L	L	—	
3599	29	—	42.6	—	—	—	Placenta Praevia Type III Cephalo-pelvic Disproportion	"	"	—	—	—	—	2700	L	S.B.	—	Monster, Terata Anadiadyna
3746	28	—	39.2	—	7:00	—	Contracted Pelvis	"	"	—	—	—	—	3310	L	L	—	Elective C.S.
3782	32	—	39.5	—	—	—	Placenta Praevia Type III Local Pre-eclampsia	"	"	—	—	—	—	2400	L	S.B.	—	Elective C.S.
3899	21	—	43.5	—	—	—	Postmaturity; Contracted Pelvis	Spinal	"	—	—	—	—	3150	L	L	—	Elective C.S.
4323	23	—	39.5	—	—	—	Contracted Pelvis	"	"	—	—	—	—	3200	L	L	—	"
6311	36	—	32.1	—	—	—	Severe Accidental hemorrhage	Local	C.C.S.	—	—	—	—	1300	L	S.B.	2	"
6649	36	—	40.4	—	—	—	Prolapsd Cord; Cervical Fibroid	Spinal	Norton's extra-peri- toneal C.S.	—	—	—	—	2551	L	Mac.	5	"
6958	35	—	?	—	—	3:00	Obstructing Labour Brow Presentation; Failed Forceps	"	L.S.C.S.	—	—	—	—	3200	L	N.D.	—	Failed Forceps done out side.
7483	38	—	38	—	—	—	Severe Pre-eclampsia	Local	C.C.S.	—	—	—	—	2460	L	L	1	Elective C.S.
7492	21	—	?	—	34:00	—	Shoulder Presentation; Contracted Pelvis	Spinal	L.S.C.S.	—	—	—	—	3770	L	L	—	"
7494	38	—	38.6	—	—	—	Severe Hypertension	Local	C.C.S.	—	—	—	—	3345	L	L	—	Elective C.S.
7818	30	—	42.3	—	—	—	Placenta Praevia Type IV	Spinal	L.S.C.S.	—	—	—	—	4910	L	L	—	Elective C.S.
7955	27	—	42.2	—	48:00	—	Contracted Pelvis; Foetal Distress; Uterine Inertia	"	L.S.C.S.	—	—	—	—	4050	L	L	2	Prolonged Labour 48 hours
7999	41	1	34.5	—	18:00	—	Shoulder Presentation; Proapsed Left Arm; Bandl's Ring	"	C.C.S.	—	—	—	—	1800	L	N.D.	—	Accidental Hemorrhage Revealed
8088	35	—	41	—	—	—	Severe Pre-eclampsia	Local	"	—	—	—	—	2880	L	L	—	Elective C.S.
8112	30	—	36.5	—	—	—	Accidental Hemorrhage, Connected, Severe	"	L.S.C.S.	—	—	—	—	1850	L	L	—	"
8236	35	1	40	—	—	—		Spinal	L.S.C.S.	—	—	—	—	3400	L	S.B.	—	"

TABLE 29

FORCEPS DELIVERIES

There were 199 cases of forceps deliveries, 184 in Booked and 15 in Non-booked cases; 134 in primiparae and 65 in multiparae.

There were no maternal deaths.

There were 4 stillbirths, all Booked; 12 neo-natal deaths, 8 in Booked and 4 in Non-booked cases.

Forceps were applied 6 times in twin pregnancy.

Reg. No.	Age	Previous Pregnancies Before 28 wks.	Maturity	Indication	Duration of Labour 1st Stage	Duration of Labour 2nd Stage	Weight of Child gm.	M.	Result	C.	Remarks
BOOKED											
6787/55	24	—	44-1	P.E.; Prolonged 2nd Stage; D.T.A.	11:15	2:30	4000	L	L	L	O.T. Position. M.R. + M.F.
24/56	26	—	40-5	Prolonged 2nd Stage; D.T.A.	7:10	1:50	3650	L	L	L	M.R. + M.F.
90	24	—	40-2	"	11:30	2:40	2948	L	L	L	"
132	28	—	39-2	Prolonged 2nd Stage; Maternal Distress	12:20	2:55	3289	L	L	L	"
212	38	—	39-1	Prolonged 2nd Stage; Foetal Distress	10:45	1:35	2650	L	L	L	P.E., M.F.
224	25	—	40	Prolonged 2nd Stage; D.T.A.	24:20	2:55	3500	L	L	L	M.R. + M.F.
547	26	—	38-3	"	9:20	3:05	2970	L	L	L	"
579	19	—	41-3	A.C. Head	8:50	0:20	3120	L	L	L	L.F.
598	24	—	38	"	22:30	0:16	2600	L	L	L	"
652	36	—	40-5	Prolonged 2nd Stage; D.T.A.	8:00	1:55	3850	L	L	L	M.R. + M.T.
700	26	—	36	"	7:45	3:00	3212	L	L	L	L.F.
738	28	—	38-3	Prolonged 2nd Stage; Foetal Distress	19:40	2:10	2750	L	L	L	M.R. + M.F.
825	28	—	40	P.E.; Maternal and Foetal Distress	41:40	0:45	2985	L	L	L	"
835	23	—	41-3	Prolonged 2nd Stage; P.E.	12:20	2:20	3420	L	L	L	L.F.
868	29	—	40-3	Maternal and Foetal Distress	47:40	4:30	2980	L	L	L	"
881	23	—	45-5	Intra-partum Eclampsia	6:15	0:11	2960	L	L	N.D.	M.R. + M.F.
938	28	—	40-5	Prolonged 2nd Stage; Foetal Distress	65:50	2:05	4430	L	L	L	"
995	22	—	43	Prolonged 2nd Stage; P.E.	15:15	2:25	3160	L	L	L	M.R. + M.F.
1001	23	—	42-1	Prolonged 2nd Stage	15:14	2:20	2850	L	L	L	M.R. + M.F.
1029	24	—	42-1	Prolonged 2nd Stage; Foetal Distress	58:25	2:40	4000	L	L	L	"
1045	28	—	39-3	"	26:15	2:25	2665	L	L	L	L.F.
1061	30	—	39-5	Foetal Distress	18:00	0:55	3340	L	L	L	M.R. + M.F.
1082	34	—	40-2	"	7:00	7:00	3864	L	L	L	P.P.H.; M.R. + M.F.
1083	36	—	35-3	Prolonged 2nd Stage; D.T.A.	31:35	2:50	2600	L	L	L	M.R. + M.F.
1128	25	—	38-2	Prolonged 2nd Stage	2:55	2:50	5900	L	L	L	L.F.
1153	37	—	32-4	Prolonged 2nd Stage; D.T.A.;	13:30	2:05	5255	L	L	L	M.R. + M.F.
1324	27	—	37-5	Foetal Distress	24:40	2:40	2651	L	L	L	M.R. + M.F.

FORCEPS DELIVERIES (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	After 28 wks.	Maturity	Indication	Duration of 1st Stage	Duration of Labour 2nd Stage	Weight of Child gm.	M.	Result C.	Remarks
1339	25	—	2	3	Prolonged 2nd Stage; D.T.A.	13:40	1:35	3830	L	L	M.R. + M.F.
1511	21	—	—	45	Prolonged 2nd Stage	11:05	2:00	2892	L	L	M.F.
1575	26	—	—	42-5	Prolonged 2nd Stage; Foetal Distress	6:15	4:00	2800	L	L	1st. Twin; M.F.
1677	28	—	—	39	Prolonged 2nd Stage; D.T.A.	68:25	3:05	3800	L	L	M.R. + M.F.
1684	23	—	—	44	Prolonged 2nd Stage	12:15	2:50	3820	L	L	M.F.
1716	40	1	9	41	Prolonged 2nd Stage; D.T.A.	11:45	1:50	4420	L	L	M.R. + M.F.
1737	31	—	—	40	Prolonged 2nd Stage; Foetal Distress; Uterine Inertia	10:30	2:03	3560	L	L	M.F.
1769	27	—	—	42	Maternal Exhaustion	11:50	0:29	3175	L	S.B.	L.J.
1784	31	1	3	39	Prolonged 2nd Stage	15:45	1:05	3300	L	L	"
1790	24	—	—	40.6	Prolonged 2nd Stage; Foetal Distress	34:20	2:30	3270	L	L	"
1873	27	—	—	42-5	Severe P.E.	5:50	0:45	3005	L	L	M.R. + M.F.
1954	40	2	5	40	Prolonged 2nd Stage;	24:55	1:45	3050	L	L	"
1993	40	2	6	42-1	Prolonged 2nd Stage;	6:30	1:40	4800	L	L	M.F. Hypothyroidism
2002	38	3	6	42-2	Foetal Distress; P.E.	5:30	1:45	3520	L	L	M.R. + M.F.
2014	24	—	—	40-3	Prolonged 2nd Stage;	17:00	2:20	2948	L	L	M.J.
2016	52	—	—	41-6	Foetal Distress	65:45	0:40	2950	L	L	"
2068	27	—	—	39-5	Foetal Distress;	66:30	0:20	2590	L	L	L.J.
2075	27	—	—	41-2	Maternal Exhaustion	21:00	2:15	3090	L	N.D.	M.R. + M.F.
2085	23	—	—	37-5	Prolonged 2nd Stage	15:40	3:05	3220	L	L	L.F.
2088	36	—	2	39-1	Prolonged 2nd Stage; Foetal Distress	8:10	1:55	3200	L	L	M.R. + M.F.
2092	21	—	—	46	Prolonged 2nd Stage; P.E.	5:10	2:55	3200	L	L	"
2144	21	—	—	40-3	Prolonged 2nd Stage;	29:30	2:55	3300	L	L	"
2153	31	—	—	40-4	Foetal Distress;	27:35	0:55	3100	L	L	M.I.
2198	25	—	—	40-3	Secondary Uterine Inertia	15:45	1:00	2700	L	L	"
2202	24	—	—	39	Foetal Distress	8:00	1:45	3300	L	L	M.R. + M.F.
2207	28	—	—	39-5	"	7:43	0:25	2608	L	L	"
2221	22	—	—	36-1	"			2650	L	L	"
2268	30	—	—	40-1	Foetal Distress;	60:50	1:35	3780	L	S.B.	M.F.
2332	27	—	5	40-1	Mixed Accidental Hemorrhage	28:05	0:30	2332	L	L	L.F.
2333	20	—	—	39-3	Aberrant Edemapsia	18:00	1:45	2900	L	L	"
2333	20	—	—	43	Maternal Distress;				L	L	"
2356	27	—	—	39-2	Foetal Distress	29:40	2:25	3409	L	L	M.J.
2463	29	—	—	43-1	Prolonged 2nd Stage	6:40	1:45	2810	L	L	L.F.
2542	36	—	—	37-2	Prolonged 2nd Stage	16:50	1:20	3060	L	N.D.	M.R. + M.F. 2nd Twin
2671	33	—	3	40	Prolonged 2nd Stage;	16:50	1:30	3450	L	L	M.R. + M.F.
2827	36	—	—	42-2	Foetal Distress; D.T.A. Elderly Primigravida	17:05	2:10	2693	L	L	M.F.

FORCEPS DELIVERIES (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Pregnancies After 28 wks.	Maturity	Indication	Duration of 1st Stage	Duration of Labour 2nd Stage	Weight of Child	Result	Remarks
2811	34	2	—	42-1	Prolonged 2nd Stage;	23:25	2:55	3900	L	M.F.
2907	27	1	—	42.5	Foetal and Maternal Distress	18:55	0:20	2840	L	L.F. Previous Hysterotomy
2942	38	—	2	37-3	Foetal and Maternal Distress	4:40	0:25	3280	L	M.F.
2946	29	2	—	42	Mitral Stenosis; P.E.	27:37	2:53	3250	L	M.F. + M.F.
2954	23	—	—	41-3	Prolonged 2nd Stage;	29:10	2:50	3160	L	M.F.
3021	28	2	4	41	Foetal Distress; D.T.A.	8:45	1:40	3500	L	M.F. + M.F.
3035	21	—	—	39	Prolonged 2nd Stage	7:50	2:40	2930	L	M.F.
3049	30	2	5	41	Prolonged 2nd Stage	73:10	2:00	3550	L	M.F. + M.F.; P.P.H.; Manual Removal of Placenta.
3116	21	—	—	38	"	5:25	2:45	2340	L	M.F. + M.F.
3139	25	—	—	40-2	Prolonged 2nd Stage;	66:15	1:03	3050	L	M.F.
3155	28	—	—	36-5	Foetal and Maternal Distress	13:15	2:05	2466	L	M.F. + M.F.
3320	24	—	—	38-6	Prolonged 2nd Stage; P.E.; D.T.A.	12:40	2:35	3200	L	M.F.
3446	25	—	—	38-2	Maternal Distress	49:00	3:07	3030	L	L.F.
3463	27	—	—	43	Prolonged 2nd Stage	8:50	2:20	2880	L	M.F. + M.F.
3500	26	—	2	41-6	"	7:35	0:20	3420	L	M.F.
3578	26	1	2	?	Uterine Inertia, Secondary;	6:15	1:30	3030	L	L.F. Hydrops
3641	31	—	1	38-4	Prolonged 2nd Stage	7:00	0:35	2680	L	L.F.
3656	25	—	—	39-1	Brow Presentation	26:20	2:05	2960	L	M.F. + M.F.
3743	24	—	—	41-5	Foetal and Maternal Distress; D.T.A.	46:00	2:30	3000	L	M.F.
3799	31	—	—	36	Prolonged 2nd Stage	46:15	(1) 2:30 (2) 2:50	(1) 2300 (2) 2180	L	M.F. + M.F.
3833	43	4	3	39-5	"	49:45	0:15	3570	L	L.F.
3863	29	—	—	39-6	Foetal Distress; P.E.;	6:30	2:20	2920	L	M.F.
3921	28	—	1	36-3	Uterine Inertia	15:25	2:00	3300	L	L.F.
3926	42	1	—	40-5	Prolonged 2nd Stage; Foetal Distress	28:30	4:10	4850	L	M.F. + M.F. Hydramnios
4031	31	3	—	40-3	Prolonged 2nd Stage; D.T.A.	3:10	2:40	3005	L	M.F. + M.F.
4056	42	—	7	40-2	Prolonged 2nd Stage	7:50	0:30	3550	L	M.F.
4084	26	—	—	39-6	Mitral Stenosis	46:05	2:45	3345	L	L.F.
4179	28	—	—	40-3	Prolonged 2nd Stage; D.T.A.	61:45	1:15	3062	L	M.F.
4204	24	—	—	41-5	Foetal Distress	3:10	3:40	2500	L	M.F. 2nd Twin
4352	38	3	8	38-4	Prolonged 2nd Stage;	10:00	1:35	2948	L	M.F. + M.F.
4414	29	1	1	41	Secondary Uterine Inertia	18:10	2:55	3200	L	"
4435	23	—	—	40-4	Prolonged 2nd Stage	17:25	0:40	3200	L	"
4437	39	—	—	44	Previous L.S.C.S.				L	M.F.
					Borderline Contracted Pelvis					

FORCEPS DELIVERIES (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	After 28 wks.	Maternity	Indication	Duration of 1st Stage	Duration of 2nd Stage	Weight of Child	Result	Remarks
4563	34	—	1	39-5	Prolonged 2nd Stage	9:15	2:05	3125	L	M.F.
4682	24	1	—	42-4	Uterine Inertia Prolonged 2nd Stage Fetal Distress	10:55	2:15	2640	L	M.R. + M.F.
4797	27	—	—	39-5	Deep Transverse Arrest	18:25	2:45	3180	L	M.F. 1st Twin
4800	24	—	—	37-3	Prolonged 2nd Stage	1:15	1:55	2650	L	L.F.
4845	27	—	—	36-3	Uterine Inertia Prolonged 2nd Stage Secondary Uterine Inertia	7:00	2:15	3280	L	L.F.
4890	28	—	—	41-5	Pre-eclampsia Prolonged 2nd Stage	16:20	2:15	3170	L	M.R. + M.F.
4903	26	—	—	42	Uterine Inertia Prolonged 2nd Stage	12:00	2:25	3800	L	"
4925	43	—	5	39-6	Deep Transverse Arrest Fetal Distress Prolonged 2nd Stage Pre-eclampsia	16:00	2:30	4200	L	"
4958	26	—	1	41	Deep Transverse Arrest Prolonged 2nd Stage	10:10	2:20	3300	L	N.D.
4995	23	—	—	38-4	Retraction Ring Prolonged 2nd Stage	10:55	2:45	3150	L	L.F.
5045	35	—	1	41-3	Secondary Uterine Inertia Prolonged 2nd Stage Fetal Distress	18:00	1:35	3175	L	M.R. + M.F.
5292	27	—	—	35	"	19:00	2:35	2520	L	M.F.
5315	27	—	—	41-6	A.C. Head	15:35	0:40	4000	L	L.F.
5330	30	—	—	39-4	Mitral Stenosis and Ventricular Tachycardia	9:45	0:35	2280	L	M.R. + M.F.
5370	32	—	—	44-3	"	94:30	0:15	3020	L	M.F. for 1st Twin
5393	21	—	—	37	Eclampsia	4:45	2:15	3170	L	P.F., L.F.
5494	30	—	—	39-2	Prolonged 2nd Stage Secondary Uterine Inertia	11:45	2:30	2900	L	L.F.
5582	23	—	5	40-3	Uterine Inertia Prolonged 2nd Stage Uterine Inertia	17:45	3:15	3440	L	M.R. + M.F.
5720	21	—	—	39-3	"	4:40	2:50	3480	L	"
5762	21	—	—	39-6	"	7:00	3:40	3100	L	L.F.
5763	35	—	4	39-3	"	2:10	2:20	3000	L	M.R. + M.F. for O.T.
5804	38	—	1	40	Prolonged 2nd Stage Fetal Distress	38:30	2:05	3430	L	S.B. M.R. + M.F.
5853	23	—	—	40-1	Prolonged 2nd Stage	20:30	2:30	3230	L	M.F.
5897	28	—	—	40-1	Prolonged 2nd Stage Maternal & Fetal Distress	21:30	2:10	3005	L	"

FORCEPS DELIVERIES (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks., 28 wks., After	Maturity	Indication	Duration of Labour 1st Stage 2nd Stage	Weight of Child gm.	Result M. C.	Remarks			
5948	38	1	5	40-2	Prolonged 2nd Stage	7:05	1:40	3720	L	L	M.F.
5968	23	—	—	34-1	Foetal Distress Maternal Distress	32:00	1:00	3770	L	L	"
5991	32	1	2	39-4	Primary Uterine Inertia	12:40	1:10	3350	L	L	L.F.
5995	29	—	5	40-2	Prolonged 2nd Stage	6:35	1:25	3450	L	L	L.F.
6136	31	—	—	42-1	Foetal Distress	24:40	0:20	2935	L	L	M.F.
6144	30	—	—	37-1	Elderly Primigravida Foetal Distress	33:00	3:05	3420	L	L	"
6158	26	—	—	41-2	Foetal Distress	20:40	0:20	2550	L	L	L.F.
6166	38	2	—	38	Prolapse of Cord	2:15	1:50	2750	L	L	L.F.
6196	26	—	—	41-3	Prolonged 2nd Stage	27:40	3:10	3374	L	L	M.R. + M.F.
6341	24	—	—	39-3	"	13:50	2:30	3050	L	L	+ M.F. for 1st Twin
6492	26	—	1	39-6	"	12:30	1:45	2510	L	L	M.F.
6493	36	—	—	40-4	P.E.; Elderly Primigravida	16:30	1:15	3550	L	L	M.F.
6566	40	—	—	39-6	Prolonged 2nd Stage; Severe P.E.; Elderly Primigravida	8:30	2:40	2450	L	L	L.F.
6599	22	—	—	42	Maternal and Foetal Distress;	52:30	1:30	3374	L	L	L.F.
6600	45	—	—	38-4	Uterine Inertia	2:25	1:25	3340	L	L	L.F.
6644	21	—	—	40	Prolonged 2nd Stage	8:20	2:10	4040	L	L	L.F.
6668	23	—	—	40	Severe P.E.; Prolonged 2nd Stage	13:50	1:50	3000	L	L	P.E. M.F.
6680	27	—	—	41-2	Foetal Distress	36:45	2:35	2760	L	L	M.R. + M.F. for O.T. Position
6761	24	—	—	40-2	Prolonged 2nd Stage; Foetal Distress	29:30	0:50	3320	L	L	M.R. + M.F.
6764	37	4	—	39-4	P.O.P.; P.E.	23:10	2:15	3150	L	L	M.R. + M.F.
6818	27	—	—	39-5	Prolonged 2nd Stage; Foetal Distress	20:10	2:38	3515	L	L	L.F.
6844	32	—	1	41-1	Prolonged 2nd Stage; Contracted Outlet	10:30	1:20	2948	L	L	M.F.
6925	23	—	—	40-6	Maternal Distress	19:15	0:15	3260	L	L	L.F.
6929	23	—	1	39	Previous C.S.;	10:00	1:35	2608	L	L	L.F.
6935	27	—	—	41-3	Prolonged 2nd Stage	9:40	0:07	3175	L	L	L.F.
6948	23	—	—	39-5	A.C. Head	5:30	2:28	2302	L	L	M.F.
6955	21	—	—	42-3	Prolonged 2nd Stage; Foetal Distress	42:05	2:05	3000	L	L	M.R. + M.F.; P.E.
6957	26	—	—	2	Foetal Distress	8:15	0:35	2200	L	L	M.R. + M.F.
6963	24	—	1	37	P.E.; Prolonged 2nd Stage	10:30	1:35	3370	L	N.D.	M.F.; Hydramnios.
6970	25	—	—	40-1	Prolonged 2nd Stage	14:30	2:40	3200	L	L	M.F.
6975	26	—	—	39-1	P.E.; Mild Contracted Pelvis	12:50	0:40	2722	L	L	M.R. + M.F.; Manual Removal of Placenta.
7011	26	—	1	39-2	Maternal Distress; Prolonged 2nd Stage;	19:15	1:30	3222	L	L	M.R. + M.F.; Manual Removal of Placenta.
7024	23	—	—	39-5	Previous C.S. Elective	29:30	0:30	2778	L	L	P.E.; Elective L.F.

FORCEPS DELIVERIES (Continued)

Reg. No.	Age	Previous Pregnancies		Maturity	Indication	Duration of Labour		Weight of Child gm.	Result	Remarks
		Before 28 wks.	After 28 wks.			1st Stage	2nd Stage			
7042	32	---	8	41	Prolonged 2nd Stage	7:25	1:50	3005	L	M.F.
7071	21	---	---	39.5	A.C. Head	4:00	0:45	3250	L	L.F.
7182	37	---	6	42	Foetal Distress	8:40	0:25	3470	L	P.P.H.; M.F.
7145	30	---	---	35-1	Prolonged 2nd Stage; Previous C.S.	4:20	2:35	2850	L	M.F.; P.P.H.
7320	23	---	---	39.2	Prolonged 2nd Stage	24:24	2:20	3800	L	L.F.
7390	28	---	---	39.5	Prolonged 2nd Stage; Foetal Distress	40:25	2:10	3053	L	L.F.
7495	36	---	---	37	Eclampsia; Prolonged 2nd Stage	19:30	2:45	2750	L	M.F.
7521	28	---	---	37.1	Prolonged 2nd Stage	30:45	2:35	3300	L	M.F.
7536	20	---	---	37.3	Severe P.E.; Prolonged 2nd Stage	14:20	2:00	2480	L	M.F.
7546	26	---	---	39.3	A.C. Head	24:40	0:10	2600	L	L.F.
7636	28	---	---	38	Prolonged 2nd Stage	7:15	2:15	2400	L	M.R. + M.F.
7652	25	---	---	41.2	Prolonged 2nd Stage; Foetal Distress	4:15	2:15	3400	L	M.F.
7679	21	---	---	44	Prolonged 2nd Stage; Contracted Outlet	5:15	2:25	3250	L	M.F.
7705	24	---	---	38.6	Prolonged 2nd Stage	6:35	2:35	2950	L	L.F.
7720	39	---	---	39.4	Elderly Primigravida	22:00	1:35	3446	L	M.R. + L.F.
7742	22	---	1	41.6	Previous C.S.; Mild Contracted Pelvis	11:50	1:05	3289	L	M.R. + M.F.
7796	25	---	---	40	P.E.; Foetal Distress	19:25	0:15	3060	L	L.F.
7827	33	---	---	40	Prolonged 2nd Stage	15:20	3:30	3200	L	M.F.
7922	23	---	---	43	Prolonged 2nd Stage; Foetal Distress	9:00	2:00	3400	L	L.F.
7931	24	---	1	39.3	Previous C.S.; P.E.	16:15	0:35	3402	L	L.F.
7945	31	---	---	43	Prolonged 2nd Stage	37:30	2:40	2340	L	M.R. + M.F.
7971	38	---	7	43	" "	23:50	1:20	4050	L	M.R. + M.F. for O.T. Post- tion
8008	25	---	1	40.3	Previous C.S.	8:15	0:25	2900	L	M.F.; P.P.H.
8016	22	---	---	40.3	Prolonged 2nd Stage	31:30	2:35	2700	L	M.F.
8017	26	---	---	40.1	Prolonged 2nd Stage; Uterine Inertia	25:45	3:30	3657	L	M.R. + M.F.
8020	27	---	---	38.1	Elective	7:30	1:30	2800	L	Elective L.F.
8024	27	---	1	39.4	Prolonged 2nd Stage	20:40	1:25	3220	L	L.F.; P.P.H.
8051	42	---	5	34.1	" "	5:15	1:55	4320	L	M.R. + M.F. for O.T. Post- tion
8053	25	---	---	40.5	Elective	7:10	0:25	3289	L	Elective L.F.; Manual Removal of Placenta; Vaginal Septum.
8072	20	---	---	43.6	Eclampsia	6:15	2:00	3000	L	L.F.
8087	32	---	---	38.4	Prolonged 2nd Stage	14:45	2:30	3005	L	M.R. + M.F. for O.T. Post- tion; P.P.H.
8231	30	---	---	37.1	" "	12:00	2:20	2090	L	M.F.

FORCEPS DELIVERIES (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	After 28 wks.	Maturity	Indication	Duration 1st Stage	Duration of Labour 2nd Stage	Weight of Child	Result	C.	Remarks
NON-BOOKED											
733	22	—	—	40	Prolonged 2nd Stage	82:20	2:45	3260	L	L	M.R. + M.F.
913	36	—	—	42.5	Deep Transverse Arrest	9:15	4:15	2550	L	N.D.	Anacmia; M.R. + M.F.
1140	28	—	—	39	Prolonged 2nd Stage	50:00	0:45	3000	L	N.D.	L.F.
1482	25	—	—	41.3	Foetal Distress A.C. Head	9:10	0:42	2870	L	L	M.F.
1919	27	—	—	39.6	Foetal Distress	22:40	2:40	2948	L	L	M.R. + M.F.
2316	24	—	—	36.3	Prolonged 2nd Stage	14:15	6:55	3062	L	N.D.	M.R. + M.F.
3110	25	—	1	39.5	O.P. Position Severe Pre-Eclampsia Maternal Distress	?	4:10	2850	L	L	M.R. + M.F.
3129	23	—	—	42	Deep Transverse Arrest Prolonged 2nd Stage Pre-Eclampsia	25:00	3:05	3400	L	L	M.F.
3390	27	—	—	44	Mild Contracted Pelvis Foetal Distress	5:30	1:55	3740	L	L	M.F.
3466	35	—	—	41.2	Prolonged 2nd Stage Foetal Distress	16:20	3:10	3200	L	L	M.F.
3704	23	—	—	35.3	Elderly Primigravida Prolonged 2nd Stage Contracted Pelvis	51:30	2:15	2400	L	N.D.	M.F.
4519	33	—	5	36	Prolonged 2nd Stage Prolonged 2nd Stage Uterine Inertia	3:05	2:00	2850	L	L	M.F.
5021	25	—	—	41.1	Eclampsia	21:15	0:20	2948	L	L	L.F.
5022	30	—	—	38	Prolonged 2nd Stage	5:40	1:57	3470	L	L	M.R. + M.F.
8148	35	—	4	40.2	Prolonged 2nd Stage; Uterine Inertia	16:00	3:50	3620	L	L	M.F.; P.P.H.

TABLE 30

RUPTURE OF UTERUS

There were 7 cases of rupture of uterus, 6 in Booked and one in Non-booked cases.

There was one maternal death.

All infants except 2 were lost. These 2 infants were delivered from mothers, in whom, the occult and incomplete rupture was diagnosed at the time of Caesarean Section.

Reg. No.	Age Before Maturity	Pregnancies Before 28 wks. 28 wks.	Time of Rupture	Type of Rupture	Site of Rupture	Cause of Rupture	Type of Labour	Duration of Labour	Weight of Baby	Result M. C.	Remarks						
								1st Stage									
								2nd Stage	Operation								
								3rd Stage									
BOOKED																	
1409	33	—	5	42	7 hrs. After Onset of Labour	Complete	Posterior Wall of Lower Segment	Neglected Labour	Spont.	7 hrs.	—	—	D	D			
4022	25	—	1	40-2	—	Incomplete	Previous C.C.S. Scar	Previous C.C.S.	—	—	—	—	—	L	L	Previous C.C.S. at another Hospital	
4321	37	—	4	39-6	5:30 hrs After Onset of Labour	"	Right Lateral Wall of Lower Segment	Unknown	"	3:45	2:50	—	(1) L.S.C.S. Subtotal Hysterectomy (3) Right Salpingo-omphrectomy Subtotal Hysterectomy	3270	L	S.B.	
5374	36	—	2	35-3	8:50 hrs. After Onset of Labour	Complete	Transverse Rupture of Right Posterior Wall of Upper Segment	"	"	8:00	—	—	—	2440	L	S.B.	
5547	39	1	1	34-4	—	Incomplete	Occult Rupture in Old C.C.S. Scar	Previous C.C.S. Scar	"	21:10	3:10	—	L.S.C.S.	100	L	L	Failed Trial Labour
6290	34	—	7	39-2	16:30 hrs. After Onset of Labour	Complete	Right Anterior Wall of Lower Segment	Grand Multipara	"	15:30	1:30	—	Subtotal Hysterectomy	3459	L	S.B.	Shock
NON-BOOKED																	
1453	38	—	6	43-5	22:30 hrs. After Onset of Labour	"	Anterior Wall of Lower Segment Extending to the Cervical Canal	(1) Pitocin given by General Practitioner During 2nd Stage of Labour (2) Contracted Inlet (O.C. 9.7 cm) (T.D. 11.9 cm)	"	16:40	—	—	L.S.C.S. Subtotal Hysterectomy	?	L	S.B.	

TABLE 31

VERSION IN LABOUR

There were 18 cases, 13 in Booked and 5 Non-booked cases.

There were no maternal deaths.

There were 8 stillbirths, 5 Booked and 3 Non-booked; 2 neo-natal deaths, one Booked and one Non-booked.

There were 6 second twins in this series.

Reg. No.	Age	Previous Pregnancies Before After 28 wks. 28 wks.	Maturity	Indication	Bipolar, External, or Internal	Weight of Child Em.	Result M, C.	Remarks
395	34	—	35-4	Shoulder Presentation	Internal Version	2470	L	P.E. Accidental Haemorrhage
407	39	—	30-1	"	Bipolar Version	1013	L S.B.	Manual Removal of Placenta
660	39	1	33-3	Compound Presentation	"	1869	L S.B.	Mitral Stenosis + Aortic Regurgitation Class IV
1649	31	—	36-4	Prolapse of Cord	"	2565	L	Placenta Praevia Type II
3871	31	—	39-3	Shoulder Presentation	"	2830	L	Prolapsed Cord
3874	29	—	40	Oblique Lie	Internal Version	3740	L	2nd Twin
4797	27	—	39-5	Shoulder Presentation	Internal Version	1900	L	P.P.S.
5028	29	—	35-6	Failed Forceps	Internal Version	2200	L S.B.	2nd Twin
5370	32	—	44-3	Shoulder Presentation	Internal Version	2350	L	2nd Twin; A.P. Eclampsia; Internal Version, of Placenta, P.P.H.
6492	26	—	39-6	Eclampsia	Internal Version	2500	L	Breech Extraction; Manual Removal of Placenta, P.P.H.
7873	26	1	29-4	Compound Presentation	Internal Version	1400	L	2nd Twin
7895	27	—	3-1	Placenta Praevia	"	2460	L S.B.	Blood Transfusion 1500c.c.
8312	30	—	30-1	Type III. Transverse Lie	"	2220	L	2nd Twin
				Shoulder Presentation	Internal Version		L Mac.	
				Transverse Lie	"		L	
NON-BOOKED								
1706	25	—	3	Shoulder Presentation	Bipolar Version	1474	L S.B.	P.F.
2837	32	—	40-1	Shoulder Presentation	Internal Version	1950	L N.D.	2nd Twin
3605	27	—	40-4	Oblique Lie	Internal Version	2588	L S.B.	P.E.
3862	26	—	1	Shoulder Presentation	Bipolar Version	1250	L S.B.	2nd Twin
5124	30	—	40-4	Shoulder Presentation	Internal Version	3325	L	Prolapse of Cord

TABLE 32
EMBRYOTOMY AND CRANIOTOMY

There were two Booked cases.
There were no maternal deaths

Reg No	Age	Previous Pregnancies Before 28 wks	After 28 wks	Maturity	Indication	Type of Pelvis	Duration of Labour 1st Stage	2nd Stage	Type of Operation	Weight of Child gm	Result to Mother	Remarks
407	39	—	3	30 1	Shoulder presentation	Normal	27 5	—	Bipolar Version with Craniotomy	1013	I	Pre Eclampsia, Mixed Accidental Haemorrhage, Vaginal Lochia
4138	28	—	1	?	Foetal Ascites	—	20 45	0 0	Perforation of Abdomen	2460	L	

BOOKED

TABLE 33

PUERPERAL PYREXIA, GENITAL INFECTION.

This table has been prepared from the records of hospital cases who developed fever whilst in the wards.

The standard used was that laid down by the Puerperal Pyrexia Regulations (England and Wales, 1951) which are defined as including "any febrile condition occurring in a woman in whom a temperature of 100.4°F. (38 degrees C.) or more has occurred within fourteen days after confinement or miscarriage".

There were 62 cases—Incidence of 0.82%, of puerperal pyrexia due to genital infection, 58 in Booked and 4 Non-booked cases. There were no maternal deaths.

Reg. No.	Age	Maturity	Method of Delivery Operation, Induction Manual Removal etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in Days	Results	Remarks
BOOKED									
235	23*	42	Spont.	Perineal Wound Infection	—	Antibiotics	1	L	
407	39	30-1	Bipolar Version Craniotomy	Endometritis	—	Chemotherapy	3	L	P.E. Mixed Rhage
825	28	40	M.R. + M.F.	Perineal Wound Infection	—	"	6	L	
938	28*	40-5	"	Perineal Wound Infection	—	Antibiotics	3	L	Prolonged Labour D.A.A.
1339	25	?	"	Perineal Wound Infection	—	"	1	L	
1511	24*	40	M.F.	Parametritis	—	"	6	L	
1537	23*	38	Spont.	Endometritis	—	"	1	L	
1540	19*	40-5	"	"	—	"	7	L	
1541	25*	40-2	L.S.C.S.	"	—	"	7	L	
1564	27*	42-1	Spont.	Perineal Wound Infection	—	"	3	L	Sterile Culture
2529	29	43-3	"	"	—	"	2	L	
2530	24*	40-1	"	Epistomy, Wound Infection	—	"	1	L	
3136	18*	40-3	"	Endometritis	—	"	4	L	
3146	29	?	"	Perineal Wound Infection	—	"	2	L	
3175	27	38-1	"	"	—	"	2	L	
3305	34	41-1	"	Endometritis and Parametritis	—	"	3	L	

* Primigravida

PUERPERAL PYREXIA, GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery Operation, Induction Manual Removal etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in Days	Recovery	Remarks
3387	24*	34-1	Spont.	Epistomy Wound Infection	—	Antibiotics	3	L.	
3405	24*	38-3	"	Perineal Wound Infection	—	"	1	L.	
3498	38	38-3	"	Endometritis + Breast Engorgement	—	"	2	L.	
3686	32	40-2	"	Endometritis and Parametritis	—	"	2	L.	
4017	28*	35	"	"	—	"	2	L.	
4230	27	38-2	"	"	—	"	1	L.	
4346	25	35	"	"	—	"	2	L.	
4384	30	39-2	"	"	—	"	1	L.	
4494	29	41	"	Parametritis Left Parametritis, Left Pelvic Abscess	—	"	25	L.	Culture Sterile, D.A.A.
4944	30*	39-4	"	Perineal Wound Infection	—	"	1	L.	
4958	26	41	M.F.	Endometritis	—	"	1	L.	
5069	19*	39-1	Spont.	Sepsis in Wounds in Both Labia Majora & Around Introitus	—	"	4	L.	D.A.A.
5075	26*	36-2	"	Perineal Wound Infection	—	"	1	L.	
5288	31*	35-4	"	Endometritis	—	"	2	L.	
5313	23*	30-5	"	"	Staph. Aureus	"	3	L.	
5347	27*	38	"	Perineal Wound Infection	—	"	1	L.	
5610	23	40-2	"	Parametritis Endometritis	Strept. Viridans	Antibiotics Chemotherapy	1	L.	
5748	18*	37	"	Epistomy Wound Infection	—	"	2	L.	
5833	23*	40-1	M.F.	"	—	"	1	L.	
5946	37	38-3	Spont.	Parametritis	—	"	3	L.	
6135	25*	39-5	"	Endometritis	—	"	4	L.	
6136	31*	42-1	M.F.	Epistomy Wound Infection	B. Coli	"	4	L.	
6144	30*	37-1	M.F.	Perineal Wound Infection	B. Coli	"	2	L.	
6348	28	40-5	Spont.	"	—	Chemotherapy	1	L.	
6520	21*	39-3	"	"	—	Antibiotics	3	L.	
6581	24*	40-5	"	"	—	"	3	L.	
6741	26*	38-4	Spont.	Perineal Wound Infection	—	"	3	L.	
6748	34	41-6	"	Endometritis	—	"	2	L.	
6909	29	41-3	"	"	—	"	2	L.	

* Primigravida

PUERPERAL PYREXIA, GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery Operation, Induction Manual Removal etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in Day	Results	Remarks
6826	20*	38	Spont.	Retained Foreign Body In Vagina	--	Antibiotics	3	L	
6975	26	39-1	M.F.	Perineal Wound	--	.	1	L	
7008	43	37-6	Spont.	Endometritis	--	.	3	L	
7263	23*	42	..	Perineal Wound Infection	Septic Haemolyticus	.	3	L	
7484	25*	38-3	..	Endometritis	--	.	3	L	
7513	38	40-4	..	Endometritis	--	.	3	L	
7542	21*	42-5	..	Perineal Wound Infection	--	..	4	L	
7559	28*	40-2	..	Endometritis	--	.	2	L	
7574	27	37-5	..	Endometritis	--	.	1	L	
7608	24*	36	..	Perineal Wound Infection	--	.	3	L	
7680	28*	41-4	..	Endometritis	.	.	2	L	
7817	26	39-1	..	Perineal Wound Infection	--	.	1	L	
7919	28*	39-3	..	Endometritis	--	.	3	L	
NON-BOOKED									
638	22*	40-2	L.S.C.S.	Endometritis	--	Antibiotic	3	L	
1425	22*	41	Spont.	Septic Episiotomy Wound	--	..	1	L	
3110	25	39-5	M.R. + M.F.	Episiotomy Wound Suppurated	--	..	1	L	
6280	20*	37-4	Spont.	Perineal Wound Infection	--	Chemotherapy	4	L	

* Primigravida

TABLE 34

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION

There were 365 cases—an incidence of 4.83% of puerperal pyrexia due to extra-genital infections, 346 in Booked and 19 in Non-booked cases.

276 cases were delivered spontaneously; 28 with forceps, 2 with internal version and one with subtotal hysterectomy.

Caesarean Sections were performed on 58 cases, 51 Booked and 7 Non-booked.

Reg. No.	Age	Maturity	Method of Delivery, Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in days	Result	Remarks
39	33	37-5	Spont.	Breast Engorgement	—	Expectant	1	L	Sterile Culture
69	20*	39-5	"	Cystitis	—	Chemotherapy	1	L	
96	23*	38-4	"	Breast Engorgement	—	Expectant	1	L	
183	36	39-3	"	"	—	"	1	L	
196	19*	40-1	"	"	—	Antibiotics	2	L	
222	31	?	Spont. Manual Removal of Placenta	"	—	"	2	L	
237	18*	38-1	L.S.C.S.	"	—	Expectant	1	L	
248	21*	?	Spont.	"	—	"	1	L	
314	24	40-2	"	Unknown	—	"	2	L	
334	32*	39-4	L.S.C.S.	Pyelitis	B. Coli	Chemotherapy	5	L	Infective C S
376	25	38-4	Spont.	Unknown	—	Antibiotics	2	L	Pulmonary T B
405	32	37-3	"	"	—	Expectant	1	L	P P S
481	28	40-1	"	Urinary Infection	Staph. pyogenes aureus	Antibiotics	4	L	
506	23	30	L.S.C.S.	Pyelitis	B. Coli	Chemotherapy	3	L	Placenta Praevia Type III.
519	23*	37-3	Spont.	Breast Engorgement	—	Expectant	1	L	
540	25	38-5	"	"	—	"	2	L	
575	35	40-2	"	Unknown	—	"	2	L	P.E.
576	35	38	"	"	—	"	2	L	
583	23*	42-1	"	Breast Engorgement	—	Chemotherapy	1	L	
609	23	42-4	"	Unknown	—	Antibiotics	1	L	
677	28*	38-5	"	Breast Engorgement	—	Expectant	1	L	
837	33	44-1	"	"	—	"	1	L	
868	29*	40-5	Low Forceps.	"	—	"	1	L	
876	31	43-6	L.S.C.S.	" Cystitis	B. Coli	Antibiotics	2	L	

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery: Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in days	Result	Remarks
894	24	?	L.S.C.S.	Unknown	—	Chemotherapy	1	L	
935	28	40	Breast Engorgement	Breast Engorgement	—	Expectant	1	L	
942	23*	37-1	Spont.	Unknown	—	—	1	L	
967	34	35	"	Breast Engorgement	—	Expectant	1	L	
985	24*	40	"	"	—	"	1	L	
1005	18*	36-4	L.S.C.S.	Unknown	—	Antibiotics	3	L	
1007	26	40-5	Spont.	Breast Engorgement	—	Expectant	1	L	Active Pulmonary T.B.
1031	22*	40-4	"	Unknown	—	—	1	L	
1089	31*	40-3	L.S.C.S.	Post-operative Reaction	—	Antibiotics	1	L	
1111	29	3	Spont.	Breast Engorgement	—	Expectant	1	L	
1340	44	38-2	"	"	—	"	1	L	
1366	24*	41-2	"	"	—	"	1	L	
1378	18*	42	"	"	—	"	1	L	
1402	29	39-3	"	"	—	"	1	L	
1414	27	38-3	L.S.C.S.	Post-operative Reaction	—	Antibiotics	1	L	
1421	30*	38-1	"	Breast Engorgement	—	Expectant	1	L	F.B.
1427	22*	40-1	Spont.	Filariasis	Microfilariae	Hextrazan	5	L	
1439	24	39	"	Breast Engorgement	Banctoflu	Expectant	1	L	
1443	30	36-6	"	"	—	"	1	L	
1455	33	38-6	"	Unknown	—	"	1	L	
1475	21*	37-3	"	Breast Engorgement	—	Expectant	1	L	
1489	24	37-6	"	"	—	"	1	L	
1571	26*	37-2	Spont.	"	—	"	1	L	P.P.H. (1,000 c.c. Blood Loss).
1575	26*	42-5	(1) Compound — Assisted (2) Vertex — Mid Forceps	Unknown	—	Antibiotics	1	L	Twin Pregnancy
1680	28*	39-3	Breech Assisted	Breast Engorgement	—	Expectant	1	L	
1697	27*	38-2	Spont.	Cystitis and Urinary Infection	B. Coli	Antibiotics	1	L	
1702	27	40	"	Breast Engorgement	B. Coli	"	2	L	
1735	29	40-4	"	Breast Engorgement	—	"	1	L	
1819	31	41-4	"	Unknown	—	Expectant	2	L	
1830	19*	44-4	"	Breast Engorgement	—	"	2	L	
1918	41	34-4	"	"	—	"	1	L	
1931	19*	35-6	L.S.C.S.	"	—	—	1	L	
1936	37	41-1	Spont.	Unknown	—	Expectant	1	L	
2074	37	36-2	"	Breast Engorgement	—	Antibiotics	1	L	
2110	26	?	"	"	—	"	1	L	
2158	27	41-2	"	"	—	"	1	L	
2198	25*	40-5	"	"	—	"	1	L	
2225	24	38-2	Mid Forceps	"	—	"	1	L	
2233	25	38-6	Spont.	"	—	"	1	L	
2248	21*	38-4	"	Lobar Pneumonia	Pneumococci	Chemotherapy	1	L	
			"	Breast Engorgement	—	Expectant	1	L	

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery, Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms if Known	Treatment	Months in days	Result	Remarks
2321	30	34-3	Spont.	Breast Engorgement	—	Expectant	2	L.	
2341	30	41	"	Breast Engorgement	—	"	1	L.	
2365	30	38-1	"	Acute Bronchitis	—	Antibiotics	1	L.	
2380	29	40-2	"	Breast Engorgement	—	Expectant	1	L.	
2400	24	41-6	"	"	—	"	1	L.	
2420	25*	40-2	"	"	—	Antibiotics	1	L.	
2469	38	7	"	Retained Abdominal Swab	—	"	5	L.	P.P.S.
2495	20*	44	"	Furteritis	—	"	1	L.	
2504	22	40-4	"	Unknown	—	Expectant	1	L.	
2534	35	34-4	"	"	—	Antibiotics	1	L.	
2542	26	37-1	"	"	—	"	1	L.	
2558	25	40	(1) Spont. + M.F. (2) M.R. + M.F.	Infectious Hepatitis	—	Expectant	3	L.	Twin Pregnancy
2577	24*	39-5	Spont.	Breast Engorgement	—	Antibiotics	1	L.	D.A.A.
2588	22*	39-3	"	Unknown	—	Expectant	1	L.	
2602	31	44-5	"	Breast Engorgement	—	"	1	L.	
2603	26*	40-6	"	Bronchitis	—	Expectant	2	L.	
2604	23	41-3	"	Breast Engorgement	—	"	1	L.	
2633	30	38	"	Unknown	—	Expectant	1	L.	
2639	23	37-3	"	Breast Engorgement	—	"	1	L.	
2641	31	37-5	"	"	—	"	2	L.	
2652	29	40	"	Bronchitis	—	Antibiotics	1	L.	
2665	23*	40	"	Breast Engorgement	—	Prophylactic Antibiotics	4	L.	D.A.A.
2683	25*	39	"	"	—	Expectant	1	L.	
2704	33	39-5	"	Unknown	—	"	1	L.	
2721	22*	39-5	"	Breast Engorgement	—	Expectant	1	L.	
2771	42	35-5	"	Unknown	—	Antibiotics	1	L.	D.A.A.
2787	33*	39-5	"	Urinary Infection	B. Coli	"	1	L.	Twin Pregnancy
2806	39	32-4	"	Unknown	—	"	1	L.	
2845	25	39	"	Breast Engorgement	—	"	1	L.	
2878	25*	39-3	"	"	—	"	1	L.	
2880	26	39-3	"	Breast Engorgement	—	"	1	L.	
2897	39	26*	"	"	—	"	1	L.	
2935	37	43	"	"	—	"	1	L.	
2980	29	39-3	"	"	—	"	1	L.	
2999	35	40-4	"	Breast Engorgement	—	Antibiotics	2	L.	
3049	30	41	M.R. + M.F.	Unknown	—	Expectant	1	L.	
3057	25	34-3	L.S.C.S.	Breast Engorgement	—	Antibiotics	1	L.	
3078	32	38-6	Spont.	Unknown	—	Expectant	1	L.	
3086	28	38-6	L.S.C.S.	Breast Engorgement	—	Antibiotics	1	L.	
3102	26*	41-4	Spont.	Unknown	—	"	2	L.	
3109	23*	38-2	L.S.C.S.	Breast Engorgement	—	Expectant	1	L.	Congestive Heart Failure P.P.S.

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery: Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in days	Result	Remarks
3118	37	39.5	Spont.	Breast Engorgement	—	Expectant	1	L	
3133	33	39.6	"	Unknown	—	Antibiotics	1	L	P.F.
3139	25*	40.2	Mid Forceps	Infection of Respirating Tract	—	"	3	L	P.F.
3155	28*	36.5	M.R. + M.L.	Urinary Infection	—	"	3	L	P.F.
3166	28	38.2	Spont.	Unknown	—	"	1	L	
3168	34	41.3	"	"	—	"	1	L	
3176	22*	35.4	"	Breast Engorgement	—	Expectant	1	L	
3257	23*	40.2	"	Unknown	—	"	1	L	
3283	30	37	"	Urinary Infection	—	Antibiotics	1	L	
3317	37	38.5	"	Breast Engorgement	—	Expectant	1	L	
3320	21*	38.6	"	Unknown	—	"	1	L	
3339	21*	40.4	"	Breast Engorgement	—	Antibiotics	1	L	
3359	27	34.4	L.S.C.S.	Unknown	—	"	1	L	
3417	19*	39.2	Spont.	Breast Engorgement	—	Antibiotics	1	L	
3439	22*	37.2	"	"	—	Expectant	1	L	
3489	24*	38.5	"	"	—	"	1	L	
3506	26*	36.4	"	Unknown	—	"	1	L	
3554	28	38.3	"	Urinary Infection	B. Faecalis Alkaligenes	Chemotherapy	2	L	
3632	27	39.2	"	Breast Engorgement	—	Expectant	1	L	
3723	24*	40.6	"	"	—	"	1	L	
3734	32	39	"	"	—	"	1	L	
3741	25*	38	"	"	—	"	1	L	
3743	24*	41.5	Mid Forceps	"	—	"	1	L	
3755	24	36.4	Spont.	"	—	"	1	L	
3796	29*	36.6	"	"	—	"	1	L	
3811	20*	38.6	"	"	—	"	1	L	
3819	23	38.5	"	Unknown	—	"	1	L	
3841	38	38.1	"	Breast Engorgement	—	Expectant	3	L	
3863	29*	39.6	Mid Forceps	"	—	"	1	L	Mitral Stenosis—Class II
3885	32	41.3	Spont.	"	—	"	1	L	
3922	30	37	"	Unknown	—	"	1	L	
3931	21*	39.2	"	Breast Engorgement	—	Expectant	4	L	
3944	36	39	"	Breast Engorgement	—	Expectant	1	L	
3957	27*	36.2	"	Urinary Infection	—	Antibiotics	1	L	
4022	25	40.2	"	Breast Engorgement	—	Expectant	1	L	
4052	42	40.2	L.S.C.S.	"	—	Antibiotics	1	L	Elective C.S.
4075	32	39.1	Mid Forceps	"	—	Expectant	1	L	Mitral Stenosis—Class I.
4087	26	35.5	Spont.	"	—	"	1	L	P.F.
4109	30*	39.2	"	Unknown	—	"	1	L	
4119	29	39.2	"	Ludwig's Angina	—	Antibiotics	2	L	
4133	21*	43	"	Breast Engorgement	—	Expectant	1	L	
4191	39	26.4	L.S.C.S.	Injection Abscess Over Left Buttock	—	Antibiotics	1	L	P.P.S.

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery, Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms if known	Treatment	Morbidity in days	Result	Remarks
4196	28	42-6	Spont.	Breast Engorgement	—	Expectant	1	L	
4204	24*	41-5	(1) Spont. (2) Mid Forceps	"	—	"	1	L	Twin Pregnancy
4206	26*	41-4	Spont.	Unknown	—	"	1	L	
4352	38	38-4	Mid Forceps	Breast Engorgement	—	Antibiotics	2	L	
4379	23*	37-1	Spont.	"	—	Expectant	1	L	
4414	29	41	M.R. + M.F.	" Pyelitis	B. Coli	Antibiotics	2	L	
4449	26	39-6	Spont.	Breast Engorgement	—	Expectant	1	L	
4451	25*	39-3	"	Unknown	—	"	1	L	
4492	19*	37-1	"	Urinary Infection	B. Coli	Antibiotics	1	L	
4497	30*	40-3	L.S.C.S.	Post-operative Reaction	—	"	1	L	
4549	29	41	Spont.	Breast Engorgement	—	Expectant	4	L	
4504	21*	25-3	Spont.	Urinary Infection	B. Coli	Antibiotics	4	L	
4587	20*	40-5	"	Breast Engorgement	—	Expectant	1	L	P.E.
4598	26	"	"	"	—	"	1	L	P.E.
4604	36*	39-2	"	"	—	"	1	L	
4635	30*	40-2	L.S.C.S.	" Enteritis	—	Antibiotics	2	L	
4640	22*	37-2	Spont.	Breast Engorgement	—	"	4	L	
4648	20*	38-6	"	Mild Cystitis	B. Coli	(hemo)therapy	5	L	
4649	32	37-3	"	Breast Engorgement	—	Expectant	1	L	
4660	33	40	"	Post operative Reaction	—	Antibiotics	1	L	P.P.S.
4682	24*	42-4	M.R. + M.F.	" Unknown	—	"	1	L	
4698	27*	36-4	Spont.	"	—	"	1	L	
4716	36	38-6	"	Enteritis	—	"	2	L	
4738	25*	37-2	"	"	—	"	5	L	
4741	34	36-1	"	Unknown	—	(hemo)therapy	2	L	
4748	24	38-6	"	"	—	"	1	L	
4748	27	35-3	"	"	—	"	1	L	
4776	27	35-3	"	"	—	"	1	L	
4797	27*	39-5	(1) Mid Forceps (2) Internal Version	"	—	Antibiotics	1	L	Twin Pregnancy
4821	33	35-5	Spont.	Post-operative Reaction	—	"	2	L	P.P.S.
4845	27*	36-3	Low Forceps	"	—	Prophylactic Antibiotics	2	L	
4855	21*	38-3	Spont.	"	—	Antibiotics	1	L	
4910	30	41	"	"	—	"	1	L	
4924	24*	38-3	"	Bronchitis	—	"	2	L	
4939	27	39-6	"	Breast Engorgement	—	"	1	L	
4950	28*	39-5	"	Unknown	—	Antibiotics	3	L	
5018	28	"	L.S.C.S.	Breast Engorgement	—	"	1	L	
4967	24*	41-2	Spont.	Post-operative	—	"	1	L	
5041	28	40-1	"	Unknown	—	Expectant	1	L	
5124	46	41-1	"	Breast Engorgement	—	(hemo)therapy	2	L	Essential Hypertension.
5195	22*	39-3	"	"	—	Expectant	2	L	D.A.A.

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery: Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in days	Result	Remarks
5233	26	39-6	L.S.C.S.	Post-operative Reaction	—	Antibiotics	1	L.	
5065	20*	38	Spont.	Unknown	—	"	11	L.	Edema of, One FH, Left Heart Failure, Anaemia, Aplasticosis B. P.F.
5251	38*	39-3	L.S.C.S.	Post-operative Reaction	—	Antibiotics	1	L.	
5254	28	41-1	Spont.	Breast Engorgement	—	Expectant	1	L.	
5256	25*	41-4	"	Unknown	—	"	1	L.	
5261	24*	41-5	"	Breast Engorgement	—	Expectant	1	L.	
5273	37	35-1	"	Aggravation of Pulmonary T.B.	—	Antibiotics	1	L.	
5208	25	39-6	"	Breast Engorgement	—	Expectant	2	L.	
5322	20*	41-3	"	"	—	"	1	L.	
5350	30*	39-4	M.R. + M.F.	Unknown	—	Antibiotics	1	L.	Mitral Stenosis Class IV
5360	25*	39	L.S.C.S.	Breast Engorgement	—	"	1	L.	P.F.
5370	32*	44-3	(1) Internal Version (2) Mid Forceps	"	—	"	1	L.	Twin Pregnancy. Manual Removal of Placenta
5393	21*	27	Low Forceps	"	—	Expectant	1	L.	P.F.
5422	22	34-1	Spont.	"	—	"	1	L.	
5424	24*	37-1	"	Unknown	—	"	1	L.	
5426	25*	35-4	"	Breast Engorgement	—	Antibiotics	5	L.	
5447	30*	38-1	L.S.C.S.	Post-operative Reaction	—	"	1	L.	
5456	37*	38-4	Spont.	Breast Engorgement	—	Expectant	1	L.	
5472	27	42	"	Unknown	—	"	1	L.	
5512	21*	39-4	"	Breast Engorgement	—	Expectant	2	L.	
5513	30*	39-3	"	"	—	"	1	L.	
5547	39	34-4	"	"	—	Chemotherapy	2	L.	
5555	36	40-3	L.S.C.S.	Cystitis	—	Antibiotics	2	L.	
5559	37	24	Spont.	Unknown	—	"	2	L.	
5580	21*	40-6	"	"	—	Expectant	1	L.	Abortion. P.P.S., Mitral Stenosis—Class I
5608	25	39-3	"	Breast Engorgement	—	"	1	L.	
5621	24*	36	"	Post-operative Reaction	—	Antibiotics	1	L.	
5639	23*	41-4	L.S.C.S.	Unknown	—	"	1	L.	D.A.A.
5646	24	?	Spont.	Breast Engorgement	—	"	1	L.	
5647	25*	38-4	"	Thrombophlebitis	—	Expectant	1	L.	
5663	40	?	"	Allergy	—	Chemotherapy	1	L.	
5670	23	33-5	C.C.S.	Pneumonia	—	Antibiotics	5	L.	
5682	26*	37	L.S.C.S.	Unknown	—	"	1	L.	
5695	22*	40-4	Spont.	Breast Engorgement	—	Expectant	2	L.	
5733	28*	29-2	"	"	—	"	2	L.	
5743	32	31-2	"	Upper Respiratory Tract Infection	—	Antibiotics	1	L.	

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery: Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in days	Result	Remarks
5774	39	40-6	Spont. L.S.C.S.	Breast Engorgement	—	Expectant	1	L	
5778	20*	36-5		Post-operative Reaction, Multiple Boils in Buttock	—	Chemotherapy	4	L	Post-partum Pyrexiosis, Transferred to Mental Hospital
5781	34	41-2	Spont.	Breast Engorgement	—	Expectant	2	L	
5824	39	34-4	"	Breast Engorgement	—	"	1	L	
5832	29*	39	"	Breast Engorgement	—	"	1	L	
5856	26*	39-2	L.S.C.S.	Breast Engorgement	B. Coli	Antibiotics	2	L	
5885	23*	38-2	"	Urinary Infection	—	"	2	L	
5887	23*	40-5	Spont.	Breast Engorgement	—	"	3	L	
5918	28	40-5	"	"	—	Expectant	1	L	
5942	40	40-3	"	Unknown	—	"	1	L	
5946	37	38-3	"	"	—	"	1	L	
5971	19*	38-1	"	Breast Engorgement	—	Expectant	1	L	
5972	31	37	"	Unknown	—	"	1	L	
5983	26	35-5	"	Breast Engorgement	—	Expectant	1	L	
5995	29	40-2	L.S.C.S.	Post-operative Reaction	—	Antibiotics	1	L	
6088	38*	41-2	Spont.	Breast Engorgement	B. Coli	Chemotherapy	3	L	
6027	23*	44	"	Chronic Lymphatic Leukaemia	—	Antibiotics	3	L	D.A.A.
6048	28*	39-3	"	Breast Engorgement	—	"	1	L	
6056	38	41	"	Leukaemia	—	Expectant	1	L	
6057	28	35-4	"	Pyelonephritis	—	Antibiotics	1	L	
6058	27*	42	"	Breast Engorgement	—	Expectant	1	L	Pulmonary T.B.
6076	31	40-1	"	Urinary Infection	B. Coli	Antibiotics	1	L	
6093	27*	40-3	L.S.C.S.	Post-operative Reaction	—	"	5	L	P.P.S.
6133	24*	35	Spont.	Breast Engorgement	—	Expectant	1	L	
6152	20*	41-3	L.S.C.S.	"	—	"	1	L	P.I., Twin Pregnancy
6158	26*	41-2	Spont.	Unknown	—	"	1	L	
6186	33	38	"	Breast Engorgement	—	Expectant	1	L	
6251	21*	39-4	"	Unknown	—	"	1	L	
6276	36	40-5	"	Post-operative Reaction	—	Expectant	2	L	
6300	34	39-2	L.S.C.S.	Breast Engorgement	—	Antibiotics	2	L	
6308	31*	40-4	"	Unknown	—	"	2	L	
6309	22*	34-6	Subtotal Hysterectomy	"	—	"	2	L	2 Exhaustive C.S., P.P.S., Ruptured Uterus Complete
6309	24*	40-1	"	Breast Engorgement	—	Expectant	5	L	
6314	22	40-5	"	Unknown	—	Antibiotics	1	L	
6318	32*	41-3	L.S.C.S.	Breast Engorgement	—	Expectant	3	L	
6328	38	35-6	Spont.	"	—	"	1	L	
6352	25*	39	"	Urinary Infection	B. Coli	Antibiotics	3	L	D.A.A. P.P.S.
6391	31	40-3	"	Post-operative Reaction	—	"	1	L	
6402	32	33-3	L.S.C.S.	Breast Engorgement	—	Expectant	1	L	

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery: Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms if Known	Treatment	Morbidity in days	Result	Remarks
6428	22	38-2	L.S.C.S.	Breast Engorgement	—	Expectant	1	L	
6477	36	37-4	Spont.	Unknown	—	Chemotherapy	1	L	
6475	27*	41-1	"	Breast Engorgement	—	Expectant	2	L	
6403	36*	40-4	Mid Forceps	"	—	"	1	L	
6504	33*	39-4	Spont.	"	—	Expectant	2	L	
6523	28*	40	"	"	—	"	1	L	
6575	25*	38-4	"	"	—	"	3	L	Pulmonary T. B.
6528	34	39	L.S.C.S.	Post-operative Reaction	—	Antibiotics	1	L	
6541	21*	40-4	"	"	—	Expectant	1	L	
6580	23*	39-2	Spont.	Breast Engorgement	—	"	2	L	
6601	21	35-4	C.C.S.	"	—	"	1	L	
6617	28	36-1	L.S.C.S.	Post-operative Reaction	—	Chemotherapy	1	L	
6644	21*	40	Low Forceps	Drip Reaction	—	Antibiotics	1	L	
6701	23*	38-1	L.S.C.S.	Breast Engorgement	—	"	1	L	
6718	28	41-4	Spont.	"	—	Expectant	1	L	Ante Partum Eclampsia 10 Fits before Admission
6734	23	39-5	"	"	—	"	1	L	
6738	26*	39-6	"	"	—	"	3	L	
6772	26*	39-4	L.S.C.S.	Post-operative Reaction	—	Antibiotics	1	L	
6804	33	39-4	Spont.	Breast Engorgement	—	Expectant	2	L	
6824	25	40-3	"	"	—	"	1	L	
6831	27*	37-5	"	Urinary Infection	B. Coli	Antibiotics	1	L	
6847	40	39-3	"	Unknown	—	"	1	L	
6887	24*	40	"	Breast Engorgement	—	Expectant	1	L	
6976	25*	41-1	"	Unknown	—	"	1	L	
7042	32	36-4	"	Breast Engorgement	—	Expectant	1	L	
7048	31	38-3	Mid Forceps	Post-operative Reaction	—	Antibiotics	1	L	
7059	23*	41-2	Spont.	Cystitis	B. Coli	"	2	L	
7085	31	38-3	"	Unknown	—	"	1	L	
7087	26*	35-5	"	Breast Engorgement	—	Expectant	1	L	
7121	27	44-1	"	"	—	"	1	L	
7151	22*	43-1	"	"	—	"	2	L	
7167	22*	40-1	"	"	—	"	1	L	
7173	40	?	"	"	—	"	1	L	
7182	57	42	Mid Forceps	Unknown	—	Expectant	1	L	
7187	30*	37-6	Spont.	Breast Engorgement	—	Antibiotics	1	L	
7192	34	?	"	Unknown	—	"	2	L	
7267	22*	38-1	"	Urinary Infection	Strept. Faecalis	Expectant	1	L	
7326	37	34	"	Breast Engorgement	—	Antibiotics	3	L	
7371	40	40-6	L.S.C.S.	Peritonitis	—	Expectant	1	L	
7395	36*	37	Mid Forceps	Cystitis	B. Coli	Antibiotics	6	L	P.P.S. Blood Secondary 1,500 c.c.
7413	36	38-4	Spont.	Cellulitis Left Arm	—	"	2	L	Antepartum Eclampsia
7416	25	43	"	Pneumonia	—	"	1	L	

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg. No.	Age	Maturity	Method of Delivery: Operation, Induction, Manual Removal, etc.	Cause of Temperature	Organisms of Known	Treatment	Morbidity in days	Result	Remarks
7447	20*	38-5	Spont.	Breast Engorgement	—	Expectant	1	L	
7536	20*	37-3	Mid. Forceps	" " Unknown	—	"	2	L	P. L.
7550	26	42-2	L.S.C.S.	Breast Engorgement	—	Expectant	1	L	
7579	28	36-5	Spont.	" " Retention of Urine	—	Antibiotics	5	L	
7597	22*	33-4	"	" " Unknown	—	Carbachol			
7607	29	38	"	" " Unknown	—	—	1	L	Mural Stenosis with Con- gestive Heart Failure
7615	26	36-4	"	Breast Engorgement	—	Expectant	1	L	
7621	33	40-6	"	Urinary Infection	B. Coli	Chemotherapy	1	L	
7709	29	38-6	"	Typhoid Fever	—	Antibiotics	10	L	Transferred to Infectious Hospital
7727	22	36	"	Drug Sensitivity	—	Expectant	1	L	
7742	22	41-6	M.R. + M.F.	Breast Engorgement	—	"	1	L	
7768	19*	39	Spont.	" " "	—	"	1	L	
7808	24*	39-2	"	" " "	—	"	2	L	
7814	27	41-3	"	" " "	—	"	2	L	
7824	27*	39-5	Spont.	" " Unknown	—	Antibiotics	2	L	
7839	34	38-1	"	Breast Engorgement	—	Expectant	1	L	
7851	24*	40-2	"	" " Cystitis	—	"	1	L	
7852	32	41-5	"	Breast Engorgement	B. Coli	Antibiotics	4	L	Pulmonary T.B.
7900	34*	36-4	"	" " "	—	Expectant	1	L	D. A. A.
7903	26*	39	Breech-Assisted	Breast Engorgement	—	"	1	L	
7909	25	36-1	Spont.	" " Unknown	—	"	1	L	
7926	27*	40	"	Breast Engorgement	—	Expectant	2	L	
7934	36*	40	"	" " Infection of Abdominal Incision	—	"	1	L	
7944	30	38-3	C.C.S.	" " Infection of Abdominal Incision	—	Antibiotics	2	L	Naegel's Pelvis
7960	32	40-5	Spont.	Urinary Infection	B. Coli	Chemotherapy	2	L	P. P. S.
7964	27	?	L.S.C.S.	Infection of Abdominal Incision	—	Antibiotics	3	L	Placenta Previa Type III
7970	28	38	Spont.	Urinary Infection	—	Expectant	2	L	P. L.
8005	26*	37	"	Breast Engorgement	—	Antibiotics	1	L	
8012	25*	38-4	"	" " "	—	Expectant	2	L	
8027	25	35-4	"	" " "	—	Antibiotics	3	L	
8051	42	34-1	M.R. + M.F.	Breast Engorgement and Infection of Areola	—	Expectant	1	L	
8067	33	41	Spont.	Breast Engorgement	—	"	1	L	
8072	20*	43-6	Low Forceps	" " Eclampsia	—	Antibiotics	2	L	Ante Partum Eclampsia — 1 P. L.
8086	24*	41-2	Spont.	Breast Engorgement	—	Expectant	1	L	
8087	32*	38-4	M.R. + M.F.	Urinary Infection	—	Chemotherapy	4	L	P. P. H.—1,000 c.c. Cervical Tear Repaired
8104	25*	39-2	Spont.	Breast Engorgement	—	Post-operative	1	L	
8144	27	40-4	"	Retention of Urine	—	Antibiotics	21	L	
8179	27*	39-1	"	Breast Engorgement	—	"	1	L	

* Primigravida

PUERPERAL PYREXIA, EXTRA-GENITAL INFECTION (Continued)

Reg No	Age	Maternity	Method of Delivery, Operation, Induction, Manual Removal, etc.	Course of Temperature	Organisms if Known	Treatment	Months in days	Case	Remarks
8209	24*	39 3	L S C S	Breast Engorgement and Urinary Infection	B. Coli	Antibiotics	4	I	
8351	29	41 2	"	Breast Engorgement	"	Expectant	2	I	
8373	29	38	"	Urinary Infection	"	Chemotherapy	7	I	
8377	25*	37	"	Urinary Infection	B. Coli		4	L	
NON BOOKED									
761	21*	40 3	I S C S	Breast Engorgement	"	Expectant	1	L	P. I. Twin Pregnancy
792	35	30 4	Spont	"	"	"	1	I	worm Infestation
1166	26	33 3	"	"	"	"	1	I	Twin Pregnancy Both Breech
1837	38	34 3	L S C S	Post operative Reaction	"	Antibiotics	1	I	
1933	32	37 5	Spont	Eclampsia	"	"	2	I	
2414	23*	38	"	Breast Engorgement	"	Expectant	3	L	
2610	28*	32 4	I S C S	Post operative Reaction	"	Antibiotics	1	I	
3956	28	39	Spont	Breast Engorgement	"	Expectant	1	I	
4152	21*	37	"	Unknown	"	"	1	I	
4519	33	36	"	"	"	Antibiotics	1	I	
5021	25*	41 4	Low Forceps	Breast Engorgement	"	Expectant	1	I	
5022	39	38	M R + M F	"	"	"	1	I	
5965	36	39 5	Spont	"	"	"	1	I	
6311	36	32 1	C. C. S	Post operative Reaction	"	Antibiotics	2	I	D. A. A.
6502	26*	38 1	Spont	Unknown	"	"	1	I	
6649	36	40 4	Norton's Extra peritoneal Caesarean Section	"	"	"	5	I	Fibroid of Cervix
7483	38	35	C. C. S	Breast Engorgement	"	Expectant	1	I	Severe P. I.
7835	33	40 5	Spont	Cystitis	P. Coli	Antibiotics	2	I	
7955	27*	42 4	I S C S	Breast Engorgement	"	Expectant	2	I	

* Primigravida

MATERNAL MORTALITY

There were 4 maternal deaths, 2 Booked and 2 Non-booked cases. These 2 Booked cases died undelivered. They were both over 28 weeks pregnant. The maternal mortality taking it as number of delivered patients is nil, but these 2 fatalities represent a 0.024% maternal mortality for the 8,165 Booked admissions. The 2 deaths among the 195 Non-booked deliveries after the 28th week of pregnancy gives an incidence of 1.02 deaths per 1,000 deliveries.

Causes of Maternal Deaths.

(For short notes on each case, see pages 120-122).

BOOKED:

- | | |
|-------------|---|
| 1. No. 1409 | Ruptured uterus. |
| 2. No. 6485 | Massive haemorrhage from fistula between
aorta and oesophagus. |

NON-BOOKED:

- | | |
|-------------|--------------------------|
| 1. No. 3085 | Eclampsia of pregnancy. |
| 2. No. 3834 | Post-partum haemorrhage. |

DETAILS OF MATERNAL DEATHS

Reg. No. 1409, Booked. Age 33, gravida 6, had had 5 previous normal spontaneous deliveries; birth weights 5-6 pounds. Last delivery 2 years previously.

She attended ante-natal clinic regularly and findings were normal. She was admitted at 42nd week (4/6/56) at 4.15 a.m. in first stage of labour. Examination showed B.P. 100/70 mm Hg and pulse 84 per minute; foetus in vertex R.O.A. position, foetal head not engaged, foetal heart present. Membranes ruptured spontaneously three hours after admission when the os was 4 cm dilated. Pains were moderate, 1 in 5 minutes, and became stronger six hours after admission. Pethidine 100 mg was given intramuscularly at 10.00 a.m. At 11.00 a.m. the patient was in good condition. The pain was less severe but still came at 5 minute intervals. At 11.30 a.m. "the patient was not looking so well." No pulse or blood pressure was recorded at this time. At 11.40 a.m., a vaginal examination showed the os fully dilated, the vertex presenting at low level. At 12.25 p.m. the foetal heart was not heard. There was no mention of the type of uterine contraction or alteration in shape of the uterus at this time. The patient appeared in peripheral circulatory failure with pulse 130/min, B.P. 50/30 mm Hg. She was fully conscious. A nor-adrenaline drip was started. Blood was drawn and sent for typing and crossmatching. Free fluid was demonstrated at 1.30 p.m. Pulse became thready and blood pressure was not registrable. No external bleeding was seen coming from the vagina. Her condition deteriorated in spite of 1 pint of rapid blood transfusion. She vomited at 2.10 p.m., gasped at 2.15 p.m. and expired at 2.20 p.m., undelivered.

Postmortem findings: 2,500 cc of blood in peritoneal cavity. 4 inch tear in posterior wall of uterus and one arm prolapsed through the ruptured site, but the foetal head was well engaged in the R.O.A. position.

Reg. No. 3085, Non-Booked. Age 27, gravida 2. Patient had one full-time normal spontaneous delivery one year previously. She was admitted to hospital on 7th July, 1956 at 12.30 p.m. with the history of blurring of vision, fever, anorexia and vomiting for 4 days; oedema of legs, yellowish discoloration of skin and sclera and passage of dark brown urine for 2 days. L.M.P. was unknown. Examination showed a mentally confused, drowsy, incooperative, and jaundiced patient, with marked ankle oedema. The temperature was 98.4°F, the pulse rate 76/minute, and B.P. 120/90 mm Hg. The uterus was enlarged to about 28 weeks size of pregnancy. The urine contained trace of protein, hyaline casts, and bile pigment but no urobilinogen.

At 5.45 a.m. of the 8th July, 1956, patient had incontinence of urine and became semi-comatous. She developed 4 atypical fits (at 6 a.m., 6.15 a.m. and 10.45 p.m. of the 8th, and 9.40 a.m. of the 9th July, 1956 respectively.) She was given 20% glucose drip and cortisone 40 mg. every 6 hours. She delivered a live female infant of 2300 grams spontaneously at 9.50 a.m. on the 9th of July, 1956. She bled 500 cc after delivery. The placenta showed yellow staining. Her condition deteriorated rapidly after delivery. She died one hour later.

Postmortem findings: Severe jaundice.

Oedema of ankles.

Liver: small, flabby, with multiple small yellowish necrotic spots.

Kidneys: parachymatous degeneration of the convoluted tubules.

Cause of death: Hepatic failure, acute necrosis of liver.

Reg. No. 3834, Non-Booked. Age 26, gravida 2. She was referred here from the Eastern Maternity Hospital in very poor condition. The notes accompanying patient stated "Patient had a low forceps delivery at noon yesterday under ether anaesthesia because of slightly contracted outlet, prolonged labour, and foetal distress. Following the delivery, the vagina was found to have a large tear, and the patient had 30 oz of postpartum haemorrhage. She promptly went into shock. 1 litre of glucose was given. The blood pressure came back to 110/60 mm Hg. The tear was repaired. Four hours later, the patient suddenly became cyanotic and dyspnoeic, B.P. not registrable, pulse rate 150/minute and thready. General examination was otherwise negative. She was given oxygen, glucose infusion, nicamide and adrenaline. Blood pressure gradually returned from 60/30 mm Hg to 110/50 mm Hg. Blood transfusion was given between 11 p.m. and 2 a.m. but was discontinued after giving about 350 cc because the patient developed a temperature of 105°F. At 9 a.m. this morning the pulse rate was 116/minute, still thready. She was still on a 20% glucose drip."

Examination on admission to Tsan Yuk Hospital showed the B.P. 90/60 mm Hg, the pulse rate 128/minute, temperature 101°F, dyspnoea, rales and rhonchi over the left lower chest, tenderness over the right costal margin, flaccid paralysis of both lower extremities, bilateral loss of ankle jerk and knee jerk, absence of Babinski, loss of pain sensation in both lower extremities from L₂ downwards, and very poor general condition. Patient was still conscious. One pint of blood transfusion was given, followed by intravenous dextrose and saline of 3500 cc spread out over a period of 57 hours. The blood pressure came up to 110/60 mm Hg, and the pulse rate to 110 per minute at 7 p.m. Her condition deteriorated again on the following day. She became dyspnoeic and incontinent of urine and faeces. She became drowsy and comatose the 3rd morning and expired at 8.10 p.m. of the 3rd day. Blood pressure and pulse rate remained normal up to last two hours of life.

Postmortem findings: Puerperal uterus with laceration of anterior lip of cervix, repaired; necrosis of pituitary gland; congestion and oedema of lungs; focal necrosis of liver; congestion in medulla of kidneys and fibrous nodule in right kidney.

Reg. No. 6485, Booked. Age 27, gravida 1. L.M.P. 22 January, 1956 and E.D.C. 29 October, 1956. Patient had been in good health all along and attended the antenatal clinic regularly. X-ray pelvimetry at 39th week showed gynaecoid pelvis with slightly contracted measurements.

On 2 November, 1956, at 2 a.m., the patient had sudden onset of constricting pain in the interscapular region. Four hours later, she vomited fresh blood, amount estimated at 800 cc. Patient lost consciousness but recovered spontaneously later. She was brought into hospital at 8.30 a.m. in fairly good condition. B.P. was 110/70 mm Hg and pulse rate 100/minute. Soon after admission, she had another attack of haematemesis of 500 cc and became severely shocked. The abdomen was noticed to be gradually distending. Two and a half pints of blood transfusion were given, following by a nor-adrenaline drip. Her condition improved. B.P. came back to 120/80 mm Hg and radial pulse reappeared at the rate of 110/minute. But she soon had another bout of haematemesis, which was followed by almost continuous vomiting of fresh blood. The abdomen became more and more distended. The patient expired 4½ hours after admission, undelivered.

Postmortem findings: Full-time intra-uterine pregnancy.

Fistula between right side of oesophagus
(about 4 inches below the upper rim
of the larynx and the arch of aorta.)

Extensive haemorrhage into the stomach,
3 lbs of blood clots. The entire length
of the small intestine was filled with
blood.

Cause of death: Exsanguination.

Fistula was thought to have resulted from
a fish bone.

SPECIAL CASE
CASE REPORT OF ABDOMINAL PREGNANCY

Reg. No. 5093, Booked

32 year old, gravida 2, para 1, had one full term normal spontaneous delivery 2 years previously. She first attended antenatal clinic when she was 29 weeks pregnant. Examination then showed thin abdominal wall and easily felt foetal parts, especially around the umbilical region. Foetus was lying transversely with the foetal head just below the umbilicus. X-ray of abdomen showed a single foetus lying transversely and curled up like a ball. Position of foetus remained unchanged during the four subsequent antenatal visits. It was decided to do an Elective Caesarean Section one week before term owing to the thin uterine and abdominal walls. Patient came to hospital when she was 39 $\frac{1}{7}$ weeks pregnant complaining of weak abdominal pains starting some one hour before admission. Examination revealed an average sized foetus lying transversely, the head was felt at umbilicus. Foetal movements could be seen and foetal parts could be distinctly felt through the abdominal wall. No vaginal bleeding and no uterine contractions were detected. Extrauterine pregnancy was suspected and laparotomy performed 12 hours after admission. The uterus was found to be enlarged to the size of a three months pregnancy and a transparent extrauterine sac containing a full term foetus and placenta was found attached to the right side of the fundus of the uterus. A portion of the right Fallopian tube, its fimbria and a flattened ovary were identified on its surface. The sac was paper thin, not connected with the uterine cavity and appeared to be coming from the right cornu of the uterus. The sac was opened. A living normal female infant weighing 3,430 grams (7 lbs. & 9 ozs.) was delivered in excellent condition. The right cornu of the uterus was excised with the sac and placenta. Left uterine tube and ovary were normal. Post operative course was uneventful. Both mother and baby were discharged well on the 14th day after operation.

PAEDIATRIC REPORT

The overall infant loss for the 7,635 Live Births and Still Births is 35.49 per 1,000 Live Births and Still Births. The overall Stillbirth rate is 15.59 per 1,000 Live Births and Still Births. The overall Neonatal rate is 20.04 per 1,000 Live Births.

The Stillbirth rate is 15.03 per 1,000 Live Births and Still Births in the Booked and 87.63 per 1,000 Live Births in the Non-Booked cases. The Neonatal rate is 17.57 per 1,000 Live Births in the Booked and 129.94 in the Non-Booked cases.

(I).

Birth Weight Groups in Grams	Live Birth		Still-birth		Neo-natal Death	
	Booked	Non-Booked	Booked	Non Booked	Booked	Non-Booked
< 1,000	3	1	8	—	2	1
1,001—1,500	26	4	11	6	23	4
1,501—2,000	77	17	19	2	30	6
2,001—2,499	357	20	23	1	36	5
2,500—4,500	6,866	134	39	10	38	7
4,501 & >	10	1	—	—	1	—
Total	7,339	177	100	19	130	23
		7,516		119		153
		7,635*		272		

Note: * This includes 28 babies admitted after delivery. Of these 28 babies, one (#4889/56NB) died on 3rd day because of aspiration of food into bronchi & bronchopneumonia, and one (#7321/56H) was a 1525 gm. macerated foetus. There was no maternal death in the 28 cases and no twins.

(II). Neo-natal Mortality Rate per 1,000 Live Births:—

Birth Weight Groups in Grams	BOOKED		NON-BOOKED	
	No. of Death	Mortality Rate	No. of Death	Mortality Rate
< 1,000	2	666.66	1	1,000.00
1,001—1,500	23	884.61	4	1,000.00
1,501—2,000	30	389.61	6	352.94
2,001—2,499	36	100.84	5	250.00
2,500—4,500	38	5.53	7	5.22
4,501 & >	1	100.00	—	—
TOTAL	130	—	23	—

PAEDIATRIC REPORT (Continued)

2. Causes of Mortality of Live Births after 7 days:—47 babies.

Birth Weight Groups in Grams	B or NB	Bronchopneumonia	Subdural Haemorrhage	Prematurity	Atelectasis	Aspiration of foreign body	Jaundice	Haemorrhage into lungs	Postmortem not done
< 1,000	NB	—	—	—	—	—	—	—	—
1,001 — 1,500	B	4	—	2	1	—	1	—	—
1,501 — 2,000	NB	—	—	1	1	2	1	—	—
2,001 — 2,499	NB	3	3	1	1	—	—	—	—
2,500 — 4,500	NB	1	—	—	2	1	—	1	—
	NB	9	—	—	—	—	—	—	—
	NB	5	3	—	—	—	—	—	—
	NB	1	1	—	—	—	—	—	—
4,501 & >	NB	—	—	—	—	—	—	—	—

B. Still Births: There were 119 Still Births. Postmortem was obtained in 95.0% of the cases.

Birth Weight Groups in Grams	B or NB	*No Abnormality other than Non aerated lungs	Maceration	Anucleated body	Other Focal Inomalities	Perinatal Injury	Subdural Haemorrhage	Atelectasis	Broncho pneumonia	Postmortem not done
< 1,000	B	2	5	—	—	—	—	—	—	1
1,001 — 1,500	NB	—	—	—	—	—	—	—	—	—
	B	2	7	2	—	—	—	—	—	—
1,501 — 2,000	NB	4	1	1	—	—	—	—	—	—
	B	4	5	1	3	2	—	—	—	4
2,001 — 2,499	NB	2	—	—	—	—	—	—	—	—
	B	5	15	1	—	1	—	—	1	—
2,500 — 4,500	NB	21	8	2	4	2	—	1	—	1
	B	6	1	—	1	—	2	1	—	—
	NB	46	42	7	8	5	2	2	1	6
TOTAL										

Note: * Please read following table listing the maternal or foetal complications in connection with the S.B.

PAEDIATRIC REPORT (Continued)

* Table showing maternal or foetal complication in the 46 cases of Stillbirths.

Accidental Haemorrhage (14)	#1818/56NB #2764/56NB #3067/56NB #3941/56B #7606/56B	#2512/56B #2771/56B #3603/56NB #6001/56B #9236/56NB	#2713/56B #2771/56B #4531/56B #6311/56NB
Placenta Praevia (6)	#1706/56NB #3086/56B	#2499/56B #4191/56B	#2842/56NB #4684/56B
Ruptured Uterus (4)	#1453/56NB #6290/56B	#4321/56B	#5374/56B
Prolapsed Cord (4)	#3205/56B #7647/56B	#5028/56B	#6531/56B
Prematurity (1)	#1885/56B		
Shoulder Presentation (1)	#3862/56NB		
Breech (3)	#4135/56NB (prematurity) #391/56B	#6745/56NB #5633/56B	#6575/56B
Pre-eclampsia and/or Hypertension (2)			
Ruptured Splenic Aneurism (1)	#8342/56B		
Prolonged 2nd stage & Instrumental Delivery (2)	#1769/56B	#5804/56B	
Unknown maternal or foetal complications (8)	#1131/56B #3206/56B #7967/56B	#3172/56B #6536/56B #7933/56B	#4802/56B #6803/56B

STATISTICAL SUMMARY

	Booked	Non-booked	Total
Live births	7,339	177	7,516
Stillbirths (over 28 wks gestation)	100	19	119
Stillbirth rate per 1,000 live and stillbirths ..			15.59
Neonatal deaths (all gestations)	130	23	153
Neonatal death rate per 1,000 live and stillbirth			20.04
Neonatal deaths (excluding those with birth weight under 1,000 grams)	128	22	150
Noenatal death rate (excluding those with birth weight under 1,000 grams)			19.65
Premature live births (below 2,500 grams) ..	463	42	505
Deaths among premature live births	91	16	107
Premature mortality rate (% of premature live births)			21.18%
Congenital Malformation:			
Among live births	53	1	54
Among stillbirths	14	1	15
Among premature live births	8	—	8
Birth Trauma:			
Among live births	25	9	34
Among stillbirths	9	2	11
Among premature live births	12	4	16
Infections:			
Among live births	46	5	51
Among premature live births	25	4	29

TABLE 35

STILLBIRTHS

There were altogether 119 stillbirths, or 15.59 per 1,000, 100 in Booked and 19 in Non booked cases.

Reg. No	Birth Weight gm	Maturity of foetus	Sex	Method of Delivery	Maternal factors	Foetal factors	P M Findings (if no P M State, not done)	Remarks
BOOKED								
363	2250	34.4	M	Spont	Accidental Haemorrhage	—	No Advanced Maceration of Internal Organs	
391	2590	35	M	"	Essential Hypertension	—	No Gross Abnormality of Internal Organs	
407	1013	30.1	F	Bipolar Version and Craniotomy	Pulmonary TB Mixed Accidental Haemorrhage	—	Advanced Maceration of Internal Organs	
428	3250	36.2	M	Spont	—	—	No Gross Abnormality of Internal Organs	
591	2650	37.2	F.	"	—	Breech	Hydrops foetalis	
660	1800	33.3	F	Bipolar Podalic Version	Mitral Stenosis and Aortic Regurgitation	(Compound Presentation)	Non-terted Lungs	
662	1850	31.1	M	Spont	P E	Hydranmios	Ictericum of Ictericum Subdural and Subarachnoid Haemorrhage	
836	2400	39.6	F	"	P E	—	Non-terted Lungs	
1130	550	29.1	F	"	—	—	Ictericum of Ictericum Subarachnoid and Subdural Haemorrhage	
1131	2250	37.5	M	"	—	—	Non-terted Lungs	
1260	2700	43	F	"	—	—	Advanced Maceration of Internal Organs	
1336	1040	35.2	F	"	—	Breech	Ictericum of Ictericum Subarachnoid and Subdural Haemorrhage	
1450	800	32.1	M	"	—	—	Non-terted Lungs	
1663	2041	33.3	M	"	—	—	Anencephalic Foetus	
1769	3175	42.1	F	L.F.	Accidental Haemorrhage	—	Advanced Maceration	
1851	1620	29.6	M	Spont	—	Breech	Macerated Foetus	
							No Other Findings	
							Hydrops foetalis	
							Non-terted Lungs	

STILLBIRTHS (Continued)

Reg. No.	Birth Weight gm.	Maturity of foetus	Sex	Method of Delivery	Maternal factors	Foetal factors	P. M. Findings (if no P. M. State "not done")	Remarks
1885	726	28	F.	Spont.	---	---	No Gross Abnormality of Internal Organs	
2268	3780	40-1	M.	M.F.	---	O.P.	Laceration of Tentorium Massive Cephalohaematoma Subarachnoid and Subdural Haemorrhage	
2286	2370	32-6	F.	Spont.	P.F.	---	Macerated Foetus	
2308	2520	29-2	M.	"	---	---	Anencephalic Foetus	
2323	3570	45	M.	"	---	L.M.A.	Macerated Foetus	
2359	1700	35-6	M.	Manipulative Delivery	---	---	Laceration of Tentorium	
2363	2500	35	F.	Spont.	---	---	Anencephalic Foetus	
2374	1790	34-1	F.	"	---	---	Not Done	
2388	3340	44-1	M.	"	---	R.M.P.	Cephalohaematoma	
2440	2550	35-4	M.	"	---	---	Non aerated Lungs	
2499	3175	37-6	M.	Willitt's Forceps	Placenta Praevia III	---	Non aerated Lungs	
2512	3525	44-1	M.	Spont.	Concealed	---	Anencephaly	
2534	1750	34-4	M.	"	Accidental Haemorrhage	L.M.A.	Haemorrhage in Lungs	
2587	2280	35-3	F.	"	---	Hydramnios	No Gross Abnormality of Internal Organs	
2713	3500	37-5	F.	"	Accidental Haemorrhage	---	"	
2771	{ 1899 1701 }	35-5	{ M. M. }	"	---	L.O.P. → P.O.P. Uniovular Twin Breech Pregnancy	Non aerated Lungs	
2795	1400	?	F.	"	---	---	Macerated Foetus	
2845	2800	39	M.	"	---	Breech	Non-aerated Lungs	
3086	2722	36-4	M.	L.S.C.S.	Placenta Praevia IV	---	Macerated Foetus	
3172	2000	36	M.	Willitt's Forceps	---	---	Non-aerated Lungs	
3205	2700	41-1	F.	Spont.	---	Prolapse of Cord	Non-aerated Lungs	
3278	3400	41-2	F.	"	---	Binovular Twins Pregnancy—1st Twin Uniovular Twins	No Other Abnormality Found	
3404	3120	38-5	F.	"	---	Pregnancy—2nd Twin Prolapsed Cord	Complete Atelectasis (Bil.) No Other Abnormality Found	
3578	3030	?	M.	L.F.	---	Face Presentation	No Other Abnormality	
3621	1000	34-3	---	"	---	Uniovular Twins Pregnancy—2nd Twin Papyraceous	Hydrops Foetalis Macerated Foetus	
3643	2180	38-2	F.	"	---	Face Presentation	Anencephaly, Atelectasis Subpleural Haemorrhage	

STILLBIRTHS (Continued)

Reg. No.	Birth Weight gm.	Maturity of foetus	Sex	Method of Delivery	Maternal factors	Foetal factors	P. M. Findings (if no P. M. State 'not done')	Remarks
3844	2100	35.5	F.	Spont.	---	---	Macerated Foetus	
4092	2050	40.2	M.	"	---	---	"	
4109	2700	39.2	M.	"	---	---	"	
4136	3250	40.2	F.	"	---	---	Non-aerated Lungs No Other Abnormality	
4138	2400	?	F.	Embryotomy- Perforation	---	---	Macerated Foetus	
4191	900	26.4	M.	I. S. C. S.	Placenta Praevia IV	---	Non-aerated Lungs	
4204	2650	41.5	M.	Spont.	---	Uniovular Twin	Non-aerated Lungs No Other Abnormality	
4244	1700	38.4	M.	"	---	Pregnancy—1st Twin	Macerated Foetus	
4321	3270	39.6	F.	I. S. C. S.	Incomplete Rupture of Uterus (Contracted Pelvis)	---	Non-aerated Lungs No Other Abnormality	
4391	1800	35.4	M.	Spont.	---	Breech	Hydrocephalus	
4531	2920	36.2	M.	"	Accidental Haemorrhage	---	Non-aerated Lungs	
4678	510	30	M.	"	---	---	Non-aerated Lungs	
4679	680	30.5	F.	"	---	---	Subpleural Haemorrhage (Bil.)	
4684	3178	36	F.	"	Placenta Praevia I	---	Not Done	
4892	2850	43.4	M.	"	---	---	Macerated Foetus	
4901	2600	37.3	M.	"	---	---	Non-aerated Lungs	
4901	1800	37.3	M.	"	---	---	No Other Abnormality	
5028	2200	35.6	M.	Internal Version Breech Extraction	---	Uniovular Twin	Maceration	
5189	2300	?	F.	Spont.	---	1st Twin 2nd Twin	"	
5206	1200	37	F.	"	---	Pregnancy Shoulder Presentation Prolapsed Left Arm & Cord	Non-aerated Lungs No Other Abnormality	
5374	2440	35.3	M.	"	Rupture of Uterus	Breech	Maceration	
5410	1550	34.1	F.	Laparotomy for Ruptured Uterus and Subtotal Hysterectomy	---	---	Non-aerated Lungs No Other Abnormality	
5633	3730	41.5	M.	Spont.	P.F.	Breech	Maceration	
5800	2180	32	M.	"	---	Breech, Hydramnios	Non-aerated Lungs No Other Abnormality	
5804	3430	40	M.	M.R. + M.F.	---	---	Maceration	
5842	3170	36.5	F.	Spont.	---	Shoulder Presentation	Non-aerated Lungs No Other Abnormality	
5852	1200	33.5	F.	"	---	---	Maceration	

STILLBIRTHS (Continued)

Reg. No.	Birth Weight gm.	Maturity of foetus	Sex	Method of Delivery	Maternal factors	Foetal factors	P. M. Findings (if no P. M. State "not done")	Remark.
5941	2465	35-5	M.	Spont.	Accidental Haemorrhage	---	Non-aerated Lungs No Other Abnormality	
6001	2200	38-3	M.	"	"	Breech	Maceration	
6023	2700	30-2	M.	Subtotal Hysterotomy	Rupture of Uterus	---	Non-aerated Lungs No Other Abnormality	
6290	3459	39-2	M.	"	Contracted Pelvis, P.P.H.	Breech	Maceration	
6468	1005	27-4	M.	Spont.	---	Prolapscd Cord	Non-aerated Lungs No Other Abnormality	
6531	3410	39	F.	"	---	---	"	
6536	3540	43	M.	"	---	---	"	
6575	2750	39-6	M.	"	---	Breech	Maceration	
6648	3200	42	M.	"	---	---	"	
6731	1500	27-1	M.	"	V.D.R.L.:Positive	Breech	"	
6735	700	30-3	F.	"	---	---	"	
6795	2190	34-2	F.	"	---	---	"	
6803	1500	39-6	F.	"	---	---	"	
6825	1800	35	M.	"	"U": Shaped Uterus	---	Non-aerated Lungs No Other Abnormality	
6880	1680	?	F.	"	---	---	Maceration	
6911	2400	39	M.	"	---	---	Not Done	
7126	1650	35	M.	"	---	---	Maceration	
7321	1525	39	F.	"	---	---	Not Done	
7358	1455	38-5	F.	"	---	---	Maceration	
7457	3350	36-2	F.	"	---	---	Anencephaly	
7531	2280	36-1	M.	P.E.	---	---	---	
7606	2600	35-6	M.	C.C.S.	Accidental Haemorrhage	---	Maceration No Other Abnormality Non-aerated Lungs	B.B.A.
7774	1780	35	M.	Spont.	---	Breech	Maceration	
7838	1700	33	F.	"	---	"	"	
7873	1400	29-4	F.	Internal Version	V.D.R.L.:Positive Placenta Praevia III	"	"	
7895	2400	37-3	F.	Internal Version Breech Extraction	---	Shoulder Presentation	"	
7967	1800	42-1	M.	Spont.	---	---	Non-aerated Lungs No Other Abnormality	
7647	2680	38-2	F.	"	---	Prolapscd Cord	"	
7933	2700	37-1	M.	"	Contracted Pelvis Intra-Peritoneal Haemorrhage	---	"	
8242	3500	39-3	M.	"	---	---	"	
NON-BOOKED								
1453	2500	43-5	M.	L.S.C.S.	Rupture of Uterus	---	Non-aerated Lungs	
1548	1480	39-1	M.	Spont.	Pulmonary T.B.	---	Anencephalic Foetus	
1700	1474	?	M.	Bipolar Version	Placenta Praevia III	Shoulder Presentation	Non-aerated Lungs	

STILLBIRTHS (Continued)

Reg. No.	Birth Weight gm.	Maturity of foetus	Sex	Method of Delivery	Maternal factors	Foetal factors	P. M. Findings (If no P. M. State "not done")	Remarks
1818	2700	41	F.	Willett's Forceps	Accidental Haemorrhage	—	No Gross Abnormality of Internal Organs	
2710	1300	30	M.	Spont.	Accidental Haemorrhage	Hydramnios	Macerated Foetus	
2764	2560	29	M.	"	Concealed	—	Non-acrated Lungs	
2842	1770	32	M.	"	Accidental Haemorrhage	Prolapsc of Cord	"	
3067	1250	?	M.	"	Placenta Praevia II	—	No Other Abnormality	
3323	3400	40.6	M.	"	Accidental Haemorrhage	—	Subdural Haemorrhage	
3599	2700	42.6	F.	L.S.C.S.	—	Conjoined Twins, Hydramnios	Atelectasis (Bil) Not Done	
3603	3600	40.3	M.	Spont.	Accidental Haemorrhage	—	Non-acrated Lungs	
3605	2588	40.4	M.	Internal Version	P. E.	Shoulder Presentation	Subpleural Haemorrhage (Bil.)	
3782	2400	39.5	F.	Breech Extraction L.S.C.S.	P. E.	—	Subdural Haemorrhage	
3862	1250	?	F.	Bipolar Version	Placenta Praevia III	Shoulder Presentation	Complete Atelectasis (Bil.)	
4135	1450	34.6	M.	Spont.	P.L.	Breech	No Other Abnormality	
6311	1300	32.1	M.	C.C.S.	Accidental Haemorrhage	—	Subdural Haemorrhage	
6649	2551	40.4	M.	Norton's Extra- Peritoneal Caesarean Section	Fibroid of Cervix	Shoulder Presentation	Prematurity	
6745	2800	39.1	F.	Spont.	Contracted Pelvis	Breech	Non-acrated Lungs	
8236	3400	40	M.	L.S.C.S.	Accidental Haemorrhage	—	No Other Abnormality	

TABLE 36
NEO-NATAL DEATHS

There were altogether 153 neo-natal deaths, or 20.03 per 1,000, 130 in Booked and 23 in Non-booked cases.

Reg. No.	Birth Weight gm.	Maturity foetus	Sex	Method of Delivery	Maternal factors	Complications in infant	P. M. Findings (if no P. M. State "not done")	Age of death	Method of feeding	Remarks
BOOKED										
101	2890	38.5	F.	L.S.C.S.	—	Breech	Atelectasis with Aspiration of Foreign Body into Lungs	4 days	Breast	
113	1700	27.4	F.	Spont.	Mitral Stenosis	—	Atelectasis of Lungs	2 "	Artificial	
149	2880	33.1	M.	"	—	—	Severe Prematurity	3 "	"	
198	1500	30.3	F.	"	—	—	Torsion of Tentorium	25 "	"	
215	2200	36.1	M.	"	P.E.	—	Torsion of Tentorium	12 hrs.	—	
291	2200	39.5	F.	"	"	—	Laceration of Tentorium	1 day	—	
340	2800	40	M.	"	—	Shoulder Presentation	"	3 days	Artificial	
350	1820	37.1	F.	"	P.E.	Breech	Subdural Haemorrhage	10 "	"	
506	1250	30	M.	L.S.C.S.	Placenta Praevia III	—	Torsion of Tentorium	5 hrs.	—	
623	1870	35	M.	Spont.	—	—	Atelectasis of Lungs	3 days	Artificial	
674	1000	28.3	F.	"	Placenta Praevia I	—	"	7 "	"	
790	2950	38.4	M.	"	—	Face Presentation	Anencephaly	4 hrs.	—	
794	2890	38.1	M.	"	Pulmonary T.B.	—	Bronchopneumonia	12 days	Artificial	
881	2960	45.5	M.	M.R. + M.F.	I.P. Eclampsia	R.O.T.	Aspiration of Foreign Body	3 "	"	
967	2000	35	F.	Spont.	Mitral Stenosis	—	Haemorrhage into Lungs	20 "	"	
1075	3130	39.2	F.	"	—	—	Atelectasis of Lungs	4 "	"	
1178	2100	33.4	F.	"	—	—	"	9 "	"	
1534	1900	34	M.	"	—	—	"	3 "	"	
1735	3750	40.4	M.	"	—	—	Tentorial Tear	7 "	"	
1719	2750	40.6	M.	L.S.C.S.	Contracted Pelvis	L.O.P.	Atelectasis of Lungs	3 "	"	
2075	3090	41.2	M.	M.R. + M.F.	—	O.P.	Tentorial Tear	4 "	"	
2180	3000	37.5	M.	Spont.	—	Twins	Hydrocephalus	3 "	Breast	1st Twin
2232	7460	32.4	M.	Breech	—	Pregnancy	Atelectasis	4 "	Artificial	2nd Twin
			M.	Assisted	—	—	"	2 "	"	
2236	2800	41.5	M.	Spont.	Pulmonary T.B.	—	Haemorrhage into Lungs	2 "	"	2nd Twin
2355	2020	35.6	M.	"	—	Rhinovular Twins	Atelectasis	7 "	Breast	
2505	1750	33.5	M.	"	—	—	Prematurity	7 "	"	
2542	3300	37.1	M.	Spont.	—	Pregnancy	Severe Jaundice	5 "	Breast and Artificial	1st Twin
	3050	37.2	M.	M.R. + M.F.	—	Uniovular Twins	Tentorial Tear	8 "	"	2nd Twin
2657	2500	33.5	M.	L.S.C.S.	Placenta Praevia III	Pregnancy	Hydrocephalous	2 hrs.	—	
2706	1250	31.2	F.	Spont.	—	—	Prematurity	3 days	Artificial	

NEO-NATAL DEATHS (Continued)

Reg. No.	Birth Weight gm.	Maturity of foetus	Sex	Method of Delivery	Maternal factors	Complications in infant	P. M. Findings (if no P. M. State "not done")	Age of death	Method of feeding	Remarks
2826	1900	34.1	F.	Spont.	---	---	Atelectasis of Lungs	9 hrs.	---	
2882	1160	29.4	F.	"	---	---	"	22 days	Artificial	
2927	2790	42.2	F.	"	Mitral Stenosis	Breech	Atelectasis of Lungs	3 days	"	
2991	2320	39	F.	"	---	---	Bronchopneumonia	11 "	"	
3025	1280	28.3	F.	"	---	---	Atelectasis of Lungs	1 day	---	
3065	2350	34.5	F.	"	---	---	Hæmorrhage into Lungs	6 days	Artificial	
3139	3050	40.2	M.	M.F.	P.F.	---	Aspiration of Foreign Body into Lungs	4 "	"	
3161	2600	40.1	M.	Spont.	---	---	Atelectasis of Lungs	6 "	---	
3165	3000	41.2	M.	"	Anæmia	---	"	5 hrs.	---	
3171	2850	43	M.	"	---	---	"	6 days	Breast	
3176	1500	35.4	F.	"	---	---	"	9 hrs.	---	
3189	2320	38.6	F.	"	---	---	"	8 days	Breast and Artificial	
3317	1970	38.5	M.	"	---	Breech	Bronchopneumonia	4 "	"	
3442	2680	39.1	F.	"	---	Breech	"	5 "	"	
3532	1420	28.5	F.	L.S.C.S.	Placenta Praevia III	---	Atelectasis of Lungs	7 hrs.	"	
3543	1300	28.3	M.	Spont.	---	Uniovular Twins	Prematurity	2 days	---	
3621	2630	34.3	M.	"	---	Pregnancy	Bronchopneumonia	4 "	Breast and Artificial	1st Twin
3661	3150	40.1	M.	"	Contracted Pelvis	---	Atelectasis of Lungs	3 hrs	"	
3667	960	29.1	M.	"	---	---	"	3 d ys	---	
3736	1830	31	M.	"	---	---	"	10 "	Artificial	
3835	2650	41.5	M.	"	---	Hydræmios	Atresia of Oesophagus, Oesophageo tracheal Fistula	10 hrs	"	
3905	1980	36.5	M.	"	---	Uniovular Twins	Jaundice	9 d ys	Artificial	2nd Twin
4034	1190	29.5	M.	"	---	Pregnancy	Prematurity	2 "	---	1st Twin
	1240		M.	"	---	Uniovular Twins	Subdural Hæmorrhage	2 "	---	2nd Twin
4099	2080	40	M.	"	---	---	"	5 "	Artificial	
4158	1800		F.	"	---	---	Jaundice	3 "	"	
4205	2080	43.3	M.	"	---	---	Bronchopneumonia	17 "	"	
4221	2150	40.3	F.	"	---	---	Prematurity	3 "	"	
4234	2260	34.4	F.	"	---	---	Bronchopneumonia	18 "	"	
4256	2000	34	F.	"	---	---	Prematurity	19 "	"	
4326	1785	31	M.	"	---	---	Bronchopneumonia	9 "	"	
4344	2950	41.4	M.	"	Pulmonary T.B.	---	"	3 hrs	"	
4354	2770	42.6	F.	"	---	---	"	11 days	Artificial	

NEO-NATAL DEATHS (Continued)

Reg. No.	Birth Weight gm.	Maturity of foetus.	Sex	Method of Delivery	Maternal factors	Complications in infant	P. M. Findings (if no P. M. State "not done")	Age of death	Method of feeding	Remarks
4413	1760	32-6	M.	Spont.	—	—	Bronchopneumonia	1 day	—	1st Twin
4460	1300	35-5	F.	"	—	Uniovular Twins	Prematurity	11 days	Artificial	2nd Twin
	1660		F.	"	—	Pregnancy	Aspiration of Foreign Body	"	"	
4516	1910	34-4	M.	"	—	Uniovular Twins	Bronchopneumonia	17 "	"	1st Twin
4570	1770	44-4	F.	"	—	Pregnancy	Bronchopneumonia	11 days	"	2nd Twin
	2110		F.	"	—	—	"	6 "	"	Breech
4585	1260	36	M.	"	—	—	Atelectasis of Lungs	13 hrs.	—	
4658	1200	26-2	F.	"	—	—	Bronchopneumonia	12 days	—	
4670	1730	38-1	M.	"	—	Breech	Atelectasis of Lungs	5 hrs.	—	
4741	2410	36-1	M.	"	—	—	Subdural Haemorrhage	1 day	—	
4797	1900	39-5	F.	Internal Version	—	Binovular Twins	Tentorial Tear	20 hrs.	—	2nd Twin
4879	1900	35-5	M.	Spont.	—	Pregnancy	Not Done	7 days	Artificial	
4958	3200	41	F.	M.F.	—	—	Subdural Haemorrhage	15 "	"	
5020	1820	34-4	M.	Spont.	—	Breech	"	1 day	—	
5090	2420	36-2	M.	"	Cervical Polyp	—	"	7 days	Breast	
5163	2085	41-1	M.	"	—	—	Cephalo-Haematoma	4 "	Artificial	
5200	2130	36-1	M.	"	—	Uniovular Twins	Not Done	1 hr.	—	1st Twin
5231	2280	37	F.	"	—	Pregnancy	Bronchopneumonia	16 days	Artificial	
5240	1180	29	F.	"	Placenta Praevia II	Breech	Subdural Haemorrhage	2 hrs.	—	
5246	1400	31-1	F.	Spont.	—	Prolapsed Cord	Bronchopneumonia	6 days	Artificial	1st Twin
	1440		F.	"	—	Uniovular Twins	"	14 "	"	2nd Twin
5330	2280	39-4	F.	M.R. + M.F.	Mitral Stenosis	—	Not Done	11 "	Breast	
5310	3150	43	F.	Spont.	—	Unilateral Harelip (L.)	Aspiration of Foreign Body into Lungs	4 "	Artificial	
5520	2268	36-6	M.	"	—	Hypoplasia	"	11 "	"	
5526	1600	32-2	M.	"	—	—	Atelectasis of Lungs	1 day	—	
5599	2300	32-5	M.	L.S.C.S.	Contracted Pelvis	—	Haemorrhage into Lungs	16 days	Artificial	
5600	2320	34-6	F.	Spont.	—	Uniovular Twins	Bronchopneumonia	11 "	"	
5650	2400	34-4	M.	"	P.E.	Pregnancy	Subdural Haemorrhage	4 "	"	1st Twin
5663	2570	?	M.	C.C.S.	Placenta Praevia II	—	Bronchopneumonia	12 "	"	
5670	2150	33-5	F.	L.S.C.S.	—	—	"	9 "	"	
5686	2005	31-3	M.	Spont.	Pulmonary T.B.	—	"	11 "	"	
5777	1330	?	M.	"	—	Binovular Twins	"	9 "	"	1st Twin
						Pregnancy	"	"	"	

NEO-NATAL DEATHS (Continued)

Reg. No.	Birth Weight gm.	Maturity of foetus	Sex	Method of Delivery	Maternal factors	Complications in infants	P. M. Findings (if no P. M. State "not done")	Age of death	Method of feeding	Remarks
5777	1390	?	M.	Spont.	—	Binovular Twins Pregnancy	Bronchopneumonia	14 days	Artificial	2nd Twin Shoulder Presentation
5793	2140	35-2	M.	"	—	—	"	6 "	"	
5852	3250	39	F.	L.S.C.S.	—	Breech	"	20 "	Breast and Artificial	
5854	2300	36.6	M.	Spont.	—	—	Not Done	13 "	Artificial	
5966	2360	35-6	M.	Spont.	—	Breech	Bronchopneumonia	7 days	"	
5989	2230	36-4	M.	"	—	—	"	12 "	"	
6003	2500	36	M.	"	—	—	"	8 "	"	
6117	1700	34-6	M.	"	—	Uniovular Twins	Bronchopneumonia	5 "	"	1st Twin
	2100		M.	"	—	Pregnancy	Subdural Haemorrhage	6 "	"	2nd Twin
6205	2260	35	F.	"	V.D.R.L. Positive	—	Prematurity	2 "	Artificial	
6389	2050	36	F.	"	"	—	Subdural Haemorrhage	2 "	"	
6435	1160	35-6	F.	L.S.C.S.	Contracted Pelvis Placenta Praevia IV	Breech	Atelectasis of Lungs	3½ hrs	"	
6472	2050	37-4	M.	Spont.	—	—	Subdural Haemorrhage	1 day	"	
6678	1760	32-3	M.	"	—	—	"	1 "	Artificial	
6755	1990	36-3	F.	"	—	—	"	6 days	"	
6765	1560	35-6	M.	"	—	—	"	3 "	Breast	
6790	2650	40-6	M.	"	—	—	Not Done	2 "	"	
6868	3400	38-7	F.	P.E.	P.E.	—	Subdural Haemorrhage	3 "	Artificial	
7124	3005	40-1	F.	Spont.	—	—	Atelectasis of Lungs	2 "	Artificial	
7197	2410	37-2	F.	"	—	—	Tenorial Tear	4 "	"	
6963	5370	37	M.	M.F.	Hydramnios	Breech	Jaundice	16 "	Breast	
7322	1446	31-2	F.	L.S.C.S.	P.E.	—	Prematurity	9 hrs	Artificial	
7426	1350	32-2	F.	Spont.	—	—	Subdural Haemorrhage	2 days	"	
7519	2000	40-1	M.	"	—	—	Bronchopneumonia	11 "	"	
7592	1200	30-2	F.	L.S.C.S.	Accidental Haemorrhage	Breech	"	"	"	
7714	2450	37	M.	Spont.	—	—	Atelectasis of Lungs	2 "	Breast	
7845	4500	38-2	F.	L.S.C.S.	—	Conjoint Twins	Monster	32 hrs.	"	
7938	1550	36	F.	Spont.	—	—	Subdural Haemorrhage	21 days	Artificial	
7944	3860	38-3	M.	C.C.S.	—	—	"	7 "	"	
8061	4005	44-4	F.	Spont.	Naegle's Pelvis	Breech	Not Done	3 "	"	
8072	3009	43-6	M.	L.F.	A.P.—F. clampsia	—	Bronchopneumonia	13 "	"	
8077	1630	38-4	F.	Spont.	—	Uniovular Twins Pregnancy	Subdural Haemorrhage	18 "	"	2nd Twin
8287	2200	?	M.	"	P.E.	—	Bronchopneumonia	9 "	"	
8301	2400	37-4	F.	"	—	—	Subdural Haemorrhage	3 "	Breast	

NEO-NATAL DEATHS (Continued)

Reg. No.	Birth Weight gm.	Maturity of foetus	Sex	Method of Delivery	Maternal factors	Complications in infant	P. M. Findings (if no P. M. State "not done")	Age of death	Method of feeding	Remarks
NON-BOOKED										
102	1150	27-3	M.	Spont.	—	Hydramnitis	Atelectasis of Lungs	4 days	Breast	
792	930	30-4	F.	—	P.E. Anaemia	—	Aspiration of Foreign Body into Lungs	4 "	Artificial	
913	2550	42-5	M.	M.R. † M.F.	—	—	Tentorial Tear	1 hr.	—	
1140	3000	39	M.	L.F.	—	—	Atelectasis of Lungs	1 day	—	
1707	1650	37-1	M.	Spont.	—	—	Tentorial Tear	8 hrs.	—	
1925	1580	32	M.	—	—	Breech	"	1 day	—	
1932	2020	37	M.	—	—	O.P.	"	4 days	Breast	
2316	3062	36-3	M.	M.R. † M.F.	P.E.	Breech	Tentorial Tear	4 days	Artificial	
2566	1720	32-4	M.	L.S.C.S.	Placenta IV Placenta II	Breech	Subdural Haemorrhage Not Done	19 "	—	
2610	1840	32-4	F.	"	—	Prolapse of Cord	Prematurity	4 days	Artificial	
2766	1080	28-4	M.	Spont.	—	Shoulder Presentation	Tentorial Tear	2 hrs.	—	
2837	1950	40-1	F.	Internal Version, Breech Extraction	P.E.	Breech Twin Pregnancy	"	2 days	—	2nd Twin
3085	2300	?	F.	Spont.	Acute Necrosis of Liver	—	Atelectasis of Lungs	3 "	Artificial	
3704	2400	35-3	M.	M.F.	Contracted Pelvis	—	Subdural Haemorrhage	3 "	"	
4245	1380	35-1	F.	Spont.	—	Breech	Prematurity	19 "	"	
4889	2400	?	M.	"	—	—	Bronchopneumonia	3 "	"	
5476	2270	33	M.	"	V.D.R.L. Positive	—	"	4 "	"	
6252	2130	21	M.	"	—	Breech	"	19 "	"	
6958	3200	?	F.	L.S.C.S.	—	Brow Presentation	Tentorial Tear	19 hrs.	—	
7081	1620	31	F.	Spont.	Accidental Haemorrhage	—	Prematurity	4 hrs.	—	
7988	2750	37	M.	Spont.	—	Binocular Twins	Bronchopneumonia	14 days	Artificial	1st Twin
	2700		M.	"	—	Pregnancy Shoulder Presentation	Not Done	3 "	"	2nd Twin
7999	1800	34-5	F.	C.C.S.	Accidental Haemorrhage	Prolapse of Lt. Arm	Bronchopneumonia	18 "	Breast and Artificial	

TABLE 37

PREMATURE LIVE-BORN INFANTS

There were altogether 505 Premature Live Births weighing less than 2,500 grams (5½ lbs.), 463 Booked and 42 Non-booked cases.

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
BOOKED										
6958/55	2400	39.3	F.	Unknown	---	Breast	2495	18 days	L.	
1/56	2350	38.4	F.	"	---	"	2160	10 "	L.	
2	1920	35.3	F.	"	---	Artificial	2510	21 "	L.	
22	2450	39	F.	"	---	Breast	2523	7 "	L.	
51	2300	36.4	F.	"	---	Artificial	2250	5 "	L.	
38	2120	37.3	M.	"	---	Breast	2460	15 "	L.	
44	2450	40	F.	"	---	"	2381	6 "	L.	
69	2370	39.5	F.	"	---	Artificial	2200	9 "	L.	
72	2300	37.4	F.	"	---	"	2300	23 "	L.	
91	2268	39.5	F.	"	---	Breast & Artificial	2410	6 "	L.	
98	2140	34.5	M.	Pulmonary T.B.	---	Artificial	2640	34 "	L.	
113	1700	27.4	F.	Mitral Stenosis	---	Artificial	1600	2 "	N.D.	
118	2402	36.2	F.	Unknown	---	Breast	2402	12 "	L.	
126	2420	40.5	F.	"	---	"	2495	7 "	L.	
145	1510	40.1	M.	"	---	Artificial	2120	60 "	L.	
198	1500	30.3	F.	Pulmonary T.B.	---	"	1080	25 "	N.D.	Transferred to Q.M.H.
208	1950	?	M.	Unknown	---	"	2650	42 "	L.	
215	2200	36.1	M.	"	---	"	---	12 hrs.	N.D.	
230	1850	32.6	M.	P.E.	---	"	2560	37 days	L.	
230	1950	32.6	M.	"	---	"	2500	34 "	L.	
248	2220	38.3	F.	Twin Pregnancy	---	"	2500	27 "	L.	
264	2200	34	M.	Unknown	---	"	2520	22 "	L.	
274	2420	38.1	F.	"	---	Breast	2498	7 "	L.	1st Twin
279	2350	39.5	M.	"	---	"	2286	5 "	L.	2nd Twin
285	2020	34	M.	"	---	Breast	2510	35 "	L.	
291	2200	39.5	F.	P.E.	Laceration of Tentorium	Artificial	---	1 day	N.D.	
343	2450	34.4	F.	Unknown	Mild Subdural Haemorrhage	Breast	2381	7 days	L.	
350	1820	27	F.	P.E.	---	Artificial	1720	10 "	N.D.	
393	2400	34.6	M.	"	---	"	2480	24 "	L.	
395	2470	35.4	F.	"	---	"	2400	19 "	L.	
416	1850	37.1	M.	"	---	"	2570	48 "	L.	
480	2460	38.5	M.	"	---	Breast	2381	9 "	L.	

PREMATURE LIVE-BORN INFANTS, (Continued)

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
506	1250	30	M.	Placenta Praevia Type III	—	Artificial 2 hrly.	—	5 hrs.	N.D.	
520	2150	38-3	F.	Mitral Stenosis	—	" 2 "	2500	21 days	L	
542	2300	43-5	M.	Unknown	—	" 2 "	2540	22 "	L	
543	2370	37-1	F.	P.E.	Cephalo-haematoma	" 2 "	2520	24 "	L	
548	2110	37-1	F.	Bronchial Asthma	—	" 2 "	2500	26 "	L	
577	2420	40-1	F.	Unknown	—	Breast 3 "	2353	6 "	L	
580	2480	43	F.	"	—	" 4 "	2381	7 "	L	
590	2400	39-4	F.	"	—	" 4 "	2410	5 "	L	
599	2460	41	F.	"	—	Artificial 3 "	2610	23 "	L	
612	2480	37-3	F.	"	—	Breast & Artificial 3 "	2495	15 "	L	
617	2480	40	F.	"	—	Artificial 3 "	2768	15 "	L	
623	1870	35	M.	"	—	Breast 4 "	1760	3 "	N.D.	2nd Twin
649	2100	31-2	F.	P.E., Twin Pregnancy	—	Artificial 2 "	2590	33 "	L	
650	2380	28-3	F.	Unknown	—	Artificial 3 "	2353	4 "	L	
674	1000	35-5	F.	Placenta Praevia Type I	—	Breast 4 "	800	7 "	N.D.	
687	2420	37	F.	Unknown	—	Artificial 2 "	2560	32 "	L	
707	2470	37	M.	"	—	" 3 "	2410	7 "	L	
735	2000	38	M.	"	—	Breast 3 "	2530	34 "	L	
748	2360	38-2	F.	"	—	Artificial 2 "	2460	24 "	L	
771	2390	40	F.	P.E.	—	" 2 "	2580	26 "	L	
784	2450	38-4	F.	Unknown	—	Artificial 3 "	2466	12 "	L	
797	1570	33	F.	"	—	Breast 3 "	2650	52 "	L	
831	2390	37-6	M.	"	—	Artificial 2 "	2550	30 "	L	
855	1500	34-5	F.	"	—	" 2 "	2600	62 "	L	
862	2430	42-1	F.	Avitaminosis B ₂	—	Breast 4 "	2460	6 "	L	
904	2300	37-3	F.	Unknown	—	Artificial 3 "	2807	39 "	L	
905	2430	41	M.	"	—	Breast 2 "	2551	8 "	L	
966	2466	33-2	M.	"	—	" 2 "	2523	13 "	L	
967	2000	35	F.	Mitral Stenosis	—	Artificial 2 "	—	20 "	N.D.	
1014	2330	37-1	M.	Unknown	—	" 2 "	2480	23 "	L	
1033	2490	37-2	M.	"	—	" 2 "	2693	24 "	L	
1060	2190	39-3	F.	P.E.	—	" 2 "	2775	32 "	L	
1088	2450	39-6	M.	Unknown	—	Breast 4 "	2432	7 "	L	
1164	1870	31-2	M.	"	—	Artificial 3 "	2550	45 "	L	
1178	1750	33-4	F.	"	—	Artificial 2 "	2320	42 "	L	1st Twin
	2100	35-4	F.	Twin Pregnancy	—	Artificial 2 "	—	9 "	N.D.	2nd Twin
1198	2120	35-1	F.	Unknown	—	Artificial 3 "	2590	29 "	L	
1261	2150	36-5	F.	P.E.	—	" 3 "	2600	24 "	L	
1279	2110	41-6	F.	Twin Pregnancy	—	Artificial 2 "	2500	27 "	L	1st Twin
	2360	41-6	M.	"	—	Artificial 2 "	2530	24 "	L	2nd Twin
1348	2360	32-5	M.	Unknown	—	Artificial 3 "	2370	20 "	L	
1448	2470	36-3	F.	Pulmonary T.B.	—	" 3 "	2480	16 "	L	

PREMATURE LIVE-BORN INFANTS (Continued)

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
1455	2480	38-6	F.	P.E.	—	Breast 4 hrly.	2466	15 days	2	
1460	2080	39-4	M.	Unknown	—	Artificial 2 "	2530	20 "	L	
1481	2450	44-4	F.	"	—	Artificial 2 "	2500	6 "	L	
1491	2120	37-3	M.	Twin Pregnancy	—	Artificial 2 "	2500	21 "	L	2nd Twin
1504	1990	35	F.	Eclampsia	—	Artificial 2 "	2620	42 "	L	
1534	1900	34	M.	Unknown	—	" 2 "	2430	6 "	N.D.	
1587	2440	36-3	M.	Accidental Haemorrhage	—	Breast 2 "	2360	7 "	L	
1617	2466	39-1	M.	Unknown	—	Artificial 3 "	2620	31 "	L	
1662	2240	39-2	F.	P.E.	—	Artificial 3 "	2620	31 "	L	
1697	2400	38-2	M.	Bronchiectasis	—	Breast 3 "	2608	9 "	L	
1759	2310	39-5	F.	Unknown Pulmonary T.B.	—	Breast & Artificial 2 "	2520	11 "	L	
1771	2396	39-3	F.	—	—	Breast 2 "	2480	8 "	L	
1844	2200	38-6	F.	Unknown	—	Artificial 3 "	2500	22 "	L	
1899	1970	31-1	F.	P.E.	—	Artificial 3 "	2540	40 "	L	
1914	2400	40-4	F.	Unknown	—	" 3 "	2560	16 "	L	
1918	2460	34-4	M.	"	—	" 3 "	2520	18 "	L	
1949	2450	35-5	F.	"	—	Breast 3 "	2325	7 "	L	
1981	2490	38-6	F.	Pulmonary T.B.	—	Artificial 3 "	2580	17 "	L	
2055	2400	39-1	F.	Unknown	—	Breast 3 "	2325	5 "	L	
2074	2050	36-2	F.	"	—	Breast 3 "	2550	31 "	L	
2090	2320	44-3	F.	"	—	Artificial 3 "	2540	30 "	L	
2176	2300	39-4	F.	"	—	Artificial 3 "	2500	13 "	L	
2184	2140	36-6	F.	P.E.	—	Artificial 3 "	2350	48 "	L	1st Twin
2225	2420	38-2	F.	Twin Pregnancy	—	Breast 4 "	2490	13 "	L	2nd Twin
2232	1520	32-4	M.	Unknown	—	Artificial 2 "	—	4 "	N.D.	1st Twin
2250	1460	38-3	M.	Mitral Stenosis	—	Artificial 2 "	—	2 "	N.D.	2nd Twin
2258	2370	?	F.	Unknown	—	" 2 "	2268	5 "	L	
2274	2400	35-3	M.	Unknown	—	Breast 3 "	2300	19 "	L	
2293	2485	37-3	M.	"	—	Breast 2 "	2510	15 "	L	
2303	1950	36	M.	"	—	Breast 4 "	2495	4 "	L	
2350	2490	38-1	M.	"	—	Breast 2 "	2460	33 "	L	
2355	2020	35-6	M.	Pulmonary T.B.	—	Breast 2 "	2466	7 "	L	
2452	2480	34-6	F.	P.E.	—	Artificial 3 "	—	2 "	N.D.	
2465	1770	34-4	F.	Twin Pregnancy	—	" 3 "	2410	12 "	L	1st Twin
2500	2450	34-1	F.	Unknown	—	Artificial 3 "	2490	49 "	L	
2505	1750	33-5	M.	"	—	" 2 "	2800	28 "	L	
1450	1450	33-5	F.	Twin Pregnancy	—	" 2 & 3 hrly.	2340	84 "	N.D.	1st Twin
2507	2400	33	F.	Unknown	—	" 3 hrly.	2450	24 "	L	2nd Twin

PREMATURE LIVE-BORN INFANTS (Continued)

Reg. No.	Birth Weight	Maturity of Fetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
2539	2050	32	F.	Unknown	—	Artificial 2 hdy.	7520	35 days	L	
2555	2430	38.4	F.	"	—	Breast 4 "	2495	8 "	L	
2560	2000	33	M.	"	—	Artificial 2 "	2470	43 "	L	
2588	2360	39.2	F.	"	—	" 2 "	2480	14 "	L	
2605	2035	32.5	M.	"	—	" 2 "	2580	48 "	L	
2639	2350	37.3	F.	"	—	" 3 "	2460	27 "	L	
2658	2250	34.4	F.	"	—	" 3 "	2540	20 "	L	
2660	2480	36	M.	"	—	Breast 2 "	2450	8 "	L	1st Twin
2694	2270	40.4	F.	Twin Pregnancy	—	Artificial 4 "	2683	10 "	L	2nd Twin
	2140		M.	"	—	" 2 "	2381	24 "	L	
2701	2350	36.2	M.	"	—	" 2 "	2450	14 "	L	
2706	1250	31.2	F.	"	—	" 3 "	2500	10 "	N.D.	
2725	2290	39.2	F.	"	—	Breast 4 "	2510	16 "	L	
2740	2450	36.2	F.	"	—	Artificial 2 "	2400	19 "	L	1st Twin
2784	2350	38.2	M.	Twin Pregnancy	—	" 2 "	2370	49 "	L	2nd Twin
	2150		F.	"	—	" 2 "	2400	45 "	L	2nd Child
2818	2070	42.1	M.	Triplets Pregnancy	—	—	—	9 hrs	N.D.	
2826	1900	23	F.	Unknown	—	Artificial 2 "	—	6 days	N.D.	
2883	1160	29.4	F.	"	—	" 3 "	2495	10 "	L	
2886	2470	40.2	F.	P.E.	—	Breast 3 "	2455	14 "	L	
2897	2400	39	M.	Unknown	—	" 2 "	2470	36 "	L	
2956	2040	31.5	M.	"	—	" 2 "	—	11 "	N.D.	
2991	2320	39	F.	Mitral Stenosis-Class IV	—	" 2 "	2500	17 "	L	
3003	2395	?	M.	Unknown	—	" 2 "	2380	9 "	L	
3020	3350	35.2	M.	"	—	Artificial 2 "	—	1 day	N.D.	
3025	1280	26.3	F.	"	—	" 2 "	2400	10 days	L	
3031	2430	43.5	F.	"	—	" 2 "	—	28 "	L	
3053	1900	35.3	F.	Essential Hypertension and Pre-eclampsia and Unknown	—	" 2 "	2350	—	L	
3057	1830	34.3	F.	"	—	" 2 "	2340	42 "	L	
3065	2250	34.5	F.	"	—	" 2 "	—	6 "	N.D.	
3068	2170	38.2	F.	"	—	" 2 "	2480	23 "	L	
3089	2470	35.6	F.	"	—	Breast 4 "	2211	7 "	L	
3112	2495	39.2	M.	"	—	" 4 "	2600	12 "	L	
3116	2340	38	F.	"	—	Artificial 2 "	2460	14 "	L	
3128	2370	35.2	M.	"	—	Artificial 2 "	2360	13 "	L	
3148	2000	36	F.	P.E.	—	Artificial 2 "	2400	14 "	L	1st Twin
	2180		F.	Twin Pregnancy	—	" 2 "	2370	25 "	L	2nd Twin
3155	2466	36.5	M.	P.E.	—	Breast 4 "	2495	14 "	L	
3176	1500	35.4	F.	Unknown	—	—	—	9 hrs.	N.D.	
3189	2320	38.6	F.	"	—	Artificial 2 "	—	8 days	N.D.	
3199	2100	?	F.	P.E. + T.B.	—	Breast 4 "	2523	18 "	L	
3222	2460	35.6	M.	Unknown	—	Artificial 2 "	2340	5 "	L	

PREMATURE LIVE-BORN INFANTS, (Continued)

Reg. No.	Birth Weight	Maturity of Fœtus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
3246	2450	39.4	F.	Unknown	—	Breast 3 haly.	2329	9 days	L	
3274	2430	36.3	M.	"	—	Artificial 2 "	2505	20 "	L	
3287	2410	41.6	F.	"	—	" 3 "	2650	16 "	L	
3317	1970	38.5	M.	"	Torsion of Tentorium Subdural	Artificial 2 "	—	4 "	N.D.	
3371	2450	35.4	M.	"	—	Breast 4 "	2515	8 "	L	1st Twin
3404	2120	38.5	F.	Twin Pregnancy Unknown	—	Artificial 2 "	2480	25 "	L	
3421	2470	36.3	M.	"	—	Breast 4 "	2580	24 "	L	
3431	2400	38.4	F.	"	—	Artificial 2 "	2490	17 "	L	
3450	2440	38.4	M.	"	—	Artificial 2 "	2305	5 "	L	
3458	2480	36.6	M.	"	—	Breast 3 "	2381	8 "	L	
3477	2170	34.4	M.	"	—	Artificial 2 "	2350	25 "	L	
3519	2480	41	F.	"	—	Breast & Artificial 2 "	2472	5 "	L	
3532	1420	38.5	F.	"	—	—	—	7 hrs.	N.D.	
3543	1300	28.3	M.	"	—	—	—	2 days	N.D.	
3618	1700	32.4	F.	"	—	Artificial 2 "	2290	86 "	L	
3667	960	29.1	M.	"	—	—	—	2 "	N.D.	
3675	2480	39.2	M.	"	—	Breast 4 "	2480	13 "	L	
3676	2260	38	F.	Hypertension > due to Aortic Aneurysm	—	Artificial 2 "	2381	7 "	L	
3726	2100	33.4	M.	Unknown	—	Artificial 2 "	2350	14 "	L	
3736	1830	31	M.	"	—	" 2 "	—	10 "	N.D.	
3768	2240	38.2	F.	"	—	Breast 4 "	2250	28 "	L	
3774	2495	35.4	M.	"	—	Artificial 2 "	2353	4 "	L	
3799	2300	36	M.	Twin Pregnancy	—	Artificial 2 "	2340	15 "	L	1st Twin
3839	2180	36	M.	"	—	" 2 "	2310	14 "	L	2nd Twin
3841	2460	41	M.	Unknown	—	Breast 4 "	2460	6 "	L	
3841	2150	?	F.	"	—	" 2 "	2300	47 "	L	
3880	2250	37.4	M.	"	—	" 2 "	2360	29 "	L	
3903	2030	37	F.	"	—	Breast 4 "	2000	26 "	L	
3905	2450	36.5	M.	Twin Pregnancy	—	Artificial 2 "	2244	9 "	L	1st Twin
3968	2200	39	F.	Unknown	—	Breast 4 "	—	—	N.D.	2nd Twin
3973	2350	39.5	F.	"	—	Artificial 2 "	2250	19 "	L	
4007	2450	38.1	M.	"	—	Artificial 2 "	2300	4 "	L	
4030	2160	44.3	F.	"	—	Breast 4 "	2450	3 "	L	
4034	1190	29.5	M.	Twin Pregnancy	Subdural	" 4 "	2325	3 "	L	
4041	1240	38.2	M.	Unknown	—	" 4 "	—	2 "	N.D.	1st Twin
4058	2470	38.6	M.	"	—	Breast 4 "	2335	8 "	L	2nd Twin
4064	2430	34.4	F.	"	—	" 4 "	2500	6 "	L	
4087	2480	35.5	F.	"	—	" 4 "	2466	7 "	L	
4089	2080	40	M.	"	—	Artificial 2 "	2320	5 "	L	
4106	2240	41.2	F.	"	—	" 2 "	2040	5 "	N.D.	
							2360	15 "	L	

PREMATURE LIVE-BORN INFANTS (Continued)

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
4158	1800	?	F.	Unknown	—	Breast 3 hrly.	—	3 days	N.D.	
4164	2480	36-3	F.	"	—	Artificial 2 "	2325	7 "	L	
4177	2350	39-3	M.	"	—	Artificial 2 "	2400	5 "	L	
4205	2080	43-3	M.	"	—	Breast 4 "	2495	17 "	N.D.	
4206	2380	41-4	F.	"	—	Breast 4 "	—	17 "	L	
4221	2150	30-4	M.	"	—	Artificial 2 "	—	3 "	N.D.	
4224	2260	34-4	F.	"	—	" 2 "	—	18 "	N.D.	
4256	2000	34	F.	"	—	" 2 "	—	19 "	N.D.	
4310	2300	37-5	F.	"	—	Breast 3 "	2350	6 "	L	
4311	2380	37-2	F.	"	—	Artificial 2 "	2428	5 "	L	
4326	1785	31	M.	"	—	Breast 3 "	—	9 "	N.D.	
4394	2470	42	F.	"	—	Artificial 2 "	2410	7 "	L	
4397	2470	39-1	M.	Twin Pregnancy	—	Breast 2 "	—	9 "	L	
4413	1760	32-6	M.	Unknown	—	Artificial 2 "	2523	7 "	L	
4434	2350	32-3	F.	"	—	Artificial 2 "	—	7 "	L	
4460	1300	35-5	F.	Twin Pregnancy	—	Artificial 2 "	2470	1 day	N.D.	1st Twin
4461	1660	39-4	F.	"	—	" 2 "	—	11 "	L	
4461	2460	39-4	F.	"	—	" 2 "	—	11 "	L	
4495	2150	39	F.	"	—	Breast 4 "	2495	13 "	N.D.	1st Twin
4501	2400	34-1	F.	Unknown	—	Artificial 2 "	1850	17 "	N.D.	2nd Twin
4516	2450	34-4	M.	"	—	Breast 4 "	2551	11 "	L	1st Twin
4522	2050	35-2	F.	"	—	Artificial 2 "	2430	5 "	L	2nd Twin
4570	1770	44-3	F.	Twin Pregnancy	—	Artificial 2 "	—	22 "	L	
4570	2110	44-3	F.	"	—	" 2 "	2480	34 "	L	
4585	1260	36	M.	Unknown	—	" 2 "	—	11 "	N.D.	1st Twin
4656	2080	39-5	F.	Twin Pregnancy	—	Artificial 2 "	—	6 "	N.D.	2nd Twin
4658	1200	26-2	F.	Unknown	—	Artificial 2 "	2140	13 hrs.	N.D.	
4670	1730	38-1	M.	"	—	" 2 "	—	40 days	L	
4698	1940	36-4	M.	"	—	Artificial 2 "	—	12 "	N.D.	
4705	2350	37	F.	"	—	Artificial 2 "	2350	5 hrs.	N.D.	
4710	2370	39-3	F.	"	—	Artificial 2 "	2120	11 "	L	Transferred to Q.M.H. because of signs of Respiratory Obstruction
4741	2410	36-1	M.	"	—	Breast 3 "	2325	4 "	L	
4758	2420	39-3	F.	"	—	Breast 4 "	2350	1 day	N.D.	
4760	2425	38-5	F.	"	—	" 4 "	2450	7 "	L	
4797	1900	39-5	F.	P.E. Tentorial Tear	—	" "	—	20 hrs.	N.D.	2nd Twin
4810	2400	37-2	M.	Twin Pregnancy	—	Breast 3 "	2430	5 days	L	
4829	1900	33-5	M.	Unknown	—	Artificial 2 "	—	7 "	N.D.	
4838	2370	36	F.	Pulmonary T.B.	—	Breast 2 "	2240	5 "	L	
4843	2300	40-4	F.	Unknown	—	Artificial 2 "	2260	27 "	L	D.A.A.
4861	2460	38-3	F.	"	—	Breast 4 "	2211	6 "	L	

PREMATURE LIVE-BORN INFANTS (Continued)

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
4891	1900	39.4	F.	P.F.	—	Artificial 2 hrly.	2300	36 days	L	
4916	2350	32.2	F.	Unknown	—	" 2 "	2220	40 "	L	
4926	2355	39.4	M.	"	—	" 2 "	2300	10 "	L	
5020	1820	34.4	M.	"	Subdural	Breast 4 "	2450	1 day	N.D.	
5034	2450	40.4	F.	"	Haemorrhage	Artificial 2 "	2240	8 days	L	D A A
5048	2155	38.5	F.	"	—	Breast 4 "	2295	4 "	L	
5080	2450	"	F.	"	—	" 2 "	—	"	N.D.	
5090	2430	36.2	M.	"	—	" 2 "	2120	11 "	L	
5091	2400	37.5	F.	"	—	Artificial 2 "	2495	6 "	L	
5143	2450	38.5	F.	"	—	Breast 4 "	2460	8 "	L	
5195	2466	39.3	M.	"	—	Artificial 4 "	2495	8 "	L	
5205	2100	34.5	F.	"	—	Breast 4 "	2495	8 "	L	
5207	2470	37.3	M.	"	—	" 4 "	2495	4 "	L	
5209	2130	36.1	M.	"	—	" 4 "	—	1 hour	N.D.	
5218	2300	38.5	M.	"	—	Artificial 2 "	2310	25 days	L	
5231	2280	37	F.	"	—	" 3 "	—	15 "	N.D.	
5240	1800	29	F.	"	—	" 2 "	2230	48 "	L	1st Twin
5246	1400	31.1	F.	Unknown	Subdural	Artificial 2 "	—	2 hrs.	N.D.	2nd Twin
5251	2495	39.3	F.	Twin Pregnancy	Haemorrhage	" 2 "	—	6 days	N.D.	
5261	2450	43.5	F.	"	Subdural	Artificial 2 "	2409	14 "	L	1st Twin
5288	1530	35.4	F.	Unknown	Haemorrhage	Breast 4 "	2495	7 "	L	2nd Twin
5326	2060	38.2	F.	"	—	" 4 "	2120	56 "	L	
5330	2280	39.4	F.	"	—	Artificial 2 "	2340	26 "	L	
5338	930	29	F.	Mitral Stenosis	—	" 2 "	—	11 "	N.D.	
5346	2450	30	F.	P.E.	—	" 3 "	2520	130 "	L	
5356	2420	38.3	F.	Unknown	—	Breast 4 "	2325	5 "	L	
5370	2350	44.3	M.	"	—	" 4 "	2353	7 "	L	
5402	2400	38.6	F.	Twin Pregnancy	—	Artificial 4 "	2650	14 "	L	
5424	2300	37.1	M.	Unknown	—	Breast 4 "	2340	9 "	L	2nd Twin
5445	2050	38.2	F.	"	—	Breast & Artificial 2 "	2310	12 "	L	
5462	2400	"	F.	"	—	Breast 4 "	1960	8 "	L	D A A
5471	2300	39.3	M.	"	—	" 4 "	2325	5 "	L	
5480	2400	37.4	M.	"	—	" 4 "	2300	7 "	L	
5498	2190	38.6	F.	"	—	Artificial 2 "	2280	30 "	L	
5514	2350	37	F.	"	—	Breast 4 "	2260	10 "	L	D A A.
5516	2090	41.1	M.	Pulmonary T.B.	—	Artificial 2 "	2290	5 "	L	
5520	2260	36.6	M.	Unknown	—	" 2 "	1700	11 "	N.D.	D A A.
5523	2320	39.2	M.	"	—	" 2 "	—	11 "	N.D.	
5526	1600	32.4	M.	"	—	Breast 4 "	2220	6 "	N.D.	D A A
5527	2440	39.5	F.	"	—	Breast 4 "	3380	9 days	L	
5558	1600	33	F.	Mitral Stenosis	—	Artificial 2 "	2100	58 "	L	

PREMATURE LIVE-BORN INFANTS (Continued)

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
5565	2450	39.2	F.	Unknown	—	Breast 4 hrly.	2300	5 days	L	
5587	2300	39.1	F.	"	—	" 4 "	2421	13 "	L	
5599	2300	32.5	M.	"	—	Artificial	—	16 "	N.D.	
5600	2320	34.6	F.	"	—	"	—	11 "	N.D.	
5621	2495	36	M.	"	—	Artificial	2827	19 "	L	
5623	2050	35	F.	"	—	Artificial 3 & Breast	2260	28 "	L	1st Twin
5650	2400	34.3	M.	Twin Pregnancy	—	Artificial 2 "	—	4 "	N.D.	2nd Twin
5659	2100	34.20	M.	Unknown	—	Artificial 5 "	2250	23 "	L	
5659	2420	39.5	M.	"	—	Artificial 2 "	2260	4 "	L	
5669	2450	38.3	M.	"	—	Breast 3 "	2500	4 "	L	
5670	2150	33.5	F.	"	—	" 4 "	—	9 "	N.D.	
5686	2005	31.3	M.	"	—	Artificial 2 "	—	11 "	N.D.	
5694	2430	34	F.	Pulmonary T.B.	—	" 2 "	2325	14 "	L	
5716	2450	37.1	F.	Unknown	—	Breast 4 "	2368	7 "	L	
5718	2350	37.5	M.	"	—	" 4 "	2250	4 "	L	
5728	2450	38.1	M.	"	—	" 4 "	2478	7 "	L	
5777	1330	38.1	M.	Twin Pregnancy	—	Artificial 2 "	—	9 "	N.D.	1st Twin
5793	1390	35.2	M.	"	—	" 2 "	—	14 "	N.D.	2nd Twin
5825	2140	36.2	M.	Unknown	—	" 2 "	—	6 "	N.D.	Transferred to Q.M.H. because of Allergic Reaction
5825	2110	36.2	F.	"	—	" 2 "	1800	13 "	L	
5854	2300	36.6	M.	"	—	" 3 "	—	13 "	N.D.	
5884	2352	41	M.	"	—	Breast 4 "	2381	7 "	L	
5886	2450	38.4	F.	"	—	Breast 4 "	2453	5 "	L	
5887	2450	40.5	F.	"	—	" 4 "	2296	7 "	L	
5913	1830	36.4	F.	Twin Pregnancy	—	Artificial 2 "	2300	25 "	L	1st Twin
5917	1800	36.4	F.	"	—	" 2 "	2220	47 "	L	2nd Twin
5932	2410	38.3	F.	Unknown	—	Breast 2 "	2353	7 "	L	1st Twin
5932	2300	38.2	F.	Twin Pregnancy	—	Artificial 2 "	2280	16 "	L	2nd Twin
5957	2090	32.6	F.	Unknown	—	" 2 "	2250	36 "	L	1st Twin
5959	1480	37	F.	"	—	" 2 "	2280	50 "	L	2nd Twin
5966	2360	35.6	M.	"	—	" 2 "	2280	8 "	L	
5967	2720	38.1	F.	Twin Pregnancy	Subdural Haemorrhage	" 3 "	2200	6 "	N.D.	2nd Twin
5989	2215	36.4	F.	"	—	Breast 4 "	2948	16 "	L	1st Twin
6028	2250	35.6	M.	Unknown	—	Artificial 2 "	2330	27 "	L	2nd Twin
6038	2320	35.6	M.	"	—	" 2 "	1840	12 "	N.D.	
6044	2370	39.5	M.	"	—	" 2 "	2100	11 "	L	
6057	2350	36.2	F.	"	—	" 2 "	2310	5 "	L	
6055	2440	35.4	M.	Pulmonary T.B.	—	Breast 3 "	2211	5 "	L	
6064	2180	42.6	F.	Unknown	—	Artificial 4 "	2268	23 "	L	
6070	2150	38.6	F.	"	—	" 2 "	1984	30 "	L	
6070	2000	38.6	M.	"	—	" 2 "	2300	25 "	L	
6070	2000	38.6	M.	"	—	" 2 "	2310	21 "	L	

PREMATURE LIVE-BORN INFANTS, (Continued)

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
6081	2440	35-2	F.	Unknown	—	Breast 4 hrly.	2455	13 days	L.	
6117	1700	34 6	M.	Twin Pregnancy	—	Artificial 2 "	—	5 "	N.D.	1st Twin
6133	2100	35	M.	"	—	Artificial 2 "	—	6 "	N.D.	2nd Twin
6145	1600	31	F.	"	—	Artificial 2 "	2260	32 "	L.	1st Twin
6161	2200	38-1	M.	Unknown	—	Breast 4 "	2100	35 "	L.	2nd Twin
6168	1850	37-4	F.	"	—	Artificial 2 "	2268	8 "	L.	
6187	2420	40-5	F.	"	—	Artificial 2 "	2608	15 "	L.	
6204	2320	42	F.	"	—	Breast 4 "	2310	30 "	L.	
6205	2260	35	F.	"	—	" 4 "	2466	4 "	L.	
6223	2400	33	M.	"	—	" 4 "	2155	23 "	L.	
6232	1575	34-5	M.	Twin Pregnancy	—	Breast 4 "	—	2 "	N.D.	
2200	2200	34-5	F.	"	—	Artificial 2 "	2353	5 "	L.	1st Twin
6316	1950	35-5	M.	Unknown	—	Artificial 2 "	2320	32 "	L.	2nd Twin
6340	2330	35-4	M.	"	—	" 2 "	2220	48 "	L.	
6378	2450	42-5	M.	Twin Pregnancy	—	" 2 "	2300	38 "	L.	
6389	2050	36	F.	Unknown	—	Breast 2 "	2290	6 "	L.	
6392	2120	38-6	M.	"	—	" 4 "	2350	7 "	L.	
6397	2160	37-4	F.	Twin Pregnancy	—	" 4 "	—	2 "	N.D.	
6399	2370	42-3	F.	"	—	Artificial 2 "	2290	21 "	L.	
6402	2130	33-3	F.	Unknown	—	" 2 "	2150	20 "	L.	1st Twin
6425	1620	33	F.	"	—	Breast 2 "	2268	10 "	L.	2nd Twin
6430	2450	38	M.	"	—	Artificial 2 "	2280	20 "	L.	
6435	2350	35-5	F.	"	—	" 2 "	—	47 "	L.	
6435	1160	35-6	F.	"	—	Breast 4 "	2381	6 "	L.	
6472	2050	37-4	F.	"	—	" 4 "	2168	7 "	L.	
6492	2150	39-6	M.	Twin Pregnancy	—	" 4 "	—	3 1/2 hrs.	N.D.	
6528	2340	36-2	M.	"	—	Artificial 2 "	2270	20 days	N.D.	
6560	2350	40-6	F.	Unknown	—	Breast 2 "	2268	7 "	L.	1st Twin
6583	2380	37-3	F.	"	—	Artificial 2 "	2450	50 "	L.	2nd Twin
6586	2450	39-6	M.	"	—	Breast 3 "	2380	6 "	L.	
6622	2440	43	M.	"	—	" 3 "	2423	24 "	L.	
6678	1760	39-6	M.	"	—	Artificial 3 "	2350	11 days	L.	
6681	2480	39-6	F.	"	—	Breast 4 "	2750	1 day	N.D.	
6686	1840	?	F.	"	—	Artificial 2 "	1780	12 days	L.	D.A.A.
6691	2200	34-2	F.	"	—	Artificial 2 "	2230	50 "	L.	
6710	2450	31-1	F.	"	—	Breast 2 "	2250	27 "	L.	
6755	1990	36-3	F.	"	—	Artificial 4 "	—	4 "	L.	
6765	1560	35-6	M.	"	—	" 2 "	—	3 "	N.D.	

Baby died age 47 days

PREMATURE LIVE-BORN INFANTS, (Continued)

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
6081	2440	35-2	F.	Unknown	—	Breast 4 hly.	2455	13 days	L.	1st Twin
6117	1700	34-6	M.	Twin Pregnancy	—	Artificial 2 "	—	5 "	N.D.	2nd Twin
6133	1900	35	M.	"	—	Artificial 2 "	2300	6 "	N.D.	2nd Twin
6145	1600	35	F.	"	—	Artificial 2 "	2300	35 "	L.	1st Twin
6161	2400	31	F.	Unknown	—	Breast 4 "	2608	8 "	L.	2nd Twin
6168	1850	38-1	M.	"	—	Artificial 2 "	2310	15 "	L.	
6187	2420	37-4	F.	"	—	Breast 4 "	2466	30 "	L.	
6204	2420	40-5	F.	"	—	Breast 4 "	2155	4 "	L.	
6205	2320	42	F.	"	—	" 4 "	—	3 "	L.	
6223	2260	35	F.	"	—	" 4 "	—	2 "	N.D.	
6232	2400	33	M.	"	—	Breast 4 "	2353	5 "	L.	1st Twin
6316	1575	34-5	F.	Twin Pregnancy	—	Artificial 2 "	2320	32 "	L.	2nd Twin
6316	2200	35-5	F.	"	—	Artificial 2 "	2220	48 "	L.	
6340	1950	35-4	M.	Unknown	—	" 2 "	2400	38 "	L.	
6378	2330	35-4	M.	"	—	" 2 "	2290	6 "	L.	
6389	2450	42-5	M.	Twin Pregnancy	—	Breast 2 "	2350	7 "	L.	2nd Twin
6389	2050	36	F.	Unknown	—	" 4 "	—	2 "	N.D.	
6392	2120	38-6	M.	"	—	Artificial 2 "	2290	21 "	L.	
6397	2160	37-4	F.	Twin Pregnancy	—	" 2 "	2150	20 "	L.	1st Twin
6397	2370	37-4	F.	"	—	Artificial 2 "	2268	10 "	L.	2nd Twin
6399	2130	42-3	F.	Unknown	—	Breast 2 "	2280	20 "	L.	
6402	1620	33-3	F.	"	—	Artificial 2 "	—	47 "	L.	Baby died age 47 days
6425	2450	38	M.	"	—	" 2 "	2181	6 "	L.	
6430	2350	35-5	F.	"	—	" 4 "	—	7 "	L.	
6435	1160	35-6	F.	"	—	" 4 "	2168	7 "	L.	
6472	2090	37-4	M.	"	—	" 4 "	—	3½ hrs.	N.D.	
6492	2150	30-6	F.	Twin Pregnancy	—	Artificial 2 "	2270	1 day	N.D.	
6528	2340	36-2	M.	"	—	Breast 2 "	2368	20 days	L.	1st Twin
6560	2150	36-2	M.	"	—	Artificial 2 "	2350	50 "	L.	2nd Twin
6560	2350	40-6	F.	Unknown	—	Artificial 2 "	2110	26 "	L.	
6563	2380	37-3	F.	"	—	Breast 3 "	2380	6 "	L.	
6566	2450	39-6	M.	"	—	" 3 "	2423	24 "	L.	
6622	2440	43	M.	"	—	Artificial 3 "	2550	11 days	L.	
6678	1760	32-3	M.	"	—	Breast 4 "	2750	12 days	L.	D.A.A.
6681	2480	39-6	F.	"	—	Artificial 2 "	1780	50 "	L.	
6686	1840	?	F.	"	—	Breast 2 "	2230	27 "	L.	
6691	2200	34-2	M.	"	—	" 2 "	2250	4 "	L.	
6710	2450	31-1	F.	"	—	Breast 4 "	—	6 "	N.D.	
6755	1990	36-3	F.	"	—	Artificial 2 "	—	5 "	N.D.	
6765	1560	35-6	M.	"	—	" 2 "	—	3 "	N.D.	

PREMATURE LIVE-BORN INFANTS (Continued)

Reg. No.	Birth Weight	Maturity of Fetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
6771	2350	36	F.	Unknown	—	Breast 2 hrly. Artificial 2 "	3296	6 days	L	
6795	2220	38.4	M.	"	—	"	2220	29 "	L	
6807	2150	37	M.	Twin Pregnancy	—	"	2018	18 "	L	2nd Twin
6819	2400	40.1	F.	Unknown	—	Breast 3 "	2280	3 "	L	
6845	2150	31.4	M.	"	—	Artificial 2 "	2300	30 "	L	
6839	2320	34.4	M.	"	—	Breast 3 "	2240	3 "	L	D.A.A.
6851	2350	38.2	F.	"	—	Artificial 2 "	2100	3 "	L	D.A.A.
6861	2070	7	F.	"	—	"	2220	23 "	L	
6869	2020	40	M.	"	—	"	—	—	L	
6888	2350	38.5	M.	"	—	Breast 4 "	2255	6 "	L	
6902	2400	40.2	F.	Pulmonary T.B.	—	"	2211	5 "	L	
6946	2302	39.5	F.	Unknown	—	"	2495	12 "	L	
6952	2350	37.4	F.	"	—	"	2325	4 "	L	
6957	2200	37	M.	"	—	"	2240	12 "	L	
6962	2300	41	F.	"	—	Artificial & Breast 2 "	2320	9 "	L	Imperforate Anus, Transferred to Q.M.H.
6976	2380	41.1	M.	"	—	Breast 2 "	2248	9 "	L	
7001	2350	39.6	M.	"	—	Breast 3 "	2495	6 "	L	
7022	2440	39	F.	"	—	"	2466	8 "	L	
7050	1850	43.4	F.	"	—	Artificial 2 "	2280	54 "	L	
7084	1645	38.4	F.	"	—	"	2290	54 "	L	
7096	2390	41.5	M.	"	—	Breast 3 "	2240	9 "	L	
7100	2100	39.6	F.	"	—	Artificial 2 "	2796	14 "	L	
7148	2400	34	M.	"	—	Breast 4 "	2410	4 "	L	
7160	2350	32.6	F.	"	—	"	2353	8 "	L	
7171	1850	36.2	F.	"	—	Artificial 2 "	1850	16 "	L	
7175	2230	39.2	F.	Twin Pregnancy	—	"	2220	24 "	L	D.A.A. 1st Twin
7183	2150	38	M.	Twin Pregnancy	—	"	2220	23 "	L	2nd Twin
7184	2230	37.2	F.	"	—	Artificial 2 "	2220	28 "	L	1st Twin
7193	1660	40	F.	Unknown	—	"	2240	32 "	L	2nd Twin
7194	1750	40	M.	"	—	"	2240	7 "	L	2nd Twin
7213	2420	40.3	F.	"	—	"	2210	65 "	L	1st Twin
7232	2320	39	M.	"	—	"	2210	58 "	L	2nd Twin
7250	2150	38.5	F.	"	—	"	2200	6 "	L	
7286	2450	38.1	F.	"	—	"	2220	30 "	L	
7297	2410	37.2	F.	"	—	Breast 4 "	2466	11 "	L	
7308	2240	36.1	F.	"	—	"	—	2 "	N.D.	
7322	1446	31.2	F.	"	—	Artificial 2 "	2400	5 "	L	N.D.
7341	2000	34	M.	"	—	"	2230	16 "	L	
7341	2300	35.2	F.	"	—	"	2220	43 "	L	
7344	2400	39.3	F.	"	—	Breast 4 "	2296	12 "	L	

PREMATURE LIVE-BORN INFANTS (Continued)

Reg. No.	Birth Weight	Maturity of Foetus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge	Day of discharge or death	Requit	Remarks
8077	2000	38 4	F.	Twin Pregnancy	—	Artificial 2 hrly.	2230	37 days	L	1st Twin
8089	1650	37	M.	Unknown	—	Breast 4 "	2401	16 "	N.D.	2nd Twin
8094	2080	37-2	F.	"	—	Artificial 3 "	2650	51 "	L	"
8127	2170	34-4	M.	"	—	" 2 "	2500	34 "	L	"
8162	2280	35	F.	"	—	Breast 4 "	2150	5 "	L	"
8225	2268	36 6	F.	"	—	" 3 "	3118	37 "	L	"
8231	2090	37-1	F.	"	—	Artificial 2 "	2070	11 "	L	D.A.A.
8272	2450	39	F.	"	—	Breast 4 "	2380	7 "	L	"
8287	2430	39	F.	"	—	" 3 "	2430	9 "	L	"
8301	2400	37-4	M.	"	—	Artificial 2 "	—	8 "	N.D.	"
8312	2200	30-1	F.	Twin Pregnancy	—	Breast 3 "	2135	3 "	N.D.	2nd Twin
8321	2320	38-5	M.	Unknown	—	Artificial 2 "	2265	5 "	L	"
NON-BOOKED										
102	1150	27-3	M.	Unknown	—	Artificial 2 hrly.	1140	3 days	N.D.	"
523	2470	33	F.	Left Hydrocephrosis and Left Ureteric Calculus	—	Artificial 2 "	2466	13 "	L	"
792	1900	30 4	F.	Twin Pregnancy	—	Breast 3 "	2750	68 "	L	1st Twin
909	1630	35-2	F.	"	—	Artificial 2 "	—	4 "	N.D.	2nd Twin
1019	1780	38 4	F.	"	—	" 3 "	2320	51 "	L	1st Twin
1166	2430	33-3	F.	Twin Pregnancy	—	" 3 "	2590	20 "	L	2nd Twin
1513	2220	36 3	F.	Unknown	—	" 2 "	2630	53 "	L	1st Twin
1707	2420	37 1	M.	"	—	Breast 3 "	2480	33 "	L	2nd Twin
1829	1650	37 5	M.	"	—	Artificial 3 "	3140	48 "	N.D.	"
1837	2050	34 3	M.	"	—	" 2 "	—	8 hrs.	L	"
1925	1340	32	M.	"	—	" 3 "	2690	45 days	L	"
2052	2000	36-5	M.	"	—	" 3 "	2500	28 "	L	"
2291	1850	31-2	F.	"	—	" 2 "	2520	1 day	N.D.	"
2484	1950	37-6	F.	"	—	" 2 "	2200	33 days	L	"
2489	2275	37 4	M.	Pulmonary T.B. Placenta Praevia	—	" 2 "	2323	29 "	L	"
2566	1720	32 4	M.	Type IV	—	" 2 "	2450	15 "	L	"
2610	1840	32 4	F.	Unknown	—	Artificial 2 "	—	4 days	N.D.	"
2743	1670	35-2	F.	Placenta Praevia Type IV	—	" 2 "	2440	75 "	L	"
2766	1080	28 4	M.	Unknown	Laceration of Tentorium	" "	—	11½ hrs.	N.D.	"

PREMATURE LIVE-BORN INFANTS (Continued)

Reg. No.	Birth Weight	Maturity of Fœtus	Sex	Cause of Premature Labour	Birth injury, if any	Method of feeding	Weight on discharge gm.	Day of discharge or death	Result	Remarks
2837	2470 1950	40-1	M. F.	Twin Pregnancy	Laceration of Tentorium	Breast and Artificial 2 hrly. Artificial 2 "	2665	14 days 2 "	L. N.D.	1st Twin 2nd Twin
2882	2350	30	M.	Unknown	—	" "	2360	15 "	L.	
3085	2300	?	F.	"	—	" "	—	2 "	N.D.	
3575	2480	35-6	M.	Twin Pregnancy	—	" "	2410	15 "	L.	1st Twin
3704	2400	35-3	M.	Unknown	Subdural Haemorrhage	Breast and Artificial 4 "	—	3 "	N.D.	
4245	1380	35-1	F.	"	—	Artificial 2 "	—	19 "	N.D.	
4440	2438	?	F.	"	—	Breast 4 "	2520	11 "	L.	
4566	2280	36-3	F.	"	—	Artificial 2 "	2300	22 "	L.	
4889	2400	?	M.	"	—	Artificial 3 "	—	3 "	N.D.	
4997	2400	36	M.	P.E.	—	" 4 "	2381	5 "	L.	B B A.
5476	2270	33	M.	Unknown	—	Breast 2 "	—	4 "	N.D.	
6094	2700	30-6	F.	"	—	Artificial 2 "	2240	7 "	L.	
6252	2130	35	M.	"	—	Breast 4 "	—	19 "	N.D.	
6380	2000	37-4	F.	"	—	Artificial 2 "	—	12 "	L.	
6872	2250	32-5	F.	"	—	" 2 "	2296	28 "	L.	
7081	1620	31	F.	"	—	" 2 "	2250	5 hrs.	N.D.	
7999	1800	34-5	F.	"	—	" 2 "	—	14 days	N.D.	
8112	1850	36-5	F.	"	—	" 2 "	2140	26 "	L.	
7483	2490	35	F.	"	—	" 2 "	2220	27 "	L.	

TABLE 38

CONGENITAL MALFORMATIONS

69 babies showed evidence of congenital malformation. 67 were Booked cases and 2 Non-booked.

Reg. No.	Nature of Malformation	Birth Weight gm.	Sex	Result	Intercurrent maternal disease	Remarks
BOOKED						
3						
10	Syndactylism, left 4th, 5th toes	4320	F.	L	P.E.	
273	Erupted tooth	3230	F.	L	---	
428	Pedunculated rudimentary digit (right thumb)	2420	F.	L	---	
478	Hydromyelia	5250	M.	S.B.	---	
591	Hydromyelia	2650	F.	S.B.	---	
608	Achondroplasia	2850	F.	L	---	
649	Cleft palate	2100	F.	L	---	
662	Talipes equinus varus (left)	1850	M.	S.B.	---	
190	Wilim's tumour	2950	M.	N.D.	---	
1089	Anencephaly	2800	M.	L	Severe P.E.	
1090	Harelip, right side	2730	M.	L	---	
1336	Harelip, bilateral; cleft palate, complete	1040	M.	S.B.	---	
1552	Erupted teeth	2600	F.	L	Pulmonary	
1571	Phymosis	2948	F.	L	---	
1590	Hare-lip, left	3325	F.	L	---	
1677	Gynacomastia, bi'ateral	3800	M.	L	---	
1851	Hydromyelia	1620	M.	S.B.	---	
2180	Hydrocephalus, harelip and cleft palate, malformation of nose, ears and eyes undescended testis, bilateral	3000	M.	N.D.	---	
2323	Anencephaly	3570	M.	S.B.	---	
2335	Hypostadus and split prepuce	2845	M.	L	---	
2388	Anencephaly	3340	M.	S.B.	---	
2402	Unilateral harelip and complete cleft palate	3200	M.	L	---	
2521	Extra thumb in left hand	3150	F.	L	---	
2534	Anencephaly	1750	M.	S.B.	---	
2657	Hydrocephalus	2500	M.	N.D.	---	
2859	Eruption of teeth	2700	M.	L	---	
2889		2850	M.	L	---	
2981	Mild degree of bilateral club foot	3150	F.	L	---	
3069	Hemiphordite	3190	F.	L	---	
3075	Cleft palate	3580	F.	L	---	
3117	Extra left thumb	2680	M.	L	---	
3569	Facialorocle	3150	F.	L	---	
3578	Hydromyelia	3030	M.	S.B.	---	
3643	Anencephaly	2180	F.	S.B.	---	
3826	Extra digit, rt. hand.	2820	M.	L	---	

2nd Twin

CONGENITAL MALFORMATIONS (Continued)

Reg. No.	Nature of Malformation	Birth Weight gm.	Sex	Result	Intercurrent maternal disease	Remarks
3835	Webbed toes; atresia of oesophagus; Oesophago-tracheal fistula	2650	M.	N.D.	---	---
4007	Imperforate anus	2450	M.	L.	---	---
4391	Hydrocephalus	1800	M.	S.B.	---	---
4343	Extra digit, left hand	2580	M.	L.	---	---
4318	Erupted tooth	2800	F.	L.	---	---
4475	Harelip, left	3000	M.	L.	---	---
4486	Extra toe, bilateral; lateral 3 toes webbed	3240	M.	L.	---	---
4663	Extra thumb, rt.	2910	M.	L.	---	---
4817	Extra thumb, left	2448	F.	L.	---	---
4932	Syringo myelocoele	2900	F.	L.	---	---
5209	Encephalocoele; extra toe, bilateral	2130	M.	N.D.	---	---
5520	Harelip, left; cleft palate, incomplete; hypospadias	2268	M.	N.D.	---	---
5582	Hypospadias	3440	M.	L.	---	---
5602	Extra thumb, left hand	3580	M.	L.	---	---
5763	Talipes equinus varus	3000	M.	L.	---	---
5929	Paraurethral cyst	2950	F.	L.	---	---
5971	Spina bifida; meningocoele	2510	F.	L.	---	---
6709	Talipes equinus valgus, left	2750	M.	L.	---	---
6869	Imperforate anus	2020	M.	L.	---	---
6922	Talipes equinus varus, bilateral	3100	M.	L.	---	---
7079	Polyp, (in front of rt. ear).	3250	M.	L.	---	---
7155	Harelip, bilateral, cleft palate	3400	M.	L.	---	---
7187	Talipes equinus valgus, bilateral	2495	M.	L.	---	---
7358	Anencephaly	1435	F.	S.B.	---	---
7397	Talips equinus valgus	3000	M.	L.	---	---
7407	Harelip, left; cleft palate	3270	M.	L.	---	---
7457	Foetal ascites	3350	F.	S.B.	---	---
7774	Monster (no eyes)	1780	M.	S.B.	---	---
7845	Conjoined twins	4500	F.	N.D.	---	---
8126	Bilateral talipes equinus varus	2840	M.	L.	---	---
8139	Cleft, Palate	2750	M.	L.	---	---
8280	Talipes equinus valgus, rt.	3330	F.	L.	---	---
NON-BOOKED						
1548	Anencephaly	1480	M.	S.B.	Pulmonary	T.B.
3599	Conjoined twins (one head, 2 bodies)	2700	F.	N.D.	---	---

TABLE 39

BIRTH TRAUMA

There were 45 cases of birth trauma, 34 Booked and 11 Non-booked.

Reg. No.	Maturity	Birth Weight gm.	Sex	Type of Injury	Method of Delivery	Cause of Injury	Result	Remarks
BOOKED								
6787/55	44-1	4000	F.	Fracture of upper 1/3 of femur	M.R. + M.F.	Unknown	L.	
291/56	39-5	2200	F.	Laceration of tentorium	Spont.	Prematurity	N.D.	
340	40	2800	M.	" "	"	Shoulder presentation and spont. delivery	N.D.	
350	37-1	1820	F.	Mild subdural haemorrhage	"	Breech	N.D.	
543	37-1	2770	F.	Cephalo-haematoma	"	Prematurity	L.	
660	33-3	1800	F.	Laceration of tentorium	"	Version	S.B.	
836	39-6	2400	F.	Haemorrhage in lungs	Bipolar podalic Version	"	S.B.	
1049	39-1	3000	M.	Laceration of tentorium	Spont.	Prematurity	L.	
1082	40-2	3864	M.	Sterno-mastoid haematoma (L.)	Assisted Breech	Assisted Breech	L.	
1260	43	2700	F.	Facial palsy, right	M.R. + M.F.	M.R. + M.F.	S.B.	
1735	40-4	3750	M.	Laceration of tentorium	Spont.	"	N.D.	
1769	42-1	3175	F.	Forepicks mark on scalp	L.F.	L.F.	S.B.	
2040	39-1	3350	M.	R. Sterno-mastoid haematoma	Breech Assisted	Breech assisted	L.	
2268	40-1	3780	M.	Laceration of tentorium left eye-lid, forehead, and right side of head	M.F.	M.F.	S.B.	
2374	34-1	1790	F.	Laceration of tentorium	Spont.	Prematurity	S.B.	
2499	37-6	3175	M.	Cephalo-haematoma	Willitt's Forceps	Willitt's forceps	S.B.	Plaecenta Praevia Type III
2542	37-2	3050	M.	Laceration of tentorium	M.R. + M.F.	M.R. + M.F.	N.D.	2nd Twin
3317	38-5	1970	M.	Torsion of tentorium; subdural haemorrhage	Spont.	Breech	N.D.	
3341	40-6	2700	F.	Cephalo-haematoma	Spont. Breech	"	L.	
3578	39	2030	M.	Torsion of tentorium	L.F.	L.F.	S.B.	Hydrops Fetalis
3635	41-5	2650	M.	Extensive subdural haemorrhage	Spont.	"	N.D.	Oesophageal tracheal Fistula 1st Twin
4034	29-5	1190	M.	Subdural haemorrhage	"	Prematurity	N.D.	
4099	40	2080	M.	" "	"	"	N.D.	
4797	39-5	1900	M.	Tentorial tear	"	Version	N.D.	2nd Twin
4958	41	3300	F.	Subdural haemorrhage	M.F.	Breech	S.B.	
5020	34-4	1820	M.	" "	Spont.	"	N.D.	
5185	38-1	2970	F.	Paralysis of right arm, spasm of right sterno-mastoid	"	"	L.	
5240	29	1180	F.	Subdural haemorrhage	"	"	N.D.	

BIRTH TRAUMA (Continued)

Reg. No.	Maturity	Birth Weight gm.	Sex	Type of Injury	Method of Delivery	Cause of Injury	Result	Remarks
5246	31-1	1440	F.	Subdural haemorrhage	Spont.	Breech	N.D.	2nd Twin
5966	35-6	2360	M.	Subdural haemorrhage (right)	M.R. + M.F.	M.R. + M.F.	N.D.	2nd Twin
6957	?	2200	M.	Left cephalo-haematoma	L.S.C.S.	M.F., before C.S.	L	
6958	?	3200	F.	Tentorial laceration	M.F.	M.F.	N.D.	
6963	37	3370	M.	" "	Spont.	?	N.D.	
7094	41-4	3400	M.	Cephalo-haematoma	Spont.		L	
NON-BOOKED								
913	42-5	2550	M.	Cephalo-haematoma	M.R. + M.F.	M.R. + M.F.	N.D.	
1140	39	3000	M.	Laceration of tentorium	L.F.	Breech, assisted	N.D.	
1707	37-1	1650	M.	Cephalo-haematoma	Spont.	—	N.D.	
4932	37	2620	M.	Laceration of tentorium	M.R. + M.F.	Breech, assisted	N.D.	
2316	36-3	3062	M.	Subdural haemorrhage, moderate	M.R. + M.F.	M.R. + M.F.	N.D.	
2766	28-4	1080	M.	Laceration of tentorium	Spont.	?	N.D.	
2837	40-1	1950	F.	" "	Internal Version and Breech Extraction	Internal version	N.D.	2nd Twin
3323	40-6	3400	M.	Subdural haemorrhage	Spont.	?	S.B.	
3605	40-4	2588	M.	" "	Internal Version	Internal version	S.B.	
3704	35-3	2400	M.	" " (right)	M.F.	M.F.	N.D.	
4189	38-4	2800	F.	Cephalo-haematoma	Spont.	Breech	L	

NEO-NATAL INFECTIONS, (Continued)

Reg No	Maturity	Birth Weight gm	Sex	Type of Infection	Source of Infection	Dominant Causal Organism	Treatment	Result	Remarks
5330	39.4	2280	F	Broncho pneumonia	Respiratory	—	Penicillin Terramycin Achromycin	ND	10 d 11 Days MR + M1
5600	34.6	2320	F	"	"	—	"	ND	11
5663	2570	2570	M	"	"	—	Achromycin	ND	12
5670	33.5	2150	F	"	"	—	Penicillin & Achromycin	ND	9 Present Present C.C.S Present Present L.S.C.S
5686	31.3	2005	M	"	"	—	Penicillin Achromycin	ND	11
5777	"	1330	M	"	"	—	Sulphatirvid Penicillin Terramycin & Eucortone	ND	9 1st Twin
5777	"	1390	M	"	"	—	Penicillin Terramycin	ND	14 Shoulder Present tion 2nd Twin
5793	35.2	2140	M	"	"	—	Penicillin Chloromycetin	ND	6
5832	39	3250	F	"	"	—	Penicillin Chloromycetin Achromycin	ND	20 L.S.C.S Breach Presentation
5966	35.6	2360	F	"	"	—	"	ND	7
5969	36.4	2230	M	"	"	—	Terramycin & Achromycin	ND	12
6003	36	2500	M	"	"	—	"	ND	8
6117	34.6	1700	M	"	"	—	Penicillin	ND	5 1st Twin
7144	39.5	2650	F	Conjunctivitis	"	—	Penicillin eye drops	L	L.S.C.S P1
7592	30.2	1200	F	Broncho pneumonia	Respiratory	—	Penicillin & Terramycin	ND	2 Hours Accident Hemorrhagic L11 clump
8072	43.6	3000	M	"	"	—	Penicillin Streptomycin	ND	13 Days
8287	"	2200	M	"	"	—	Penicillin & Terramycin	ND	9 P1
NON BOOKED									
4889	?	2400	M	Broncho pneumonia	Respiratory	—	Penicillin Achromycin	ND	13 Days BBA
5476	33	2270	M	"	"	—	Penicillin Achromycin Penicillin	ND	3 Days
6252	21	2130	M	"	"	—	Chloromycetin Garamycin	ND	19 Breach Present tion
7988	37	2750	M	"	"	—	Chloromycetin Achromycin	ND	14 1st Twin
7999	34.5	1800	F	"	"	—	Penicillin Garamycin	ND	18 C.C.S Should be Present tion Problems of Cord Accident H11 morrhagic

TABLE 41

OTHER NEO-NATAL DISEASES

There were 53 babies which suffered from other neo-natal diseases.
49 were in Booked and 4 in Non-booked cases.

Reg. No.	Maturity	Birth Weight gm.	Sex	Type of Disease	Maternal Complication	Method of Delivery	Treatment	Result	Remarks
BOOKED									
101	38.5	2890	F.	Severe Jaundice	—	L.S.C.S	Vitamin K.	N.D.	
198	30.3	1500	F.	Mild Jaundice	—	Spont.	" "	N.D.	
340	40	2800	M.	Jaundice	—	" "	Vitamin K	N.D.	
401	?	3550	M.	Mild Jaundice	—	" "	" "	L	
497	40.1	2960	M.	Jaundice	—	" "	Vitamin K. Penicillin	L	
623	35	1870	M.	" "	—	" "	Terramycin	N.D.	
715	39.3	2920	M.	" "	—	" "	Vitamin K.	L	
794	38.1	2890	M.	" "	Pulmonary T B	" "	Vitamin K. + L.M.I. Blood Penicillin	N.D.	
							Streptomycin		
							Vitamin K.		
							Thyroid		
881	45.5	2960	M.	" "	I P. Eclampsia	M R + M F	Vitamin K. Penicillin	N.D.	
937	39	2670	F.	" "	—	Spont.	Streptomycin	L	
1108	39.5	2850	M.	Intestinal Obstruction	—	" "	Vitamin K.	L	Transferred to Q V H
1178	33.4	2100	F.	Jaundice	—	" "	—	N.D.	2nd Twin
1255	40.1	2920	F.	Diarrhoea	—	" "	Vitamin K. Penicillin	L	
1320	38.4	3118	M.	Jaundice	—	" "	Terramycin Chloromycetin Cortisone	L	
1439	39	3203	F.	" "	—	" "	Vitamin K.	L	
1648	38	2950	M.	" "	—	" "	" "	L	
							Vitamin K. Chloromycetin Terramycin		
1719	40.6	2750	M.	Jaundice	—	L.S.C.S	Thyroid	N.D.	
1722	43.2	3100	M.	Diarrhoea	—	" "	Vitamin K. Chloromycetin Penicillin Terramycin	L	

OTHER NEO-NATAL DISEASE, (Continued)

Reg. No.	Maturity	Birth Weight gm.	Sex	Type of Disease	Maternal Complication	Method of Delivery	Treatment	Result	Remarks
2075	27	3090	M.	Jaundice	Terramycin	M.R. + M.F.	Penicillin Cortisone Vitamin K.	N.D.	
2355	39 6	2020	M.	"	"	Spont.	Vitamin K. Penicillin	N.D.	
2542	37-1	3300	M.	"	"	"	Vitamin K. Penicillin	N.D.	1st Twin
2542	37-1	3050	M.	"	"	M.R. + M.F.	Terramycin Vitamin K. Penicillin	N.D.	2nd Twin
2706	31-2	1250	F.	"	"	Spont.	"	N.D.	
2880	39-3	3350	M.	"	"	"	Penicillin Cortisone Vitamin K.	L	
2927	42-2	2790	F.	"	"	"	Vitamin K. I.M. Blood Cortisone	N.D.	
2963	40 1	3350	F.	"	"	"	Penicillin	L	
2982	39-3	3710	F.	"	"	"	Vitamin K. Penicillin	L	
3073	37	3390	M.	"	"	"	Terramycin Vitamin K.	L	
3130	43-3	3540	M.	"	"	"	"	L	
3161	40-1	2600	M.	"	"	"	Penicillin Vitamin K.	N.D.	
3260	38 4	3190	F.	"	"	"	Cortisone	L	
3317	38 5	1970	M.	"	"	"	Penicillin	L	
3460	41-4	3580	M.	"	"	"	Vitamin K.	L	
3510	"	2920	F.	"	"	"	"	L	
3553	38-3	2930	F.	"	"	"	"	L	
3905	36-5	1980	M.	"	"	"	Vitamin K. Penicillin	N.D.	2nd Twin
4163	36	2720	M.	Intestinal Obstruction	"	L.S.C.S.	Chloromycetin Vitamin K. H.I. Fluid Terramycin	L	Transferred to Q.M.H.
4669	41 4	3160	M.	Jaundice	"	Spont.	Myriton Vitamin C. Vitamin K. Chloromycetin Icterycin	L	

OTHER NEO-NATAL DISEASE, (Continued)

Reg. No.	Maturity	Birth Weight gm.	Sex	Type of Disease	Maternal Complication	Method of Delivery	Treatment	Result	Remarks
4705	37	2350	F.	Respiratory Obstruction	—	L.S.C.S.	Vitamin K. Terramycin Penicillin	L.	Transferred to Q.M.H.
5321	38	3280	M.	jaundice	—	Spont.	Chloromycetin Terramycin Vitamin K. Blood I.M.	L.	
5429	39-3	3150	F.	Intestinal Obstruction	—	L.S.C.S.	Cortisone Thalazole Chloromycetin Terramycin	L.	Transferred to Q.M.H.
5713	39-1	3350	M.	Meconium Ileus	—	Spont.	Terramycin Achromycin	L.	
5743	31-2	2948	M.	Diarrhoea	—	"	Chloromycetin	L.	
5981	41-5	2510	M.	jaundice	—	"	Chloromycetin Cortisone	L.	
6523	40	3250	F.	Diarrhoea	Pulmonary T.B.	"	Vitamin K	L.	
7215	40-1	3590	M.	jaundice	—	"	Penicillin	L.	
7339	35-1	2540	M.	"	—	"	Vitamin K. Terramycin	L.	
							Achromycin		
7574	37-5	3720	M.	"	—	"	Vitamin K. Blood Transfusion Cortisone	L.	
8107	39	3300	F.	Cloudiness of Rt. Cornea	—	"	Terramycin Vitamin K.	L.	Referred to Eye Clinic
NON-BOOKED									
1932	37	2620	M.	Jaundice	—	Brach. Spont.	Vitamin K.	N.D.	
2316	36-3	3062	M.	"	—	M.R. + M.F.	"	N.D.	
2610	32-4	1840	F.	"	Placenta Previa Type II	L.S.C.S.	Vitamin K Penicillin	N.D.	
2983	40	2600	M	Anaemia	P.E. Heart Failure	Spont.	Cortisone Blood Transfusion Penicillin Streptomycin	L.	Transferred to Q.M.H

HYSTERECTOMY

There were 5 hysterectomies, 4 Booked and one Non Booked All the mothers survived

Reg. No	Age	Previous Pregnancies		Indication	Operation	Results	
		Before 28 wks	After 28 wks			M	C
BOOKED							
4321	37	---	4	Ruptured Uterus (Lower Segment Rt. I. t.)	"	L	S B
5374	36	---	2	Ruptured Uterus (Posterior Wall & Upper Segment)	"	L	S B
6290	34	---	7	Ruptured Uterus (Lower Segment, Rt. I t.)	"	L	S B
6435	25	---	2	Placenta Praevia	"	L	N D
NON BOOKED							
1453	38	---	6	Ruptured Uterus (Lower Segment Rt. Lat)	Subtotal	L	S B

X-RAY DEPARTMENT STATISTICS

Type of Cases done in New Tsan Yuk Hospital in 1956.

Month	Chest		Bones		Petemetry		Abdomen		Cardiography					
	I.P.*	O.P.*	Total	I.P.	O.P.	Total	I.P.	O.P.	Total	I.P.	O.P.	Total		
January	13	480	493	12	3	15	8	99	107	13	29	42	1	1
February	21	359	380	6	2	8	6	113	119	5	26	31	3	3
March	21	717	738	2	5	7	12	97	109	20	33	53	—	—
April	39	855	894	2	2	4	10	119	129	20	25	45	—	—
May	55	844	899	—	6	6	16	128	144	19	32	51	2	2
June	28	989	1017	7	4	11	13	147	160	13	60	73	8	8
July	30	1138	1168	4	12	16	9	137	146	15	40	55	1	1
August	41	1000	1041	7	3	10	31	196	227	15	43	58	—	—
September	28	1087	1115	4	2	6	24	186	210	16	19	35	1	1
October	28	976	1004	5	2	7	31	156	187	20	23	43	—	—
November	28	1052	1080	5	2	7	48	139	187	9	20	29	—	—
December	56	793	849	3	2	5	61	97	158	17	12	29	1	4
Total	388	10290	10678	61	45	106	269	1614	1883	182	362	544	6	20

Month	Cystogram		I.I.P.		Cholecystography		Direct Placentography		Ba Meal			
	I.P.	O.P.	Total	I.P.	O.P.	Total	I.P.	O.P.	Total	I.P.	O.P.	Total
January	—	—	—	—	—	—	—	—	—	—	—	—
February	—	—	—	2	—	2	—	—	—	—	—	—
March	—	—	—	—	—	—	1	—	—	—	—	—
April	—	—	—	—	—	—	—	—	—	—	—	—
May	1	—	1	—	—	—	—	—	—	—	—	—
June	3	—	3	1	—	1	—	—	—	—	—	—
July	—	—	—	—	—	—	—	—	—	—	—	—
August	1	—	1	—	—	—	—	—	—	—	—	—
September	1	—	1	—	—	—	—	—	—	—	—	—
October	—	—	—	—	—	—	—	—	—	—	—	—
November	—	—	—	—	—	—	—	—	—	—	—	—
December	—	—	—	—	1	1	—	—	—	—	—	—
Total	6	—	6	6	1	7	2	50	7	1	—	1

* I.P. In-patient.

* O.P. Out-patient.

